

A Statistical Dynamical Model to Predict Extreme Events and Anomalous Features in Shallow Water Waves with Abrupt Depth Change

Andrew J. Majda^{a,1}, M. N. J. Moore^b, and Di Qi^{a,1}

^aDepartment of Mathematics and Center for Atmosphere and Ocean Science, Courant Institute of Mathematical Sciences, New York University, New York, NY 10012;

^bDepartment of Mathematics and Geophysical Fluid Dynamics Institute, Florida State University, Tallahassee, FL

This manuscript was compiled on November 27, 2018

Understanding and predicting extreme events and their anomalous statistics in complex nonlinear systems is a grand challenge in climate, material, and neuroscience, as well as for engineering design. Recent laboratory experiments in weakly turbulent shallow water reveal a remarkable transition from Gaussian to anomalous behavior as surface waves cross an abrupt depth change (ADC). Downstream of the ADC, PDFs of surface displacement exhibit strong positive skewness, which is accompanied by an elevated level of extreme events. Here we develop a statistical dynamical model to explain and quantitatively predict the above anomalous statistical behavior as experimental control parameters are varied. The first step is to use incoming and outgoing truncated Korteweg-de Vries (TKdV) equations matched in time at the ADC. The TKdV equation is a Hamiltonian system which induces incoming and outgoing statistical Gibbs invariant measures. The statistical matching of the known nearly Gaussian incoming Gibbs state at the ADC completely determines the predicted anomalous outgoing Gibbs state, which can be calculated by a simple sampling algorithm, verified by direct numerical simulations, and successfully captures key features of the experiment. There is even an analytic formula for the anomalous outgoing skewness. The strategy here should be useful for predicting extreme anomalous statistical behavior in other dispersive media in different settings (21, 22).

extreme anomalous event | statistical TKdV model | matching Gibbs measures | surface wave displacement and slope

Understanding and predicting extreme events and their anomalous statistics in complex nonlinear systems is a grand challenge in climate, material and neuroscience as well as for engineering design. This is a very active contemporary topic in applied mathematics with qualitative and quantitative models (1–7) and novel numerical algorithms which overcome the curse of dimensionality for extreme event prediction in large complex systems (2, 8–11). The occurrence of Rogue waves as extreme events within different physical settings of deep water (12–16) and shallow water (17–19) is an important practical topic.

Recent laboratory experiments in weakly turbulent shallow water reveal a remarkable transition from Gaussian to anomalous behavior as surface waves cross an abrupt depth change (ADC). A normally-distributed incoming wave train, upstream of the ADC, transitions to a highly non-Gaussian outgoing wave train, downstream of the ADC, that exhibits large positive skewness of the surface height and more frequent extreme events (20). Here we develop a statistical dynamical model to explain and quantitatively predict this anomalous behavior as experimental control parameters are varied. The first step is to use incoming and outgoing truncated Korteweg-de Vries (TKdV) equations matched in time at the ADC. The TKdV

equation is a Hamiltonian system which induces incoming and outgoing Gibbs invariant measures. The statistical matching of the known nearly Gaussian incoming Gibbs state at the ADC completely determines the predicted anomalous outgoing Gibbs state, which can be calculated by a simple MCMC algorithm, verified by direct numerical simulations, and successfully captures key features of the experiment. There is even an analytic formula for the anomalous outgoing skewness. The strategy here should be useful for predicting extreme anomalous statistical behavior in other dispersive media in different settings (21, 22).

1. Experiments showing anomalous wave statistics induced by an abrupt depth change

Controlled laboratory experiments were carried out in (20) to examine the statistical behavior of surface waves crossing an ADC. In these experiments, nearly unidirectional waves are generated by a paddle wheel and propagate through a long, narrow wave tank. Midway through, the waves encounter a step in the bottom topography, and thus abruptly transition to a shallower depth. The paddle wheel is forced with a pseudo-random signal intended to mimic a Gaussian random sea upstream of the ADC. In particular, the paddle angle is

Significance Statement

Understanding and predicting extreme events and their anomalous statistics in complex nonlinear systems is a grand challenge in applied sciences as well as for engineering design. Recent controlled laboratory experiments in weakly turbulent shallow water with abrupt depth change exhibit a remarkable transition from nearly Gaussian statistics to extreme anomalous statistics with large positive skewness of the surface height. We develop a statistical dynamical model to explain and quantitatively predict the anomalous statistical behavior. The incoming and outgoing waves are modeled by the truncated Korteweg-de Vries equations statistically matched at the depth change. The statistical matching of the known nearly Gaussian incoming Gibbs state completely determines the predicted anomalous outgoing Gibbs state, and successfully captures key features of the experiment.

A.J.M. designed the research. A.J.M., M.N.J.M., and D.Q. performed the research. A.J.M., M.N.J.M., and D.Q. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: qidi@cims.nyu.edu, jonjon@cims.nyu.edu

- 125 specified as 187
 126 $\theta(t) = \theta_0 + \Delta\theta \sum_{n=1}^N a_n \cos(\omega_n t + \delta_n)$, $E \sim (\Delta\theta)^2 \sum_n a_n^2 \omega_n^2$. 188
 127 where the weights a_n are Gaussian in spectral space with 189
 128 peak frequency ω_p and the phases δ_n are uniformly distributed 190
 129 random variables. The peak frequency gives rise to a characteristic wavelength λ_c which can be estimated from the 191
 130 dispersion relation. The energy E injected into the system 192
 131 is determined by the angle amplitude $\Delta\theta$, which is the main 193
 132 control parameter varied in (20). Optical measurements of the 194
 133 free surface reveal a number of surprising statistical features: 195
 134 • Distinct statistics are found between the incoming and 196
 135 outgoing wave disturbances: incoming waves display near- 197
 136 Gaussian statistics, while outgoing waves skew strongly 198
 137 towards positive displacement.
 138 • The degree of non-Gaussianity present in the outgoing 199
 139 waves depends on the injected energy E : larger energies 200
 140 generate stronger skewness in the surface displacement 201
 141 PDFs and more extreme events. 202
 142 • Compared to the incoming wave train, the power spectrum 203
 143 of the outgoing wave field decays more slowly, which 204
 144 indicates that the anomalous behavior is associated with 205
 145 an elevated level of high frequencies. 206
 146
 147 **2. Surface wave turbulence modeled by truncated KdV 207**
 148 **equation with depth dependence** 208
 149
 150 The Korteweg-de Vries (KdV) equation is a one-dimensional, 209
 151 deterministic model capable of describing (weak?) surface 210
 152 wave turbulence. More specifically, KdV is leading-order 211
 153 approximation for surface waves governed by a balance of 212
 154 nonlinear and dispersive effects, valid in an appropriate far- 213
 155 field limit (23). Moreover, KdV has been adapted to describe 214
 156 waves propagating over variable depth (23). Here, we consider 215
 157 the variable-depth KdV equation truncated at wavenumber 216
 158 Λ (with $J = 2\Lambda + 1$ grid points) in order to generate weakly 217
 159 turbulent dynamics. The surface displacement is described 218
 160 by the state variable $u_\Lambda^\pm(x, t)$ with superscript ‘-’ for the 219
 161 incoming waves and ‘+’ for the outgoing waves. The Galerkin 220
 162 truncated variable $u_\Lambda = \sum_{1 \leq |k| \leq \Lambda} \hat{u}_k(t) e^{ikx}$ is normalized 221
 163 with zero mean $\hat{u}_0 = 0$ and unit energy $2\pi \sum_{k=1}^\Lambda |\hat{u}_k|^2 = 1$, 222
 164 which are conserved quantities. Here, $u_\Lambda \equiv \mathcal{P}_\Lambda u$ denotes the 223
 165 subspace projection. The evolution of u_Λ^\pm is governed by the 224
 166 truncated KdV equation with depth change D_\pm 225
 167
$$\frac{\partial u_\Lambda^\pm}{\partial t} + \frac{D_\pm^{-3/2}}{2} E_0^{1/2} L_0^{-3/2} \frac{\partial}{\partial x} \mathcal{P}_\Lambda (u_\Lambda^\pm)^2 + D_\pm^{1/2} L_0^{-3} \frac{\partial^3 u_\Lambda^\pm}{\partial x^3} = 0. \quad [1]$$
 226
 168 Equation [1] is non-dimensionalized on the periodic domain $x \in 227$
 169 $[-\pi, \pi]$. The depth is assumed to be unit $D_- = 1$ before the 228
 170 ADC and $D_+ < 1$ after the ADC. The conserved Hamiltonian 229
 171 can be decomposed as 230
 172
$$\mathcal{H}_\Lambda^\pm = D_\pm^{-3/2} E_0^{1/2} L_0^{-3/2} H_3(u_\Lambda^\pm) - D_\pm^{1/2} L_0^{-3} H_2(u_\Lambda^\pm),$$
 231
 173
$$H_3(u) = \frac{1}{6} \int_{-\pi}^{\pi} u^3 dx, \quad H_2(u) = \frac{1}{2} \int_{-\pi}^{\pi} \left(\frac{\partial u}{\partial x} \right)^2 dx.$$
 232
 174 where we refer to H_3 as the cubic term and H_2 the quadratic 233
 175 term. We introduce parameters (E_0, L_0, Λ) based on the 234
 176 following assumptions:
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248

249 change. The reference values for the model scales can be esti-
 250 mated in the range $L_0 \in [2, 6]$ and $E_0 \in [50, 200]$. These are
 251 the values we will test in the direct numerical simulations. See
 252 details about the derivation from scale analysis in *SI Appendix*,
 253 A.

254

255 3. Equilibrium statistical mechanics for generating the 256 stationary invariant measure

257 Since the TKdV equation satisfies the Liouville property, the
 258 equilibrium invariant measure can be described by an equi-
 259 librium statistical formulism (24–26) using a Gibbs measure
 260 with the conserved energy \mathcal{E}_Λ and Hamiltonian \mathcal{H}_Λ . The equi-
 261 librium invariant measure is dictated by the conservation laws
 262 in the TKdV equation. In the case with fixed total energy E_0 ,
 263 this is the *mixed Gibbs measure* in the truncated model with
 264 microcanonical energy and canonical Hamiltonian ensembles
 265 (24)

266
$$\mathcal{G}_\theta^\pm(u_\Lambda^\pm; E_0) = C_\theta^\pm \exp(-\theta^\pm \mathcal{H}(u_\Lambda^\pm)) \delta(\mathcal{E}(u_\Lambda^\pm) - E_0), \quad [3]$$

267 with θ representing the “inverse temperature”. The distinct
 268 statistics in the upstream and downstream waves can be con-
 269 trolled by the value of θ . Negative temperature, $\theta^\pm < 0$, is
 270 the appropriate regime to predict the experiments as shown
 271 below. In the incoming flow field, the inverse temperature θ^-
 272 is chosen so that \mathcal{G}_θ^- has Gaussian statistics. Using the above
 273 invariant measures [3], the expectation of any functional $F(u)$
 274 can be computed based on the Gibbs measure

275
$$\langle F \rangle_{\mathcal{G}_\theta} \equiv \int F(u) \mathcal{G}_\theta(u) du.$$

276 The value of θ in the invariant measure is specified from $\langle H_\Lambda \rangle_{\mathcal{G}_\theta}$
 277 (24, 26). The invariant measure also predicts an equilibrium
 278 energy spectrum without running the TKdV equation directly.
 279 On the other hand, the time autocorrelation and transient
 280 statistics about the state variable u_Λ cannot be recovered from
 281 the statistical theory.

282 **Statistical matching condition in invariant measures before
 283 and after the abrupt depth change.** The Gibbs measures \mathcal{G}_θ^\pm
 284 are defined based on the different inverse temperatures θ^\pm on
 285 the two sides of the solutions

286
$$\begin{aligned} \mu_t^-(u_\Lambda^-; D_-), \quad u_\Lambda |_{t=T_{ADC}-} = u_0, \quad t < T_{ADC}; \\ \mu_t^+(u_\Lambda^+; D_+), \quad u_\Lambda |_{t=T_{ADC}+} = u_0, \quad t > T_{ADC}, \end{aligned}$$

287 where u_0 represents the deterministic matching condition be-
 288 tween the incoming and outgoing waves. The two distributions,
 289 μ_t^-, μ_t^+ should also be matched at the depth change location
 290 T_{ADC} , so that,

291
$$\mu_{t=T_{ADC}}^-(u_\Lambda) = \mu_{t=T_{ADC}}^+(u_\Lambda).$$

292 In matching the flow statistics before and after the abrupt
 293 depth change, we first use the conservation of the determinis-
 294 tic Hamiltonian H_Λ^+ after the depth change. Then assuming
 295 ergodicity (24, 25), the statistical expectation for the Hamil-
 296 tonian $\langle H_\Lambda^+ \rangle$ is conserved in time after the depth change at
 297 $t = T_{ADC}$ and should stay in the same value as the system ap-
 298 proaches equilibrium as $t \rightarrow \infty$. The final statistical matching
 299 condition to get the outgoing flow statistics with parameter
 300 θ^+ can be found by

301
$$\langle H_\Lambda^+ \rangle_{\mathcal{G}_\theta^+} = \langle H_\Lambda^+ \rangle_{\mathcal{G}_\theta^-}, \quad [4]$$

302 with the outgoing flow Hamiltonian H_Λ^+ and the Gibbs mea-
 303 sures \mathcal{G}_θ^\pm before and after the abrupt depth change.

304 4. The nearly Gaussian incoming statistical state

305 The incoming flow is always characterized by a near-Gaussian
 306 distribution in the wave displacement. It is found that a
 307 physically consistent Gibbs measure should take negative val-
 308 ues in the inverse temperature parameter $\theta < 0$, where a
 309 proper distribution function and a decaying energy spectrum
 310 are generated (see (26) and *SI Appendix*, B.1 for the explicit
 311 simulation results). The upstream Gibbs measure \mathcal{G}_θ^- with
 312 $D_- = 1$ displays a wide parameter regime in (θ^-, E_0) with
 313 near-Gaussian statistics. In the left panel of Figure 1 (a), the
 314 inflow skewness κ_3^- varies only slightly with changing values of
 315 E_0 and θ^- . The incoming flow PDF then can be determined
 316 by picking the proper parameter value θ^- in the near Gaus-
 317 sian regime with small skewness. In contrast, the downstream
 318 Gibbs measure \mathcal{G}_θ^+ with $D_+ = 0.24$ shown in the right panel
 319 of Figure 1 (a) generates much larger skewness κ_3^+ as the
 320 absolute value of θ^+ and the total energy level E_0 increases.
 321 The solid lines in Figure 1 (c) offer a further confirmation of
 322 the transition from near-Gaussian statistics with small κ_3^- to
 323 a strongly skewed distribution κ_3^+ after the depth change.

324 In the next step, the value of the downstream θ^+ is deter-
 325 mined based on the matching condition [4]. The expectation
 326 $\langle H_\Lambda^+ \rangle_{\mathcal{G}_\theta^-}$ about the incoming flow Gibbs measure can be cal-
 327 culated according to the predetermined parameter values of
 328 θ^- as well as E_0 from the previous step. For the direct nu-
 329 matical experiments shown later in Figure 2, we pick proper
 330 choices of test parameter values as $L_0 = 6, E_0 = 100$ and
 331 $\theta^- = -0.1, -0.3, -0.5$. More test cases with different system
 332 energy E_0 can be found in *SI Appendix*, B.2 where similar
 333 transition from near Gaussian symmetric PDFs to skewed
 334 PDFs in the flow state u_Λ^\pm can always be observed.

335 **Direct numerical model simulations.** Besides the prediction of
 336 equilibrium statistical measures from the equilibrium statisti-
 337 cal approach, another way to predict the downstream model
 338 statistics is through running the dynamical model [1] directly.
 339 The TKdV equation is found to be ergodic with proper mixing
 340 property as measured by the decay of autocorrelations as long
 341 as the system starts from a negative inverse temperature state
 342 as described before. For direct numerical simulations of the
 343 TKdV equations, a proper symplectic integrator is required to
 344 guarantee the Hamiltonian and energy are conserved in time.
 345 It is crucial to use the symplectic scheme to guarantee the
 346 exact conservation of the energy and Hamiltonian since they
 347 are playing the central role in generating the invariant measure
 348 and the statistical matching. The symplectic schemes used
 349 here for the time integration of the equation is the 4th-order
 350 midpoint method (27). Details about the mixing properties
 351 from different initial states and the numerical algorithm are
 352 described in *SI Appendix*, C.

353 5. Predicting extreme anomalous behavior after the 354 ADC by statistical matching

355 With the inflow statistics well described and the numerical
 356 scheme set up, we are able to predict the downstream anomalous
 357 statistics starting from the near-Gaussian incoming flow
 358 going through the abrupt depth change from $D_- = 1$ to
 359 $D_+ = 0.24$.




Fig. 1. First row: skewness from the Gibbs measures in incoming and outgoing flow states with different values of total energy E_0 and inverse temperature θ (notice the different scales in the incoming and outgoing flows); Second row: outgoing flow parameter θ^+ as a function of the incoming flow θ^- computed from the statistical matching condition with three energy level E_0 ; Last row: skewness in the outgoing flow with the matched value of θ^+ as a function of the inflow parameter θ^- (the theoretical predictions using [5] are compared).

$D_+ = 0.24$. First, we consider the statistical prediction in the downstream equilibrium measure directly from the matching condition. The downstream parameter value θ^+ is determined by solving the nonlinear equation [4] as a function of θ^+ , $F(\theta^+) = \langle H_\Lambda^+ \rangle_{\mathcal{G}_\theta^+}(\theta^+) - \langle H_\Lambda^+ \rangle_{\mathcal{G}_\theta^-} = 0$. In the numerical approach, we adopt a modified secant method avoiding the stiffness in the parameter regime (see the *SI Appendix, B.2* for the algorithm). The fitted solution is plotted in Figure 1 (b) as a function of the proposed inflow θ^- . A nonlinear θ^- - θ^+ relation is discovered from the matching condition. The downstream inverse temperature θ^+ will finally saturate at some level. The corresponding downstream skewness of the wave displacement u_Λ predicted from the statistical matching of Gibbs measures is plotted in Figure 1 (c). In general, a large positive skewness for outgoing flow κ_3^+ is predicted from the theory, while the incoming flow skewness κ_3^- is kept in a small value in a wide range of θ^- . Note that with $\theta^- \sim 0$ (that is, using the microcanonical ensemble only with energy conservation), the outflow statistics are also near Gaussian with weak skewness. The skewness in the outflow statistics grows as the inflow parameter value θ^- increases in amplitude.

For a second approach, we can use direct numerical simulations starting from the initial state sampled from the incoming flow Gibbs measure \mathcal{G}_θ^- and check the transient changes in the

model statistics. Figure 2 illustrates the change of statistics as the flow goes through the abrupt depth change. The first row plots the changes in the skewness and kurtosis for the state variable u_Λ after the depth change at $t = 0$. The PDFs in the incoming and outgoing flow states are compared with three different initial inverse temperatures θ^- . After the depth changes to $D_0 = 0.24$ abruptly at $t = 0$, both the skewness and kurtosis jump to a much larger value in a short time, implying the rapid transition to a highly skewed non-Gaussian statistical regime after the depth change. Further from Figure 2, different initial skewness (but all relatively small) is set due to the various values of θ^- . With small $\theta^- = -0.1$, the change in the skewness is not very obvious (see the second row of Figure 2 for the incoming and outgoing PDFs of u_Λ). In comparison, if the incoming flow starts from the initial parameter $\theta^- = -0.3$ and $\theta^- = -0.5$, much larger increase in the skewness is induced from the abrupt depth change. Furthermore, in the detailed plots in the third row of Figure 2 for the downstream PDFs under logarithmic scale, fat tails towards the positive direction can be observed, which represent the extreme events in the downstream flow (see also Figure 3 for the time-series of u_Λ).

As a result, the downstream statistics in final equilibrium predicted from the direct numerical simulations here agree with the equilibrium statistical mechanics prediction illustrated in Figure 1. The prediction from these two different approaches confirm each other.

6. Analytic formula for the upstream skewness after the ADC

A statistical link between the upstream and downstream energy spectra can be found for an analytical prediction of the skewness in the flow state u after the ADC. The skewness of the state variable u_j at one spatial grid point is defined as the ratio between the third and second moments

$$\kappa_3 = \langle u_j^3 \rangle_\mu / \langle u_j^2 \rangle_\mu^{\frac{3}{2}}.$$

Now we introduce mild assumptions on the distribution functions:

- The upstream equilibrium measure μ_- has a relatively small skewness so that

$$\langle H_3 \rangle_{\mu_-} = \frac{1}{6} \int_{-\pi}^{\pi} \langle u^3 \rangle_{\mu_-} dx \equiv \epsilon;$$

- The downstream equilibrium measure μ_+ is homogeneous at each physical grid point, so that the second and third moments are invariant at each grid point

$$\langle u_j^2 \rangle_{\mu_+} = \sigma^2 = \pi^{-1}, \quad \langle u_j^3 \rangle_{\mu_+} = \sigma^3 \kappa_3 = \pi^{-\frac{3}{2}} \kappa_3^+.$$

Then the skewness of the downstream state variable u_Λ^+ after the ADC is given by the difference between the inflow and outflow wave slope energy of u_x

$$\kappa_3^+ = \frac{3}{2} \pi^{\frac{1}{2}} L_0^{-\frac{3}{2}} E_0^{-\frac{1}{2}} D_+^2 \int_{-\pi}^{\pi} \left[\langle u_x^2 \rangle_{\mu_+} - \langle u_x^2 \rangle_{\mu_-} \right] dx + 3\pi^{\frac{1}{2}} \epsilon.$$

The detailed derivation is shown in *SI Appendix, B.2*. In particular, the downstream skewness with near-Gaussian inflow

497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519




Fig. 2. Changes in the statistics of the flow state going through the abrupt depth change. The initial ensemble is set with the incoming flow Gibbs measure with different inverse temperature θ^- . First row: time evolution of the skewness and kurtosis. The abrupt depth change is taking place at $t = 0$; Second row: inflow and outflow PDFs of u_Λ ; Third row: the downstream PDFs fitted with Gamma distributions with consistent variance and skewness (in log coordinate in y); Last row: energy spectra in the incoming and outgoing flows.

523

524

525 statistics $\epsilon \ll 1$ is positive if and only if the difference of the
526 incoming and outgoing wave slope energy is positive. This
527 means that there is more small scale wave slope energy in the
528 outgoing state. As an evidence, in the last row of Figure 2
529 in all the weak and strong skewness cases, the outflow energy
530 spectrum always has a slower decay rate than the inflow energy
531 spectrum which possesses stronger energy in larger scales and
532 weaker energy in the smaller scales.

533 In Figure 1 (c), we compare the accuracy of the theoretical
534 estimation [5] with numerical tests. In the regime with small
535 incoming inverse temperature θ^- , the theoretical formula offers
536 a quite accurate approximation of the third-order skewness
537 using only information from the second-order moments of the
538 wave-slope spectrum.

539

540 7. Key features from experiments captured by the sta- 541 tistical dynamical model

542 In this final section, we emphasize the crucial features generated
543 by the statistical dynamical model [1] by making comparison
544 with the experimental observations in (20). As from the scale analysis displayed in Section 2, the theory is set in
545 the same parameter regime as the experimental setup.

547

- 548 • The transition from near-Gaussian to skewed non-
549 Gaussian distribution as well as the jump in both skewness
550 and kurtosis observed in the experiment observations (Fig.
551 1 of (20)) can be characterized by the statistical model
552 simulation results (see the first and second row of Figure
553 2). Notice that the difference in the decay of third and
554 fourth moments in the far end of the downstream regime
555 from the experimental data is due to the dissipation effect
556 in the flow from the wave absorbers that is not mod-
557 eled in the statistical model here. The model simulation
558 time-series plotted in Figure 3 can be compared with

559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620

the observed time sequences from experiments (Fig. 1 of (20)). The downstream simulation generates waves with strong and frequent intermittency towards the positive displacement, while the upstream waves show symmetric displacements in two directions with at most small peaks in slow time. Even in the time-series at a single location $x = 0$, the long-time variation displays similar structures.

- The downstream PDFs in experimental data are estimated with a Gamma distribution in Fig. 2 of (20). Here in the same way, we can fit the highly skewed outgoing flow PDFs from the numerical results with the Gamma distribution

$$\rho(u; k, \alpha) = \frac{e^{-k} \alpha^{-1}}{\Gamma(k)} (k + \alpha^{-1} u)^{k-1} e^{-\alpha^{-1} u}.$$

The parameters (k, α) in the Gamma distribution are fitted according to the measured statistics in skewness and variance, that is, $\sigma^2 = k\alpha^2$, $\kappa_3 = 2/\sqrt{k}$. And the excess kurtosis of the Gamma distribution can be recovered as $\kappa_4 = 6/k$. As shown in the third row of Figure 2, excellent agreement in the PDFs with the Gamma distributions is reached in consistency with the experimental data observations. The accuracy with this approximation increases as the initial inverse temperature θ^- increases in value to generate more skewed distribution functions.

- Experimental measurements of the power spectra (Fig. 4 of (20)) reveal the downstream measurements to contain more energy at small scales, i.e. a relatively slower decay rate of the spectrum. This result is also observed in the direct numerical simulations here (detailed results shown in *SI Appendix, C.2*), as the outgoing state contains more energetic high frequencies.




Fig. 3. Realization of the downstream and upstream flow solutions u_A^\pm . Note the larger vertical scale in the downstream time-series plot.

8. Concluding discussion

We have developed a statistical dynamical model to explain and predict extreme events and anomalous features of shallow water waves crossing an abrupt depth change. The theory is based on the dynamical modeling strategy consisting of the TKdV equation matched at the abrupt depth change with conservation of energy and Hamiltonian. Predictions can be made of the extreme events and anomalous features by matching incoming and outgoing statistical Gibbs measures before and after the abrupt depth transition. The statistical matching of the known nearly Gaussian incoming Gibbs state completely

determines the predicted anomalous outgoing Gibbs state, which can be calculated by a simple sampling algorithm, verified by direct numerical simulations, and successfully captures key features of the experiment. An analytic formula for the anomalous outgoing skewness is also derived. The strategy here should be useful for predicting extreme statistical events in other dispersive media in different settings.

ACKNOWLEDGMENTS. This research of A. J. M. is partially supported by the Office of Naval Research through MURI N00014-16-1-2161. D. Q. is supported as a postdoctoral fellow on the second grant. M.N.J.M would like to acknowledge support from Simons grant 524259.

1. Majda AJ, Branicki M (2012) Lessons in uncertainty quantification for turbulent dynamical systems. *Discrete & Continuous Dynamical Systems-A* 32(9):3133–3221.
2. Mohamad MA, Sapsis TP (2018) A sequential sampling strategy for extreme event statistics in nonlinear dynamical systems. *Proceedings of the National Academy of Sciences* 115(44):11138–11143.
3. Qi D, Majda AJ (2016) Predicting fat-tailed intermittent probability distributions in passive scalar turbulence with imperfect models through empirical information theory. *Communications in Mathematical Sciences* 14(6):1687–1722.
4. Majda AJ, Tong XT (2015) Intermittency in turbulent diffusion models with a mean gradient. *Nonlinearity* 28(11):4171–4208.
5. Majda AJ, Chen N (2018) Model error, information barriers, state estimation and prediction in complex multiscale systems. *Entropy* 20(9):644.
6. Majda AJ, Tong XT (2018) Simple nonlinear models with rigorous extreme events and heavy tails. *arXiv preprint arXiv:1805.05615*.
7. Thual S, Majda AJ, Chen N, Stechmann SN (2016) Simple stochastic model for el nino with westerly wind bursts. *Proceedings of the National Academy of Sciences* 113(37):10245–10250.
8. Chen N, Majda AJ (2018) Efficient statistically accurate algorithms for the Fokker-Planck equation in large dimensions. *Journal of Computational Physics* 354:242–268.
9. Chen N, Majda AJ (2017) Beating the curse of dimension with accurate statistics for the Fokker-Planck equation in complex turbulent systems. *Proceedings of the National Academy of Sciences* 114(49):12864–12869.
10. Chen N, Majda AJ (2018) Conditional Gaussian systems for multiscale nonlinear stochastic systems: Prediction, state estimation and uncertainty quantification. *Entropy* 20(7):509.
11. Qi D, Majda AJ (2018) Predicting extreme events for passive scalar turbulence in two-layer baroclinic flows through reduced-order stochastic models. *Communications in Mathematical Sciences* 16(1):17–51.
12. Adcock TA, Taylor PH (2014) The physics of anomalous ('rogue') ocean waves. *Reports on Progress in Physics* 77(10):105901.
13. Cousins W, Sapsis TP (2015) Unsteady evolution of localized unidirectional deep-water wave groups. *Physical Review E* 91(6):063204.
14. Farazmand M, Sapsis TP (2017) Reduced-order prediction of rogue waves in two-dimensional deep-water waves. *Journal of Computational Physics* 340:418–434.
15. Onorato M, Osborne AR, Serio M, Bertone S (2001) Freak waves in random oceanic sea states. *Physical Review Letters* 86(25):5831–5834.
16. Demattei G, Grafke T, Vanden-Eijnden E (2018) Rogue waves and large deviations in deep sea. *Proceedings of the National Academy of Sciences* 115(5):855–860.
17. Sergeeva A, Pelinovsky E, Talipova T (2011) Nonlinear random wave field in shallow water: variable Korteweg-de Vries framework. *Natural Hazards and Earth System Sciences* 11(2):323–330.
18. Trulsen K, Zeng H, Gramstad O (2012) Laboratory evidence of freak waves provoked by non-uniform bathymetry. *Physics of Fluids* 24(9):097101.
19. Viotti C, Dias F (2014) Extreme waves induced by strong depth transitions: Fully nonlinear results. *Physics of Fluids* 26(5):051705.
20. Bolles CT, Speer K, Moore MNJ (2018) Anomalous wave statistics induced by abrupt depth change. *arXiv preprint arXiv:1808.07958*.
21. Solli D, Ropers C, Koopath P, Jalali B (2007) Optical rogue waves. *Nature* 450(7172):1054.
22. Höhmann R, Kuhl U, Stöckmann HJ, Kaplan L, Heller E (2010) Freak waves in the linear regime: A microwave study. *Physical review letters* 104(9):093901.
23. Johnson RS (1997) *A modern introduction to the mathematical theory of water waves*. (Cambridge university press) Vol. 19.
24. Abramov RV, Kovačič G, Majda AJ (2003) Hamiltonian structure and statistically relevant conserved quantities for the truncated Burgers-Hopf equation. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences* 56(1):1–46.
25. Majda A, Wang X (2006) *Nonlinear dynamics and statistical theories for basic geophysical flows*. (Cambridge University Press).
26. Bajars J, Frank J, Leimkuhler B (2013) Weakly coupled heat bath models for Gibbs-like invariant states in nonlinear wave equations. *Nonlinearity* 26(7):1945–1973.
27. McLachlan R (1993) Symplectic integration of hamiltonian wave equations. *Numerische Mathematik* 66(1):465–492.