Data-Driven Models for Discrete Hedging Problem

Ke Nian Supervisors: Prof. Yuying Li and Prof. Thomas. F. Coleman

David R. Cheriton School of Computer Science, University of Waterloo,

Waterloo, Canada

August 19, 2019

The Dupire formula enables us to deduce the volatility function in a local volatility model from quoted put and call options in the market.

▶ Under a risk-neutral measure, we assume

$$\frac{dS_t}{S_t} = r(t)dt + \sigma(t, S_t)dZ_t$$

▶ The forward price for delivery at time T:

$$F_t = F(t, T) = S_t e^{\int_t^T r(s)ds}$$

► We also have:

$$\frac{dF_t}{F_t} = \widetilde{\sigma}(t, F_t) dZ_t$$

$$\widetilde{\sigma}(t, F_t) = \sigma(t, F_t e^{-\int_t^T r(s)ds})$$

Introduction

- ▶ The spot price at t is: $C(T,K)e^{-\int_t^T r(s)ds}$.
- It can be shown that, with $\theta(T,K)$ be the normal density function of S_T :

$$C(T,K) = \int_{K}^{\infty} (x - K)\theta(T, x)dx$$

Breedon-Litzenberger Formulas

logo.png

Data-Driven Models for Discrete Hedging Problem

Ke Nian

If we differentiate pricing function ${\cal C}(T,K)$ twice we obtain:

$$\frac{\partial C(T,K)}{\partial K} = \Theta(T,K) - 1$$

$$\frac{\partial^2 C(T,K)}{\partial K^2} = \theta(T,K)$$

Forward Equation

logo.png

Data-Driven Models for Discrete Hedging Problem Ke Nian

Introduction

Let h be an arbitrary function and $v(t,x)={\cal E}[h({\cal F}_T)|{\cal F}_t=x],$ we can show that:

$$E[h(F_T)] = \int_0^\infty v(t, x)\theta(t, x)dx$$

Differentiate them with regards to t:

$$0 = \int_0^\infty \frac{\partial v(t, x)}{\partial t} \theta(t, x) dx + \int_0^\infty \frac{\partial \theta(t, x)}{\partial t} v(t, x) dx$$

Backward Equation

logo.png

Data-Driven Models for Discrete Hedging Problem Ke Nian

Introduction

Similar to the deduction of Black-Scholes equation, using ito's lemma, we can see that.

$$\frac{\partial v(t,x)}{\partial t} + \frac{1}{2}\widetilde{\sigma}^2(t,x)x^2 \frac{\partial^2 v(t,x)}{\partial x^2} = 0$$

Note that we are dealing with forward price, the term with interest rate is dropped.

Link Backward Equation and Forward Equation

logo.png

Data-Driven Models for Discrete Hedging Problem

Introduction

Plug in the backward equation:

$$\frac{\partial v(t,x)}{\partial t} + \frac{1}{2}\widetilde{\sigma}^2(t,x)x^2 \frac{\partial^2 v(t,x)}{\partial x^2} = 0$$

into forward equation:

$$0 = \int_0^\infty \frac{\partial v(t, x)}{\partial t} \theta(t, x) dx + \int_0^\infty \frac{\partial \theta(t, x)}{\partial t} v(t, x) dx$$

We have:

$$0 = -\int_0^\infty \frac{1}{2} \widetilde{\sigma}^2(t,x) x^2 \frac{\partial^2 v(t,x)}{\partial x^2} \theta(t,x) dx + \int_0^\infty \frac{\partial \theta(t,x)}{\partial t} v(t,x) dx$$

Using the rule of Integration By Parts twice:

$$\begin{split} &\int_{0}^{\infty} \widetilde{\sigma}^{2}(t,x)x^{2} \frac{\partial^{2}v(t,x)}{\partial x^{2}} \theta(t,x) dx \\ =& \widetilde{\sigma}^{2}(t,x)x^{2} \frac{\partial v(t,x)}{\partial x} \theta(t,x) - \int_{0}^{\infty} \frac{\partial v(t,x)}{\partial x} \frac{\partial [\widetilde{\sigma}^{2}(t,x)x^{2}\theta(t,x)]}{\partial x} dx \\ =& \int_{0}^{\infty} \frac{\partial^{2} [\widetilde{\sigma}^{2}(t,x)x^{2}\theta(t,x)]}{\partial x^{2}} v(t,x) dx \end{split}$$

Integration By Parts:

$$\int uv'dx = uv - \int u'vdx$$

Forward Equation

logo.png

Data-Driven Models for Discrete Hedging Problem Ke Nian

Introduction

Plug in:

$$\int_{0}^{\infty} \widetilde{\sigma}^{2}(t,x)x^{2} \frac{\partial^{2}v(t,x)}{\partial x^{2}} \theta(t,x)dx$$

$$= \int_{0}^{\infty} \frac{\partial^{2}[\widetilde{\sigma}^{2}(t,x)x^{2}\theta(t,x)]}{\partial x^{2}} v(t,x)dx$$

We finally have:

$$\begin{split} 0 &= \int_0^\infty \frac{1}{2} \frac{\partial^2 [\widetilde{\sigma}^2(t,x) x^2 \theta(t,x)]}{\partial x^2} v(t,x) dx - \int_0^\infty \frac{\partial \theta(t,x)}{\partial t} v(t,x) dx \\ &= \int_0^\infty [\frac{1}{2} \frac{\partial^2 [\widetilde{\sigma}^2(t,x) x^2 \theta(t,x)]}{\partial x^2} - \frac{\partial \theta(t,x)}{\partial t}] v(t,x) dx \end{split}$$

Data-Driven Models for Discrete Hedging Problem Ke Nian

Introduction

Note that, it holds for arbitrary $h(F_t)$ and v(t,x):

$$0 = \int_0^\infty \left[\frac{1}{2} \frac{\partial^2 \left[\widetilde{\sigma}^2(t, x) x^2 \theta(t, x) \right]}{\partial x^2} - \frac{\partial \theta(t, x)}{\partial t} \right] v(t, x) dx$$

We then must have:

$$\frac{\partial \theta(t,x)}{\partial t} = \frac{1}{2} \frac{\partial^2 [\widetilde{\sigma}^2(t,x) x^2 \theta(t,x)]}{\partial x^2} \tag{1}$$

Dupire's equation

logo.png

Data-Driven Models for Discrete Hedging Problem Ke Nian

Introduction

$$\frac{\partial C(T,K)}{\partial T} = \int_{K}^{\infty} (x-K) \frac{\partial \theta(T,x)}{\partial T} dx$$

$$= \int_{K}^{\infty} (x-K) \frac{1}{2} \frac{\partial^{2} [\widetilde{\sigma}^{2}(T,x)x^{2}\theta(T,x)]}{\partial x^{2}} dx \quad [eq(1)]$$

$$= -\frac{1}{2} \int_{K}^{\infty} \frac{\partial [\widetilde{\sigma}^{2}(T,x)x^{2}\theta(T,x)]}{\partial x} dx$$

$$= \frac{1}{2} \sigma^{2}(T,K)K^{2}\theta(T,K)$$

$$= \frac{1}{2} \sigma^{2}(T,K)K^{2} \frac{\partial^{2} C(T,K)}{\partial K^{2}}$$

logo.png

Data-Driven Models for Discrete Hedging Problem

Ke Nian

Given a time of grid of expiries $0 = t_0 < t_1 < \dots$ and a a set of volatility functions, we recursively solve the finite difference discretization of the Dupure forward equation:

$$\frac{C(t_{i+1},k) - C(t_i,k)}{t_{i+1} - t_i} = \frac{1}{2}\sigma(t_i,k)^2 k^2 \frac{\partial^2 C(t_{i+1},k)}{\partial k^2}$$
(2)

Introduction

logo.png

Data-Driven Models for Discrete Hedging Problem Ke Nian

We can assume $\sigma(t_i,k)$ to be a piecewise constant functions for a given t_i and the initial condition is:

$$C(t_0, k) = \max(S(t_0) - k, 0)$$

Given the grid of strike: $k_0 < k_1 < \dots < k_n$ and a expiry t_i and next expiries t_{i+1} , let $C^j_{kk} = \frac{\partial^2 C(t_{i+1},k_j)}{\partial k^2}$ and $C^j = C(t_{i+1},k_j)$. The C^j_{kk} can approximated by finite difference:

$$C_{kk}^{j} = \frac{\frac{C_{j+1} - C_{j}}{k_{j+1} - k_{j}} - \frac{C_{j} - C_{j-1}}{k_{j} - k_{j-1}}}{\frac{k_{j+1} - k_{j-1}}{2}}$$

13 Introduction

logo.png

Data-Driven Models for Discrete Hedging

> Problem Ke Nian

Introduction

Let us assume
$$C_{kk}^0 = C_{kk}^n = 0$$
. Let

$$\alpha_j = \frac{1}{(k_j - k_{j-1})(k_{j+1} - k_{j-1})}$$
$$\beta_j = \frac{1}{(k_{j+1} - k_j)(k_j - k_{j-1})}$$

$$\eta_j = \frac{1}{(k_{j+1} - k_j)(k_{j+1} - k_{j-1})}$$
$$z_j = \sigma(t_i, k_j)^2 k^2 (t_{i+1} - t_i)$$

We have:

$$C(t_i, k_1) = -z_1 \alpha_1 C(t_{i+1}, k_1)$$

+ $(1 + z_1 \beta_1) C(t_{i+1}, k_2) - z_1 \eta_1 C(t_{i+1}, k_2)$

University of Waterloo

logo.png

Data-Driven Models for Discrete Hedging Problem

Ke Nian

15)Introduction

$$\begin{bmatrix} 1 \\ -z_1\alpha_1 & 1+z_1\beta_1 & -z_1\eta_1 \\ & -z_2\alpha_2 & 1+z_2\beta_2 & -z_2\eta_2 \\ & \ddots & \ddots & \ddots \\ & & -z_{n-1}\alpha_{n-1} & 1+z_{n-1}\beta_{n-1} & -z_{n-1}\eta_{n-1} \end{bmatrix}$$

$$M^{-1} \begin{bmatrix} C(t_i, k_0) \\ C(t_i, k_1) \\ C(t_i, k_2) \\ \vdots \\ C(t_i, k_{n-1}) \\ C(t_i, k_n) \end{bmatrix} = \begin{bmatrix} C(t_{i+1}, k_0) \\ C(t_{i+1}, k_1) \\ C(t_{i+1}, k_2) \\ \vdots \\ C(t_{i+1}, k_{n-1}) \\ C(t_{i+1}, k_n) \end{bmatrix}$$

Model Calibration

logo.png

Data-Driven Models for Discrete Hedging Problem

Ke Nian

6 Introduction

$$M^{-1} \begin{bmatrix} C(t_i, k_0) \\ C(t_i, k_1) \\ C(t_i, k_2) \\ \vdots \\ C(t_i, k_{n-1}) \\ C(t_i, k_n) \end{bmatrix} = \begin{bmatrix} C(t_{i+1}, k_0) \\ C(t_{i+1}, k_1) \\ C(t_{i+1}, k_2) \\ \vdots \\ C(t_{i+1}, k_{n-1}) \\ C(t_{i+1}, k_n) \end{bmatrix}$$

We try to find M that so that $C(t_{i+1},k_j)=C_mkt(t_{i+1},k_j)$. This can be done by:

$$\inf_{\sigma(t_{i},.)} \sum_{j} (\frac{c(t_{i+1},k_{j}) - c_{mkt}(t_{i+1},k_{j})}{Vega_{bs}^{mkt}(t_{i+1},k_{j})})^{2}$$