

Discussion Club 2018-09-04:

Soil-persistent *E. coli* and Mobile elements

Nicholas Waters

September 4, 2018

National University of Ireland, Galway, Ireland
The James Hutton Institute, Dundee, Scotland



Background

What does 10 years look like to *E. coli*?

Mobile Genetic Elements

In Closing

Background



- *E. coli* has been found to persist stably in the soil
- Isolates were cultured from lysimeter leachate
- Strains were sequenced, resulting in 149 soil-persistent *E. coli* genome



• What types of *E. coli* are able to persist in soil?



- ◊ What types of *E. coli* are able to persist in soil?
- ◊ What virulence factors are harboured by these strains?



- What types of *E. coli* are able to persist in soil?
- What virulence factors are harboured by these strains?
- What can we infer about adaptation from these?



- What types of *E. coli* are able to persist in soil?
- What virulence factors are harboured by these strains?
- What can we infer about adaptation from these?
- Can we differentiate soil-persistent *E. coli* from recent contamination?

What does 10 years look like to *E. coli*?



High estimate:

$$0.013865 * 60 * 24 * 365 * 10 \approx 72\text{k generations}$$



Medium estimate: $(5.9 / 2) * 365 * 10$

$\approx 10\text{k}$ generations

Bååth 1998

(assuming generation time roughly equals half of turnover rate)



No remaining bacteria after:

- 32 days
- 22 days
- 8 weeks
- 114 days

Detection of *Escherichia coli* in sequenced soil



Escherichia coli approximately .092% prevalence in soil



Highly diverse soil communities and environmental pressures favour rapid adaptation over incremental changes



Highly diverse soil communities and environmental pressures favour rapid adaptation over incremental changes

Adaptations will occur more rapidly because of mobile elements rather than mutations to core functions

Mobile Genetic Elements

Insertion Sequence (IS) (Transposon)

Short, mobile sequence capable of jumping



- Size: 750bp – 1.5Kb
- Detection: inverted tandem repeat lollipop with a transposase and promoter.
- Maintenance: genome replication
- Mobility: encoded transposase




Figure: Insertion Sequence


A DNA lollipop containing a transposase and promoter, and other accessory genes

- Size: < 3Kb
- Detection: inverted tandem repeat, transposase, bonus genes
- Maintenance: genome replication
- Mobility: encoded transposase

A DNA fragment with IS sequences on each end


- Size: $< 5\text{Kb}$
- Detection: double IS, promoters, and accessory genes
- Maintenance: genome replication
- Mobility: encoded transposase

Insertion sequences




Transposons

Composite transposons, e.g. Tn5



Transposons lacking terminal ISs, e.g. TnA





Mini-plasmids often containing AMR gene cassettes

- Size: $< 10\text{Kb}$
- Detection: integrase, recombination site (attI), promoter, and (optionally) a cassette of resistance genes
- Maintenance: integration into host genome
- Mobility: recombination, or transfer when circularized




- Size: $>10\text{Kb}$
- Detection: GC Skew, phylogenetic analysis of ORFs
- Maintenance: genome replication
- Mobility: various



Phage parasites

- Size: 5 – 15Kb
- Detection: Look for phage proteins along with phage inhibitors
- Maintenance: lysogeny
- Mobility: phage-like particles

Phage-inducible Chromosomal Islands





Virus integrated into the host genome

- Size: ~ 50Kb
- Detection: Look for phage integrases, tail and capsule proteins
- Maintenance: genome replication
- Mobility: lysogenic and lytic phases



Freely replicating DNA not required for survival

- Size: $<1\text{Kb} - 1\text{Mb}$
- Detection: run a gel; look for circular contigs
- Maintenance: self-replicating or integrating
- Mobility: conjugation (directly or indirectly)

In Closing



- Develop mobile pangenome tool for detecting "recent" adaptations
- Characterize trends in MGEs in environmental vs enteric *E. coli*



- o <https://www.nature.com/scitable/topicpage/transposons-the-jumping-genes-518>
- o <https://www.nature.com/scitable/topicpage/transposons-the-jumping-genes-518>
- o https://www.researchgate.net/publication/283707425_The_Phage-Inducible_Chromosomal_Islands_A_Family_of_Highly_Evolved_Molecular_Parasites
- o <https://www.sciencedirect.com/science/article/pii/S0043135416302226>
- o http://www.bx.psu.edu/~ross/workmg/TranspositionCh9_files/
- o <https://www.frontiersin.org/articles/10.3389/fmicb.2018.00762/full>



NUIG Microbiology

- ☐ Dr. Fiona Brennan
- ☐ Dr. Florence Abram
- ☐ Soil and Environmental
Microbiology Research Group
- ☐ Functional Environmental
Microbiology Group



James Hutton Institute, Dundee

- ☐ Dr. Leighton Pritchard
- ☐ Dr. Ashleigh Holmes



NUIG Microbiology

- ☐ Dr. Fiona Brennan
- ☐ Dr. Florence Abram
- ☐ Soil and Environmental Microbiology Research Group
- ☐ Functional Environmental Microbiology Group



James Hutton Institute, Dundee

- ☐ Dr. Leighton Pritchard
- ☐ Dr. Ashleigh Holmes

Questions?