NUI GALWAY

Graded CodY Activity Prioritizes Metabolism and Virulence Gene Expression in *Staphylococcus aureus*

Nick Waters

December 4, 2015

Brinsmade Lab, Biology Department, Georgetown University

BACKGROUND

· Conserved among Firmicutes




Figure 1. Crystal structure of unliganded CodY (largely inactive) and in the presence of the activator isoleucine. How GTP binds remains an open question. Credit: V. Levdikov and A.J. Wilkinson

- · Conserved among Firmicutes
- Dimer




Figure 1. Crystal structure of unliganded CodY (largely inactive) and in the presence of the activator isoleucine. How GTP binds remains an open question. Credit: V. Levdikov and A.J. Wilkinson

- · Conserved among Firmicutes
- Dimer
- · HTH motif




Figure 1. Crystal structure of unliganded CodY (largely inactive) and in the presence of the activator isoleucine. How GTP binds remains an open question. Credit: V. Levdikov and A.J. Wilkinson

- · Conserved among Firmicutes
- Dimer
- · HTH motif
- · B. subtilis CodY Structure




Figure 1. Crystal structure of unliganded CodY (largely inactive) and in the presence of the activator isoleucine. How GTP binds remains an open question. Credit: V. Levdikov and A.J. Wilkinson

- · Conserved among Firmicutes
- Dimer
- · HTH motif
- · B. subtilis CodY Structure
- Global Transcriptional Regulator




Figure 1. Crystal structure of unliganded CodY (largely inactive) and in the presence of the activator isoleucine. How GTP binds remains an open question. Credit: V. Levdikov and A.J. Wilkinson

- Conserved among Firmicutes
- Dimer
- · HTH motif
- · B. subtilis CodY Structure
- Global Transcriptional Regulator
- Two Known Effectors: ILV and GTP





Figure 1. Crystal structure of unliganded CodY (largely inactive) and in the presence of the activator isoleucine. How GTP binds remains an open question. Credit: V. Levdikov and A.J. Wilkinson

- Conserved among Firmicutes
- Dimer
- · HTH motif
- · B. subtilis CodY Structure
- Global Transcriptional Regulator
- Two Known Effectors: ILV and GTP
- · Consensus Sequence: AATTTTCWGAAAATT




and in the presence of the activator isoleucine. How GTP binds

remains an open question. Credit: V. Levdikov and A.J. Wilkinson


Nosocomial




- · Nosocomial
- · Gram Positive




- · Nosocomial
- · Gram Positive
- Colonizes skin and nares asymptomatically



- · Nosocomial until ~1990
- · Gram Positive
- Colonizes skin and nares asymptomatically



LINEAGE





QUESTIONS

 \cdot Does CodY orchestrate hierarchical expression in S. aureus?

QUESTIONS

- · Does CodY orchestrate hierarchical expression in S. aureus?
- · Does CodY modulate virulence?

QUESTIONS

- · Does CodY orchestrate hierarchical expression in *S. aureus*?
- · Does CodY modulate virulence?
- · If so, what can we infer about CodY and pathogen strategy?



· Construct Panel of Variants

- · Construct Panel of Variants
- · Extract RNA from exponential cells

- · Construct Panel of Variants
- · Extract RNA from exponential cells
- · Prepare RNA-seq libraries,

- · Construct Panel of Variants
- · Extract RNA from exponential cells
- · Prepare RNA-seq libraries,
- · Illumina seqencing

- · Construct Panel of Variants
- · Extract RNA from exponential cells
- · Prepare RNA-seq libraries,
- · Illumina seqencing
- · Align reads to reference, generate counts

· Re-annotate genome

- · Re-annotate genome
- · Make analysis pipeline

- · Re-annotate genome
- · Make analysis pipeline
 - · Filter out genes

- · Re-annotate genome
- · Make analysis pipeline
 - · Filter out genes
 - · Determine and account any batch effects

- · Re-annotate genome
- · Make analysis pipeline
 - · Filter out genes
 - · Determine and account any batch effects
 - · DESeq2

- · Re-annotate genome
- · Make analysis pipeline
 - · Filter out genes
 - · Determine and account any batch effects
 - · DESeq2
 - · Clustering of expression profiles

- · Re-annotate genome
- · Make analysis pipeline
 - · Filter out genes
 - · Determine and account any batch effects
 - · DESeq2
 - · Clustering of expression profiles
 - · GO term annotation and pathway analysis

- · Re-annotate genome
- Make analysis pipeline
 - · Filter out genes
 - · Determine and account any batch effects
 - · DESeq2
 - · Clustering of expression profiles
 - · GO term annotation and pathway analysis
- Validation

- · Re-annotate genome
- Make analysis pipeline
 - · Filter out genes
 - · Determine and account any batch effects
 - · DESeq2
 - · Clustering of expression profiles
 - · GO term annotation and pathway analysis
- Validation
 - · Nuclease Assays

- · Re-annotate genome
- Make analysis pipeline
 - · Filter out genes
 - · Determine and account any batch effects
 - · DESeq2
 - · Clustering of expression profiles
 - · GO term annotation and pathway analysis
- Validation
 - · Nuclease Assays
 - · Direct CodY-DNA interactions

- · Re-annotate genome
- Make analysis pipeline
 - · Filter out genes
 - · Determine and account any batch effects
 - · DESeq2
 - · Clustering of expression profiles
 - · GO term annotation and pathway analysis
- Validation
 - · Nuclease Assays
 - · Direct CodY-DNA interactions
 - · Protease Assays

- · Re-annotate genome
- Make analysis pipeline
 - · Filter out genes
 - · Determine and account any batch effects
 - · DESeq2
 - · Clustering of expression profiles
 - · GO term annotation and pathway analysis
- Validation
 - · Nuclease Assays
 - · Direct CodY-DNA interactions
 - · Protease Assays
- · Strategic inference

RE-ANNOTATION OF UAMS-1 GENOME

GENOME KEGGER

Problem: Previous annotation used a automated process that referenced another poorly curated genome

Solution:

 Kyoto Encyclopedia of Genes and Genomes (KEGG) downloader script

GENOME KEGGER

Problem: Previous annotation used a automated process that referenced another poorly curated genome

Solution:

- Kyoto Encyclopedia of Genes and Genomes (KEGG) downloader script
- · Genbank re-annotation script

DIFFERENTIAL EXPRESSION ANALYSIS

· Configure parameters

- · Configure parameters
- \cdot Filter out genes with no signal

- · Configure parameters
- · Filter out genes with no signal
- · Determine and apply appropriate normalization technique

- · Configure parameters
- · Filter out genes with no signal
- · Determine and apply appropriate normalization technique
- DESeq2: differential expression based on negative binomial distribution

- · Configure parameters
- · Filter out genes with no signal
- Determine and apply appropriate normalization technique
- DESeq2: differential expression based on negative binomial distribution
- · Group Expression Profiles by k-means clustering

RNA-SEQ RESULTS

Results of differential expression:

· 29 genes activated by CodY

RNA-SEQ RESULTS

Results of differential expression:


- · 29 genes activated by CodY
- · 353 genes repressed by CodY

RNA-SEQ RESULTS


Results of differential expression:

- · 29 genes activated by CodY
- · 353 genes repressed by CodY
- around 83 virulence genes regulated by CodY////


GRADED CODY REPONSE




GRADED CODY REPONSE: CLUSTERED



GRADED CODY REPONSE: HEATMAP



GRADED CODY REPONSE: ALTERNATICE DUAL CLUSTERING





VALIDATION

Thermonuclease nuc


· Biofilm maturation

- · Biofilm maturation
- · Neutrophil net escapement


- · Biofilm maturation
- · Neutrophil net escapement
- · Controlled indirectly by codY through Sae TCS

- · Biofilm maturation
- · Neutrophil net escapement
- · Controlled indirectly by codY through Sae TCS
- · Measured in vitro with FRET-based assay


- · Biofilm maturation
- · Neutrophil net escapement
- · Controlled indirectly by codY through Sae TCS
- · Measured in vitro with FRET-based assay



NUCLEASE ACTIVITY



NUCLEASE ACTIVITY



NUCLEASE ACTIVITY

