

Regular maps and the Euler characteristic

Nick Gill (ou)

April 15, 2013

What is a map?

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

What is a map?

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Let $\mathcal{G} = (V, E)$ be a graph.

What is a map?

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Let $\mathcal{G} = (V, E)$ be a graph.

Let \mathcal{S} be a surface (usually, but not always, compact and without boundary).

What is a map?

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Let $\mathcal{G} = (V, E)$ be a graph.


Let \mathcal{S} be a surface (usually, but not always, compact and without boundary).

A **map** is a 'nice' embedding of \mathcal{G} in \mathcal{S} .

This isn't nice...

Regular maps
and the Euler
characteristic


Nick Gill
(OU)



... but this is.

Regular maps
and the Euler
characteristic


Nick Gill
(OU)



This isn't nice either...

Regular maps
and the Euler
characteristic


Nick Gill
(OU)



But these are all lovely...

Regular maps
and the Euler
characteristic


Nick Gill
(OU)



But these are all lovely...

Regular maps
and the Euler
characteristic


Nick Gill
(OU)



But these are all lovely...

Regular maps
and the Euler
characteristic

Nick Gill
(OU)




R2.1 $\{3, 8\}$ 16 triangles
 $\rightarrow H_3$




R2.1' $\{3, 8\}$ 6 octagons
 $\rightarrow H_3$




R4.3' $\{6, 4\}$ 12 hexagons
 $\rightarrow \{6, 3\}_{1,1}$



R9.3' $\{6, 4\}$ 32 hexagons
 $\rightarrow R2.1' \rightarrow H_3$




R11.1 $\{4, 6\}$ 60 quads
 $\rightarrow D$




R2.2 $\{4, 6\}$ 6 quads
 $\rightarrow H_3$




R2.2' $\{6, 4\}$ 4 hexagons
 $\rightarrow H_3$




R5.1' $\{8, 3\}$ 24 octagons
 $\rightarrow C$




R9.4' $\{6, 4\}$ 32 hexagons
 $\rightarrow \{4, 4\}_{2,2}$




R13.2' $\{12, 3\}$ 24 faces
 $\rightarrow R3.4' \rightarrow H_4$




R2.3 $\{4, 8\}$ 4 quads
 $\rightarrow \{4, 4\}_{1,0}$




R2.3' $\{8, 4\}$ 2 octagons
 $\rightarrow \{4, 4\}_{1,0}$



R5.3' $\{5, 4\}$ 32 pentagons
 $\rightarrow \{4, 4\}_{2,0}$



R9.9' $\{12, 4\}$ 8 faces
 $\rightarrow \{3, 6\}_{2,0}$




R17.3' $\{6, 4\}$ 64 hexagons
 $\rightarrow \{4, 4\}_{4,0}$

And this one is especially groovy...

Regular maps
and the Euler
characteristic

Nick Gill
(OU)



Automorphisms

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- Let $\mathcal{M} = (\mathcal{G}, \mathcal{S})$ be a map.

Automorphisms

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- Let $\mathcal{M} = (\mathcal{G}, \mathcal{S})$ be a map.
- We specialise from here on to the situation where \mathcal{S} is a compact surface without boundary. The 'nice' condition implies, therefore, that the graph \mathcal{G} is finite.

Automorphisms

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- Let $\mathcal{M} = (\mathcal{G}, \mathcal{S})$ be a map.
- We specialise from here on to the situation where \mathcal{S} is a compact surface without boundary. The 'nice' condition implies, therefore, that the graph \mathcal{G} is finite.
- An automorphism of \mathcal{M} is a homeomorphism of \mathcal{S} which preserves the graph \mathcal{G} .

Automorphisms

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- Let $\mathcal{M} = (\mathcal{G}, \mathcal{S})$ be a map.
- We specialise from here on to the situation where \mathcal{S} is a compact surface without boundary. The 'nice' condition implies, therefore, that the graph \mathcal{G} is finite.
- An automorphism of \mathcal{M} is a homeomorphism of \mathcal{S} which preserves the graph \mathcal{G} . $\text{Aut}(\mathcal{M})$ is the quotient of the group of automorphisms of \mathcal{M} by the subgroup of automorphisms which fix \mathcal{G} .

Automorphisms

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- Let $\mathcal{M} = (\mathcal{G}, \mathcal{S})$ be a map.
- We specialise from here on to the situation where \mathcal{S} is a compact surface without boundary. The 'nice' condition implies, therefore, that the graph \mathcal{G} is finite.
- An automorphism of \mathcal{M} is a homeomorphism of \mathcal{S} which preserves the graph \mathcal{G} . $\text{Aut}(\mathcal{M})$ is the quotient of the group of automorphisms of \mathcal{M} by the subgroup of automorphisms which fix \mathcal{G} . In particular $\text{Aut}(\mathcal{M})$ is a finite group.

Automorphisms

Regular maps
and the Euler
characteristic


Nick Gill
(OU)

- Let $\mathcal{M} = (\mathcal{G}, \mathcal{S})$ be a map.
- We specialise from here on to the situation where \mathcal{S} is a compact surface without boundary. The ‘nice’ condition implies, therefore, that the graph \mathcal{G} is finite.
- An automorphism of \mathcal{M} is a homeomorphism of \mathcal{S} which preserves the graph \mathcal{G} . $\text{Aut}(\mathcal{M})$ is the quotient of the group of automorphisms of \mathcal{M} by the subgroup of automorphisms which fix \mathcal{G} . In particular $\text{Aut}(\mathcal{M})$ is a finite group.
- **Fact:** $\text{Aut}(\mathcal{M})$ acts faithfully and semiregularly on the set of flags.

And a flag is...

Regular maps
and the Euler
characteristic

Nick Gill
(OU)



Regular maps

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- It is clear that $|\operatorname{Aut}(\mathcal{M})|$ is less than or equal to the number of flags.

Regular maps

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- It is clear that $|\operatorname{Aut}(\mathcal{M})|$ is less than or equal to the number of flags.
- If $|\operatorname{Aut}(\mathcal{M})|$ equals the number of flags, i.e. $\operatorname{Aut}(\mathcal{M})$ acts transitively on the set of flags, then we call the map \mathcal{M} **regular**.

Regular maps

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- It is clear that $|\text{Aut}(\mathcal{M})|$ is less than or equal to the number of flags.
- If $|\text{Aut}(\mathcal{M})|$ equals the number of flags, i.e. $\text{Aut}(\mathcal{M})$ acts transitively on the set of flags, then we call the map \mathcal{M} **regular**.
- We would like to classify the regular maps.

Regular maps

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- It is clear that $|\operatorname{Aut}(\mathcal{M})|$ is less than or equal to the number of flags.
- If $|\operatorname{Aut}(\mathcal{M})|$ equals the number of flags, i.e. $\operatorname{Aut}(\mathcal{M})$ acts transitively on the set of flags, then we call the map \mathcal{M} **regular**.
- We would like to classify the regular maps.
- Encouraging fact: For any $g \geq 2$, there are only a finite number of regular maps on a surface of genus g .

$g = 0$: regular maps on the sphere

Regular maps
and the Euler
characteristic



Nick Gill
(OU)

$g = 0$: regular maps on the sphere


Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Octahedron




Cube
(Hexahedron)



Tetrahedron



Icosahedron





Dodecahedron

$g = 0$: regular maps on the sphere

Regular maps
and the Euler
characteristic



Nick Gill
(OU)



$g = 0$: regular maps on the sphere

Regular maps
and the Euler
characteristic

Nick Gill
(OU)



$\text{Aut}(\mathcal{M})$ is solvable except when \mathcal{M} is the dodecahedron or icosahedron, in which case $\text{Aut}(\mathcal{M}) \cong \text{Alt}(5)$.

Where are all the interesting groups?

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

To answer this we need to define the Euler characteristic χ of a surface \mathcal{S} :

Where are all the interesting groups?

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

To answer this we need to define the Euler characteristic χ of a surface \mathcal{S} :

- Given a surface \mathcal{S} we consider a homeomorphic CW-complex to obtain $\chi = V - E + F$.

Where are all the interesting groups?

To answer this we need to define the Euler characteristic χ of a surface \mathcal{S} :

- Given a surface \mathcal{S} we consider a homeomorphic CW-complex to obtain $\chi = V - E + F$.
- Recall that

$$\chi = \begin{cases} 2 - 2g, & \mathcal{S} \text{ orientable;} \\ 2 - g, & \mathcal{S} \text{ non-orientable.} \end{cases}$$

Where are all the interesting groups?

To answer this we need to define the Euler characteristic χ of a surface \mathcal{S} :

- Given a surface \mathcal{S} we consider a homeomorphic CW-complex to obtain $\chi = V - E + F$.
- Recall that

$$\chi = \begin{cases} 2 - 2g, & \mathcal{S} \text{ orientable;} \\ 2 - g, & \mathcal{S} \text{ non-orientable.} \end{cases}$$

- Given a map $\mathcal{M} = (\mathcal{G}, \mathcal{S})$, the embedding of \mathcal{G} on \mathcal{S} yields such a homeomorphic CW-complex, and so χ can be thought of as a function of the map.

Where are all the interesting groups?

To answer this we need to define the Euler characteristic χ of a surface \mathcal{S} :

- Given a surface \mathcal{S} we consider a homeomorphic CW-complex to obtain $\chi = V - E + F$.
- Recall that

$$\chi = \begin{cases} 2 - 2g, & \mathcal{S} \text{ orientable;} \\ 2 - g, & \mathcal{S} \text{ non-orientable.} \end{cases}$$

- Given a map $\mathcal{M} = (\mathcal{G}, \mathcal{S})$, the embedding of \mathcal{G} on \mathcal{S} yields such a homeomorphic CW-complex, and so χ can be thought of as a function of the map.
- If \mathcal{M} is regular with $G = \text{Aut}(\mathcal{M})$, then

$$\chi = V - E + F = |G| \left(\frac{1}{|G_v|} - \frac{1}{|G_e|} + \frac{1}{|G_f|} \right).$$

A meta-mathematical principle

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

A meta-mathematical principle

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Suppose that G is the automorphism group of a regular map on a surface of Euler characteristic χ .

A meta-mathematical principle

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Suppose that G is the automorphism group of a regular map on a surface of Euler characteristic χ .

General principle

If G is complicated then so is χ .

A meta-mathematical principle

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Suppose that G is the automorphism group of a regular map on a surface of Euler characteristic χ .

General principle

If G is complicated then so is χ .

G is complicated \longleftrightarrow Interesting non-abelian simple groups occur as composition factors of G .

A meta-mathematical principle

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Suppose that G is the automorphism group of a regular map on a surface of Euler characteristic χ .

General principle

If G is complicated then so is χ .

G is complicated \longleftrightarrow Interesting non-abelian simple groups occur as composition factors of G .

χ is complicated \longleftrightarrow The prime factorization of χ has many primes and/or high exponents.

Two theorems

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Suppose that G is the automorphism group of a regular map on a surface of Euler characteristic χ .

Two theorems

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Suppose that G is the automorphism group of a regular map on a surface of Euler characteristic χ .

Theorem (G., 2012)

If T is a non-abelian composition factor of G and χ is divisible by exactly x distinct primes, then T is a simple group of Lie type of rank at most x .

Two theorems

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

Suppose that G is the automorphism group of a regular map on a surface of Euler characteristic χ .

Theorem (G., 2012)

If T is a non-abelian composition factor of G and χ is divisible by exactly x distinct primes, then T is a simple group of Lie type of rank at most x .

Theorem (Conder, G., Short, Širáň, 2013)

If T is a non-abelian composition factor of G and y is the maximum exponent in the prime factorization of χ , then T is a simple group of Lie type over a field of order p^a where $a \leq y + 2$.

A word about the proof of Theorem 1

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- We saw earlier that

$$\chi = V - E + F = |G| \left(\frac{1}{|G_v|} - \frac{1}{|G_e|} + \frac{1}{|G_f|} \right).$$

A word about the proof of Theorem 1

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- We saw earlier that

$$\chi = V - E + F = |G| \left(\frac{1}{|G_v|} - \frac{1}{|G_e|} + \frac{1}{|G_f|} \right).$$

- It is easy to see that $|G_e| = 4$. Furthermore G_v and G_f contain cyclic groups of index 4.

A word about the proof of Theorem 1

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- We saw earlier that

$$\chi = V - E + F = |G| \left(\frac{1}{|G_v|} - \frac{1}{|G_e|} + \frac{1}{|G_f|} \right).$$

- It is easy to see that $|G_e| = 4$. Furthermore G_v and G_f contain cyclic groups of index 4.
- Writing m and n for the order of these two cyclic groups we obtain

$$\chi = -|G| \frac{mn - 2m - 2n}{4mn} = -\frac{|G|}{4[m, n]} \left(\frac{mn - 2m - 2n}{(m, n)} \right).$$

A word about the proof of Theorem 1

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- We saw earlier that

$$\chi = V - E + F = |G| \left(\frac{1}{|G_v|} - \frac{1}{|G_e|} + \frac{1}{|G_f|} \right).$$

- It is easy to see that $|G_e| = 4$. Furthermore G_v and G_f contain cyclic groups of index 4.
- Writing m and n for the order of these two cyclic groups we obtain

$$\chi = -|G| \frac{mn - 2m - 2n}{4mn} = -\frac{|G|}{4[m, n]} \left(\frac{mn - 2m - 2n}{(m, n)} \right).$$

- To restrict the number of (odd) primes dividing χ one needs to find elements of order m and n which are divisible by most of the primes dividing $|G|$.

A word about the proof of Theorem 1

Regular maps
and the Euler
characteristic

Nick Gill
(OU)

- We saw earlier that

$$\chi = V - E + F = |G| \left(\frac{1}{|G_v|} - \frac{1}{|G_e|} + \frac{1}{|G_f|} \right).$$

- It is easy to see that $|G_e| = 4$. Furthermore G_v and G_f contain cyclic groups of index 4.
- Writing m and n for the order of these two cyclic groups we obtain

$$\chi = -|G| \frac{mn - 2m - 2n}{4mn} = -\frac{|G|}{4[m, n]} \left(\frac{mn - 2m - 2n}{(m, n)} \right).$$

- To restrict the number of (odd) primes dividing χ one needs to find elements of order m and n which are divisible by most of the primes dividing $|G|$.
- For large rank groups of Lie type this is impossible. One can establish this formally by studying the prime graph of these groups.

Thanks for coming!