

Problems for Foundations of Mathematics

Nicholas Sales

Contents

1	Equivalence Relations	2
2	Partial or Transfinite Induction	2
3	Cardinality	3

1 Equivalence Relations

Problem 1.1: Prove that it is impossible to find an equivalence relation \equiv on \mathbb{R} for which $x \equiv y \iff (x - y) \notin \mathbb{Q}$ is true.

Proof: Let $x \in \mathbb{R}$ and assume \equiv is an equivalence relation on the real numbers. Since \equiv is an equivalence relation, it follows $x \equiv x$ as equivalence relations are reflexive. This, however, implies

$$x - x = 0 \notin \mathbb{Q}$$

This is a contradiction. Thus \equiv is not an equivalence relation. \square

Problem 1.2: Let $X = \mathbb{R}$. Show that the relation \equiv defined by $x \equiv y \iff \exists r \in X \setminus \{0\} (x = ry)$ is an equivalence relation. How many equivalence classes does \equiv have?

Proof: Let $x, y, z \in X$. To show $x \equiv x$ (reflexivity), it follows

$$x = rx \quad r \in X \setminus \{0\}$$

For $r = 1$, this is satisfied, and thus \equiv is reflexive. We now show $x \equiv y \implies y \equiv x$ (symmetry). Since $x \equiv y$, it follows $x = ry$ for some $r \in \mathbb{R} \setminus \{0\}$, with some rearrangement:

$$x = ry \implies y = \frac{1}{r}x$$

Since $\mathbb{R} \setminus \{0\}$ is closed under inverses, $\frac{1}{r} \in \mathbb{R} \setminus \{0\}$, thus \equiv is symmetric. We conclude by proving $x \equiv y \wedge y \equiv z \implies x \equiv z$ (transitivity) and stating the number of equivalence classes on \equiv . Since $x \equiv y$ and $y \equiv z$, it follows $x = ry$ and $y = sz$ for some $r, s \in \mathbb{R} \setminus \{0\}$. Replacing y in $x = ry$ by sz we obtain:

$$x = r(sz) = (rs)z$$

Since $\mathbb{R} \setminus \{0\}$ is closed under multiplication $rs \in \mathbb{R} \setminus \{0\}$, and so \equiv is transitive. We find the number of equivalence classes ($[a] = \{x \in X : x \equiv a\}$). Let $a \in X$. We consider two cases where $a = 0$ and $a \neq 0$. If $a = 0$, then it follows $x = r0 \implies x = 0$ for any $r \in \mathbb{R} \setminus \{0\}$. So $[0] = \{0\}$. If $a \neq 0$, for nonzero x , we have $x = ra \implies r = \frac{x}{a}$ which is well-defined for some $r \in \mathbb{R} \setminus \{0\}$. Thus $[a] = \mathbb{R} \setminus \{0\}$. \square

2 Partial or Transfinite Induction

Problem 2.1: Let (X, \preceq) be a totally ordered set. Prove that if $x_o \in X$ is a minimal element, then it is also a minimum element.

Proof: Recall that x_o is a minimal element if there is no $x \in X$ such that $x \preceq x_o$ and $x \neq x_o$, and it is a minimum element if for any $x \in X$ it follows $x_o \preceq x$. We prove this by contradiction. Assume $x_o \in X$ is minimal but not a minimum element. Since X is totally ordered, it follows $x \preceq x_o$ or $x_o \preceq x$ for any $x \in X$. It follows $x \preceq x_o$ as x_o is assumed not to be a minimum element. It must be that $x \neq x_o$ as $x = x_o \wedge x \preceq x_o \implies x_o \preceq x$ by reflexivity of \preceq . However, if $x \neq x_o$ with $x \preceq x_o$ this is contradictory to our claim that x_o is minimal. Therefore x_o must be a minimum element. \square

Problem 2.2: Let (X, \preceq) be a totally ordered set, and $A \subseteq X$. Prove that $\sup(A)$ and $\inf(A)$, if exist, are unique.

Proof: Recall the supremum is defined as an upper bound of A ($a \preceq x$ for all $a \in A$), x , such that for any upper bound of A , x' , $x \preceq x'$. And the infimum is defined as a lower bound of A ($y \preceq a$ for all $a \in A$), y , such that for any lower bound of A , y' , $y' \preceq y$. We prove this by contradiction. Assume x, x_o are both $\sup(A)$ and y, y_o are both $\inf(A)$ with $x \neq x_o$ and $y \neq y_o$. Since x, x_o are both upper bounds of A , and likewise y, y_o are both lower bounds of A , by the definition of supremum and infimum:

$$\begin{aligned} x &\preceq x_o \text{ and } x_o \preceq x \\ y_o &\preceq y \text{ and } y \preceq y_o \end{aligned}$$

Since \preceq is antisymmetric, the above can only hold if $x = x_o$ and $y = y_o$. This contradicts our assumption that $x \neq x_o$ and $y \neq y_o$. Therefore $\sup(A)$ and $\inf(A)$ must be unique. \square

Problem 2.3: Let \prec_Y be a strict partial order on a set Y , and $f : X \rightarrow Y$ be a function. Define a relation \prec_X on X as follows. For every $x_1, x_2 \in X$:

$$x_1 \prec_X x_2 \iff f(x_1) \prec_Y f(x_2)$$

Prove that \prec_X is a strict partial order on X .

Proof: Recall that \prec_Y is a strict partial order if it is transitive and irreflexive. Let $x_1, x_2, x_3 \in X$. We first prove transitivity. Assume $x_1 \prec_X x_2$ and $x_2 \prec_X x_3$. It follows that

$$f(x_1) \prec_Y f(x_2) \text{ and } f(x_2) \prec_Y f(x_3)$$

This implies $f(x_1) \prec_Y f(x_3)$ by transitivity of \prec_Y being a strict partial order. So it also follows $x_1 \prec_X x_3$, and hence \prec_X is transitive. We now show \prec_X is irreflexive by contradiction. Assume $x_1 \prec_X x_1$. It follows by the definition of \prec_X

$$f(x_1) \prec_Y f(x_1)$$

This is, however, a contradiction as \prec_Y is a strict partial order and so is irreflexive. Thus $x_1 \not\prec_X x_1$, and since \prec_X is both transitive and irreflexive it is a strict partial order. \square

3 Cardinality

Problem 3.1: Prove the following statements by giving explicit bijections:

a) $2\mathbb{Z} \sim \mathbb{N}$

Proof: We split $2\mathbb{Z}$ into two disjoint sets $2\mathbb{Z}_{<0} = \{\dots, -6, -4, -2\}$ and $2\mathbb{Z}_{\geq 0} = \{0, 2, 4, \dots\}$ that satisfy $2\mathbb{Z}_{<0} \cup 2\mathbb{Z}_{\geq 0} = 2\mathbb{Z}$. We can define the bijection

$$f_1 : 2\mathbb{Z}_{<0} \rightarrow O \subseteq \mathbb{N}, \quad x \mapsto -x - 1$$

where $O = \{1, 3, 5, 7, \dots\}$, as the mapping of negative even integers onto odd natural numbers. Now, we can define the bijection

$$f_2 : 2\mathbb{Z}_{\geq 0} \rightarrow E \subseteq \mathbb{N}, \quad x \mapsto x + 2$$

where $E = \{2, 4, 6, 8, \dots\}$, as the mapping of positive even integers onto even natural numbers. Since O and E are disjoint and $O \cup E = \mathbb{N}$, we define

$$f : 2\mathbb{Z} \rightarrow \mathbb{N}, \quad f(x) = \begin{cases} f_1(x) & \text{if } x < 0 \\ f_2(x) & \text{if } x \geq 0 \end{cases}$$

which is a bijection as both f_1 and f_2 are bijective. Since there exists a bijection from $2\mathbb{Z}$ to \mathbb{N} , the set of even integers is equinumerous with the natural numbers. \square