

# MAT 2141 Notes

Nicholas Sales

## Contents

|          |  |          |
|----------|--|----------|
| <b>1</b> | <b>Linear Maps</b>                       | <b>2</b> |
| 1.1      | Definiton . . . . .                      | 2        |
| 1.2      | Kernel and Image . . . . .               | 2        |
| 1.3      | Vector Spaces of Linear Maps . . . . .   | 3        |
| 1.4      | Isomorphisms . . . . .                   | 3        |
| <b>2</b> | <b>Structure of Vector Spaces</b>        | <b>4</b> |
| 2.1      | Spans and Generating Sets . . . . .      | 4        |
| 2.2      | Linear Dependence/Independence . . . . . | 5        |

# 1 Linear Maps

## 1.1 Definiton

**Definition 1.1.1:** If  $V$  and  $W$  are vector spaces over the same field  $F$ , and  $T : V \rightarrow W$  is a mapping, we say  $T$  is a linear mapping if the following hold:

$$\begin{aligned}T(u + v) &= T(u) + T(v) \quad \forall u, v \in V \\T(cv) &= cT(v) \quad \forall v \in V, \forall c \in F\end{aligned}$$

**Example 1.1.1:** Let  $V = C^\infty(\mathbb{R})$ ,  $F = \mathbb{R}$ , and define  $S : C^\infty(\mathbb{R}) \rightarrow C^\infty(\mathbb{R})$  by  $(Sf)(x) = \int_0^x f(t)dt$ ,  $\forall x \in \mathbb{R}$ .

Let  $f, g \in V$  (i.e.  $f$  and  $g$  are infinitely differentiable functions) and let  $c \in F$ . Using the properties of integration, we can show additivity and homogeneity hold:

$$\begin{aligned}S(f + g)(x) &= \int_0^x f(t) + g(t)dt = \int_0^x f(t)dt + \int_0^x g(t)dt = (Sf)(x) + (Sg)(x) \\S(cf)(x) &= \int_0^x cf(t)dt = c \int_0^x f(t)dt = c(Sf)(x)\end{aligned}$$

Since  $S$  satisfies the properties of additivity and homogeneity, we conclude  $S$  is a linear map. QED.

## 1.2 Kernel and Image

**Definiton 1.2.1:** If  $T : V \rightarrow W$  is a linear map, then  $Ker(T) \subseteq V$  gets mapped to the zero vector under  $T$  and  $Im(T) \subseteq W$  is where vectors from  $V$  can actually get mapped to under  $T$  (i.e. range):

$$\begin{aligned}Ker(T) &= T^{-1}(\vec{0}) = \{x \mid Tx = \vec{0}\} \\Im(T) &= T(V) = \{Tx \mid x \in V\}\end{aligned}$$

**Theorem 1.2.1:** Suppose  $T : V \rightarrow W$  is a linear mapping,  $T$  is injective iff  $Ker(T) = \{\vec{0}\}$ .

We prove this in two directions. Assume  $T$  is injective and let  $u, v \in V$ . Since  $0v = \vec{0}$  for any vector  $v$  and  $T(0v) = 0T(v)$  by the property of linear maps, it follows a linear mapping always maps the zero vector to the zero vector. Since for any injective mapping we have  $T(u) = T(v) \implies u = v$ , it follows if  $T(u) = T(v) = \vec{0}$ , then  $u = v = \vec{0}$  and thus  $Ker(T) = \{\vec{0}\}$ .

We now prove this in the other direction. Assume  $Ker(T) = \{\vec{0}\}$ . We want to show if  $T(u) = T(v)$  for any  $u, v \in V$  then  $u = v$ . If  $T(u) = T(v)$ , it follows  $T(u) - T(v) = \vec{0}$  which simplifies to  $T(u - v) = \vec{0}$  by the property of linear maps. Since  $Ker(T) = \{\vec{0}\}$  it follows that  $T(u - v) = \vec{0} \implies u - v = \vec{0} \implies u = v$ . Thus  $T$  is injective. Since we have proved both implications this completes the proof. QED.

### 1.3 Vector Spaces of Linear Maps

**Definition 1.3.1:** If  $V, W$  are vector spaces over a field  $F$ , we define the set of linear maps from  $V$  to  $W$ :

$$\mathcal{L}(V, W) = \{T : V \rightarrow W \mid T \text{ is linear}\}$$

If  $V = W$ , we simply write  $\mathcal{L}(V)$ .

**Definition 1.3.2:** If  $A, B$ , and  $C$  are sets with mappings  $T : A \rightarrow B$  and  $S : B \rightarrow C$ , we can compose the mappings to get  $ST : A \rightarrow C$  (i.e.  $S \circ T$ ) defined by:

$$(ST)a = S(Ta) \quad \forall a \in A$$

**Theorem 1.3.1:** The composition of linear maps is itself a linear map (i.e.  $T \in \mathcal{L}(U, V)$  and  $S \in \mathcal{L}(V, W) \implies ST \in \mathcal{L}(U, W)$ ).

Let  $T \in \mathcal{L}(U, V)$  and  $S \in \mathcal{L}(V, W)$  be linear maps over a field  $F$  with  $u, v \in U$  and  $c \in F$ . By the definition of composition we have:

$$\begin{aligned}(ST)(u + v) &= S(T(u + v)) \\ (ST)(cu) &= S(T(cu))\end{aligned}$$

By the property of  $T$  being a linear map we get:

$$\begin{aligned}S(T(u + v)) &= S(T(u) + T(v)) \\ S(T(cu)) &= S(cT(u))\end{aligned}$$

And further, by the property of  $S$  being a linear map we get:

$$\begin{aligned}S(T(u) + T(v)) &= S(T(u)) + S(T(v)) \\ S(cT(u)) &= cS(T(u))\end{aligned}$$

Thus,  $ST$  satisfies additivity and homogeneity and is therefore a linear map. QED.

### 1.4 Isomorphisms

**Definition 1.4.1:** If  $T : V \rightarrow W$  is a bijective linear map, we say  $T$  is an isomorphism. We also say that  $V$  is isomorphic to  $W$  and write  $V \cong W$ .

**Theorem 1.4.1:** Suppose  $V$  and  $W$  are vector spaces. Further, assume  $T : V \rightarrow W$  is an isomorphism. It follows that  $T^{-1} : W \rightarrow V$  is also linear and hence an isomorphism.

Assume  $T : V \rightarrow W$  is an isomorphism. Since it follows that  $T$  is bijective, it is invertible. Let  $u', v' \in W$ . It follows there exists some  $u, v \in V$  such that  $T(u) = u'$  and  $T(v) = v'$ . By the property of  $T$  being linear, we have:

$$\begin{aligned}T^{-1}(u' + v') &= T^{-1}(T(u) + T(v)) = T^{-1}(T(u + v)) = u + v = T^{-1}(u') + T^{-1}(v') \\ T^{-1}(cu') &= T^{-1}(cT(u)) = T^{-1}(T(cu)) = cu = cT^{-1}(u')\end{aligned}$$

Thus  $T^{-1}$  preserves additivity and homogeneity and is therefore a linear map. And moreover, since it is the inverse of a bijective mapping, it is also bijective, and thus an isomorphism. QED.

**Theorem 1.4.2:** Isomorphisms are an equivalence relation. In other words, if  $U, V, W$  are all vector spaces over the same field  $F$ , the following properties hold:

$$\begin{aligned} U &\cong U \\ U &\cong V \implies V \cong U \\ U &\cong V \text{ and } V \cong W \implies U \cong W \end{aligned}$$

This proof is left to the non-existent reader.

## 2 Structure of Vector Spaces

### 2.1 Spans and Generating Sets

**Theorem 2.1.1:** Suppose  $A$  is a nonempty subset of a vector space  $V$  and that  $W$  is a subspace of  $V$ . Then  $\text{Span}(A)$  is the smallest subspace of  $V$  containing  $A$ . In other words:

$$\begin{aligned} A &\subseteq \text{Span}(A) \\ A &\subseteq W \implies \text{Span}(A) \subseteq W \end{aligned}$$

Let  $\alpha \in A$ . Since  $\text{Span}(A)$  is the set of all linear combinations of vectors in  $A$ , it follows the linear combination  $1\alpha = \alpha \in \text{Span}(A)$ . Thus  $A \subseteq \text{Span}(A)$ .

Assume  $A \subseteq W$  and  $w \in \text{Span}(A)$ . It follows we can write  $w = c_1a_1 + \cdots + c_na_n$  for some  $c_1, \dots, c_n \in F$  and  $a_1, \dots, a_n \in A$ . Since  $W$  is a vector space, it satisfies closure properties under addition and scalar multiplication. Further, since  $A \subseteq W$ , it follows that  $a_i + a_j \in W$  and  $c_ia_i \in W$  for any  $a_i, a_j \in A$  and  $c_i \in F$ . Therefore, it must follow that  $w = c_1a_1 + \cdots + c_na_n \in W$  and we can conclude  $\text{Span}(A) \subseteq W$ . QED.

**Definition 2.1.1:** If  $V$  is a vector space with  $A \subseteq V$ , we say that  $\text{Span}(A)$  is the subspace of  $V$  generated by  $A$ , and thus  $A$  is the generating set for  $\text{Span}(A)$ . We say that  $A$  generates  $\text{Span}(A)$ , so it follows if  $\text{Span}(A) = V$ , we say  $A$  generates  $V$ .

**Theorem 2.1.2:** Suppose  $T : V \rightarrow W$  is a linear map and  $A$  is a subset of  $V$ . Then  $\text{Span}T(A) = T(\text{Span}(A))$ .

Let  $a_1, \dots, a_n$  be the vectors in  $A$  and let  $c_1, \dots, c_n \in F$ . We prove this in two directions. First, assume  $c_1T(a_1) + \cdots + c_nT(a_n) \in \text{Span}T(A)$ . We can use the properties of linear maps to show:

$$\begin{aligned} c_1T(a_1) + \cdots + c_nT(a_n) &= T(c_1a_1) + \cdots + T(c_na_n) \\ &= T(c_1a_1 + \cdots + c_na_n) \in T(\text{Span}(A)) \end{aligned}$$

Thus it follows  $\text{Span}T(A) \subseteq T(\text{Span}(A))$ . In the other direction, assume  $T(c_1a_1 + \cdots + c_na_n) \in T(\text{Span}(A))$ . Again, by the properties of linear maps:

$$\begin{aligned} T(c_1a_1 + \cdots + c_na_n) &= T(c_1a_1) + \cdots + T(c_na_n) \\ &= c_1T(a_1) + \cdots + c_nT(a_n) \in \text{Span}T(A) \end{aligned}$$

Thus  $T(\text{Span}(A)) \subseteq \text{Span}T(A)$ . We therefore conclude  $\text{Span}T(A) = T(\text{Span}(A))$ . QED.

## 2.2 Linear Dependence/Independence

**Definition 2.2.1:** Suppose  $V$  is a vector space with  $v_1, \dots, v_n \in V$  and  $c_1, \dots, c_n \in F$ . If at least one of the  $c_i$  is nonzero and:

$$c_1v_1 + \dots + c_nv_n = \vec{0},$$

we say that the list of vectors is linearly dependent. If the  $c_i$  must all be zero for the equation to hold true, the list is said to be linearly independent.

**Definition 2.2.2:** If  $A$  is any (possibly infinite) set of vectors in a vector space  $V$ , if every finite list of distinct vectors in  $A$  is linearly independent, then  $A$  is linearly independent. However, if there is some finite list of vectors in  $A$  that is linearly dependent, then  $A$  is linearly dependent.

**Theorem 2.2.1:** Suppose  $V$  is a vector space over a field  $F$  with  $v_1, \dots, v_n \in V$ . Define the linear map  $T : F^n \rightarrow V$  by  $T(a_1, \dots, a_n) = a_1v_1 + \dots + a_nv_n$ . The list of vectors  $v_1, \dots, v_n$  are linearly dependent iff  $T$  is not injective.

We prove this in two directions. First, assume  $v_1, \dots, v_n$  are linearly dependent. Since the vectors are linearly dependent, there exists some  $a_1, \dots, a_n \in F$ , not all zero, such that  $a_1v_1 + \dots + a_nv_n = \vec{0}$ . It follows that  $(a_1, \dots, a_n) \in \text{Ker}(T) \implies \text{Ker}(T) \neq \{\vec{0}\}$ . Thus  $T$  cannot be injective.

Now, assume  $T$  is not injective. It follows there is some  $(a_1, \dots, a_n), (b_1, \dots, b_n) \in F^n$  such that  $T(a_1, \dots, a_n) = T(b_1, \dots, b_n)$  and  $(a_1, \dots, a_n) \neq (b_1, \dots, b_n)$ . We thus have:

$$\begin{aligned} T(a_1, \dots, a_n) = T(b_1, \dots, b_n) &\implies a_1v_1 + \dots + a_nv_n = b_1v_1 + \dots + b_nv_n \\ &\implies (a_1 - b_1)v_1 + \dots + (a_n - b_n)v_n = \vec{0} \end{aligned}$$

Since  $(a_1, \dots, a_n) \neq (b_1, \dots, b_n)$ , it follows that  $(a_1 - b_1, \dots, a_n - b_n) \in F^n$  is nonzero, meaning at least one component  $a_i - b_i \neq 0$ . Since  $(a_1 - b_1, \dots, a_n - b_n) \in \text{Ker}(T)$  and is nonzero, the vectors  $v_1, \dots, v_n$  are linearly dependent. QED.

**Theorem 2.2.2:** Suppose  $T : V \rightarrow W$  is an injective linear map and  $v_1, \dots, v_n$  are linearly independent vectors in  $V$ , then  $T(v_1), \dots, T(v_n)$  are linearly independent vectors in  $W$ .

Let  $c_1, \dots, c_n \in F$ . Since  $T$  is a linear map, we have:

$$c_1T(v_1) + \dots + c_nT(v_n) = \vec{0} \implies T(c_1v_1 + \dots + c_nv_n) = \vec{0}$$

Further, since  $T$  is injective and  $v_1, \dots, v_n$  are linearly independent:

$$T(c_1v_1 + \dots + c_nv_n) = \vec{0} \implies c_1v_1 + \dots + c_nv_n = \vec{0} \implies c_1 = \dots = c_n = 0$$

QED.