

Predicting the efficacy of malaria vaccines across the life-course

Supplementary Notebook 2. Averted deaths in a cohort vs. a population

Nicholas J. Savill

Philip J. Spence

Previous vaccine modelling studies have modelled intervention campaigns in stable, multi-cohort populations over a fixed time interval, usually 15 years, with constant and equal birth and death rates (e.g. Penny et al. (2016); Schmit et al. (2024)). Two reasons have been given to justify a fixed endpoint of 15 years: to be comparable to the timeframe of the Global Technical Strategy for Malaria (“World Health Organization. Global Technical Strategy for Malaria 2016–2030, 2015.” 2015; Penny et al. 2016), and to capture the negative effects of infections shifting to older children - meaning rebound and the higher risk of severe malaria-associated mortality in older children (Reyburn et al. 2005; Penny et al. 2016; Schmit et al. 2024).

Unfortunately, no data exists to refute or support the assumption that the detrimental effects of rebound will be negligible after 15 years due to the practical difficulties of obtaining such data (Pemberton-Ross et al. 2015). Even assuming that rebound is negligible after 15 years, it is only the children at the start of the campaign that will have experienced the full detriment of rebound; younger children will have experienced only a fraction of its negative effects. Thus, calculating efficacy at some fixed endpoint will truncate and reduce the magnitude of the detrimental effects of rebound at the population level. In addition, the youngest children are over-represented in the calculation of vaccine efficacy. For example, after 15 years there are 15 times more newborns in total than 15-year-olds because there have been 15 years of birth, but only one cohort has aged to 15 years old. And it is the youngest children that benefit most from vaccination due to its waning efficacy. For these reasons, modelling a vaccine campaign in a population over a fixed time period over-estimates averted cases and vaccine efficacy.

Vaccine trials, on the other hand, are longitudinal studies in single, multi-age cohorts. Efficacy is calculated for this single cohort relative to an unvaccinated control cohort. This definition directly expresses a vaccine’s efficacy as the percentage reduction in risk of morbidity and mortality over the cohort’s lifetime.

Figure 1 compares averted deaths over 20 years either in a multi-cohort population (blue bars) or a single cohort (purple bars) for R21 given to 100,000 5 month olds. Note the different meaning on the x -axis. The single cohort completes three rounds of vaccination then receives no more doses; time on the x -axis represents time since the third and last vaccine dose. For the multi-cohort population, each year a new cohort of 5 month olds complete three rounds of vaccination; time on the x -axis represents time since the start of the vaccine campaign. Hence vaccine-induced protection in the population is being “topped-up” every year, whereas in the single cohort, protection gradually declines from a maximum at $t = 0$.

Averted deaths are positive for the first five years but decline due to the waning efficacy of R21. After five years rebound occurs (negative averted deaths) due to R21 temporarily reducing the risk of infection and thereby attenuating development of naturally acquired immunity to clinical and severe malaria. After 20 years rebound is negligible and cumulative averted deaths plateaus. Vaccination in the population averts 21 deaths, whereas vaccination in the single cohort indirectly causes 45 deaths.




Figure 1: Annual averted deaths per 100,000 R21-fully vaccinated five month old children, comparing a multi-cohort population (blue) to a single cohort (purple) and assuming 1 blood infection every 2 years in a perennial setting. The multi-cohort population assumes a vaccine campaign that vaccinates 100% of five month old children each year for 20 years, while the single cohort assumes a one-time vaccination of 100% of five month old children at the start of the study. The total number of deaths averted over the study period is shown in the legend.

Figure 2 compares deaths averted and death efficacy between a multi-cohort population and a single cohort followed for 15 years across a range of blood infection rates. In all cases, the population approach overestimates averted deaths and efficacy, and is particularly poor at low transmission intensities.




Figure 2: Averted deaths per 100,000 R21-fully vaccinated five month old children and death efficacy after 15 years from the start of vaccination across a range of blood infection rates, comparing a multi-cohort population (blue) to a single cohort (purple).

Bibliography

Pemberton-Ross, Peter, Thomas A. Smith, Eva Maria Hodel, Katherine Kay, and Melissa A. Penny. 2015. "Age-Shifting in Malaria Incidence as a Result of Induced Immunological Deficit: A Simulation Study". *Malaria Journal* 14 (1): 0–14. <https://doi.org/10.1186/s12936-015-0805-1>.

- Penny, Melissa A., Robert Verity, Caitlin A. Bever, Christophe Sauboin, Katya Galactionova, Stefan Flasche, Michael T. White, et al. 2016. "Public Health Impact and Cost-Effectiveness of the RTS,S/AS01 Malaria Vaccine: A Systematic Comparison of Predictions from Four Mathematical Models". *Lancet* 387 (10016): 367–75. [https://doi.org/10.1016/S0140-6736\(15\)00725-4](https://doi.org/10.1016/S0140-6736(15)00725-4).
- Reyburn, Hugh, Redempta Mbatia, Chris Drakeley, Jane Bruce, Ilona Carneiro, Raimos Olomi, Jonathan Cox, et al. 2005. "Association of Transmission Intensity and Age with Clinical Manifestations and Case Fatality of Severe *Plasmodium Falciparum* Malaria". *JAMA* 293 (12): 1461–70. <https://doi.org/10.1001/jama.293.12.1461>.
- Schmit, Nora, Hillary M. Topazian, H. Magloire Natama, Duncan Bellamy, Ousmane Traoré, M. Athanase Somé, Toussaint Rouamba, et al. 2024. "The Public Health Impact and Cost-Effectiveness of the R21/matrix-M Malaria Vaccine: A Mathematical Modelling Study". *Lancet Infectious Diseases* 24 (5): 465–75. [https://doi.org/10.1016/S1473-3099\(23\)00816-2](https://doi.org/10.1016/S1473-3099(23)00816-2).
- "World Health Organization. Global Technical Strategy for Malaria 2016–2030, 2015." 2015. Geneva.