



Target Trials in Policy Evaluation: A Case Study in Medical Cannabis Laws

Nicholas J. (Nick) Seewald, PhD

Department of Health Policy and Management Johns Hopkins Bloomberg School of Public Health

Joint with E.E. McGinty, E.A. Stuart

Annals of Internal Medicine

Original Research

Effects of U.S. State Medical Cannabis Laws on Treatment of Chronic Noncancer Pain

Emma E. McGinty, PhD; Kayla N. Tormohlen, PhD; Nicholas J. Seewald, PhD; Mark C. Bicket, MD, PhD; Alexander D. McCourt, JD, PhD; Lainie Rutkow, JD, PhD; Sarah A. White, MS; and Elizabeth A. Stuart, PhD

This work will appear in the July issue of *Annals of Internal Medicine*.

Please limit outside discussion of substantive findings until then!

Disclosures

I have a family member employed by a cannabis distributor in Michigan.

Study funded by National Institute on Drug Abuse: Ro1DA049789 (PI: McGinty)

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.





- Cannabis industry and advocates have argued that medical cannabis could be a partial solution to the opioid overdose crisis [1]
 - Substitution of cannabis for opioids to treat chronic non-cancer pain
- Clinical guidelines do **not** recommend cannabis
- ► Chronic non-cancer pain is a qualifying condition for medical cannabis under all 38 existing state (+DC) programs [2]
- Some evidence of substitution of cannabis for prescription opioids among patients [3]
- ▶ **Question:** What are the effects of state medical cannabis laws on receipt of opioid and guideline-concordant non-opioid pain treatments for chronic non-cancer pain?
- 1. https://thecannabisindustry.org/combating-the-opioid-epidemic/
- 2. https://www.ncsl.org/health/state-cannabis-policy-enactment-database
- 3. Bicket MC, et al. JAMA Network Open. 2023.





- Necessarily limited sample size
- ► Often high variability in definitions of treatment
 - "States are the laboratories of democracy" [1]
- ▶ Hard to isolate a policy's effects when other policies go into place around the same time

▶ **Partial solution:** Be very thoughtful about design! (surprise)

Trial Emulation Framework: Estimand & Scientific Question



Hypothetical Target Trial

- Estimand is typically ATE: E[Y(1) Y(0)]
- ► "In general, what is the effect on outcomes of a state implementing a medical cannabis law versus not implementing a medical cannabis law?"

Our Policy Trial Emulation Analogue

Estimand is ATT:

$$E[Y(1) - Y(0) | A = 1]$$

- * "Among states that implemented a medical cannabis law, what was the effect of the law on outcomes relative to what would have been observed had those states not implemented a medical cannabis law?"
- Only interested in studying policies on the books, rather than hypothetical policies


(ATT = ATE under random assignment or no treatment effect heterogeneity)

Trial Emulation Framework: Units



Hypothetical Target Trial AND our Policy Trial Emulation Analogue

- ▶ 12 "treated" states implemented a medical cannabis law between 2012 and 2019 and did not also implement a recreational cannabis program in that time.
- ▶ 17 "control" states did not implement medical or recreational cannabis laws



Trial Emulation Framework: Exposure & Outcomes



Hypothetical Target Trial AND our Policy Trial Emulation Analogue

- ▶ <u>Exposure</u>: Implementation of a medical cannabis law that includes chronic non-cancer pain diagnoses as qualifying conditions for receipt of medical cannabis
- ▶ <u>Outcomes</u>: Various measures of opioid and guideline-concordant non-opioid prescribing measured in time period after policy implementation (or lack of implementation)

Trial Emulation Framework: Assignment Procedure



Hypothetical Target Trial


- Random assignment of states to implement or not implement a medical cannabis law after 4 years of baseline data collection.
- ► Unblinded: states will be aware of randomization status
- Essentially cluster-randomized (data from individuals within states)

Our Policy Trial Emulation Analogue

Nonrandom policy adoption, possibly influenced by both known and unknown state-level characteristics

Staggered Adoption of Medical Cannabis Laws





Staggered Adoption Causes Problems with Traditional Methods



Research question in medical cannabis study is about an ATT

$$E[Y(1) - Y(0) | A = 1]$$


on average over the treated states.

- ▶ Traditional policy evaluation method turns out to be *very biased* for this estimand under staggered adoption when treatment effect is time-varying (i.e., almost always) [1]
- ▶ *But:* it's okay when we look at one treated state at a time.

1. Goodman-Bacon A. J Econometrics, 2021.

"Stacking" (Serial Trial Emulation)





Fix study periods

1. Hernán MA, Robins JM. Am. J. Epidemiol. 2016.


Start with full data

2. Ben-Michael E, Feller A, Stuart EA. *Epidemiology*. 2021.

Anchor time at policy implementation

"Stacking" (Serial Trial Emulation)





Anchor time at policy implementation

Estimate state-specific effects

Aggregate state-specific effects (using, e.g., inverse-variance weighting)

- 1. Hernán MA, Robins JM. Am. J. Epidemiol. 2016.
- 2. Ben-Michael E, Feller A, Stuart EA. *Epidemiology*. 2021.

Trial Emulation Framework: Data Collection Units




Hypothetical Target Trial

- ▶ People living in exposed & unexposed states with a chronic non-cancer pain diagnosis in the 4 years prior to policy implementation.
- ► Ideally people would not be allowed to move across states, wouldn't die, and would contribute complete data
 - Avoid compositional changes over time


Our Policy Trial Emulation Analogue

- People living in the treated state or one of the untreated states with a chronic non-cancer pain diagnosis in treated state's 4-year prelaw period
- Continuously enrolled in commercial health insurance for entire 7-year study period
 - Avoid compositional changes over time
 - No reason to believe enrollment is related to implementation of cannabis law



















Trial Emulation Framework: Analytic Strategy



Hypothetical Target Trial

- "Traditional" modeling approach for cluster-randomized trial with longitudinal outcome
- Effect estimation unconfounded due to randomization


Our Policy Trial Emulation Analogue

- Stacked effect estimation
- Must account for potential confounders
 - ► Idiosyncratic in "difference-in-differences" setups
- We used the augmented synthetic control method [1]

1. Ben-Michael E, Rothstein J, Feller A. J Am Stat Assoc. 2021.

Medical Cannabis Study Results: Proportion Receiving Opioid Prescriptions





State-specific effects of medical cannabis laws on


the proportion of chronic noncancer pain patients receiving <u>any opioid prescription</u> in a given month,

on average over the first 3 years of law implementation.

Inverse-variance weighted aggregate effect:

0.05 (-0.12, 0.21) pct points

Medical Cannabis Study Results: Proportion Receiving Non-Opioid Prescriptions



State-specific effects of medical cannabis laws on

the proportion of chronic noncancer pain patients receiving <u>any non-opioid prescription</u> in a given month,

on average over the first 3 years of law implementation.

Inverse-variance weighted aggregate effect:

0.05 (-0.13, 0.23) pct points





- ▶ Trial emulation provides a nice framework for good study design
 - Careful consideration of estimand, baseline, analysis
- ▶ Avoids issues with traditional kitchen-sink modeling approaches in policy evaluation
 - State-specific estimates are useful!
- ► Can go further: might allow changing control pool if comparison states implement confounding policies (i.e., different controls for each treated state)

Acknowledgements

Co-authors: Beth McGinty, Kayla Tormohlen, Mark Bicket, Alex McCourt, Lainie Rutkow, Sarah White, Liz Stuart

NIDA R01DA049789