

---

# Kmeans++

— Boston University CS 506 - Lance Galletti —

---

# K-means - Lloyd's Algorithm

Q1: Will this algorithm always converge?

**Proof** (by contradiction): Suppose it does not converge. Then, either:

1. The minimum of the cost function is only reached in the limit (i.e. after an infinite number of iterations).

**Impossible** because we are iterating over a finite set of partitions

1. The algorithm gets stuck in a cycle / loop




**Impossible** since this would require having a clustering that has a lower cost than itself and we know:




- If  $\text{old} \neq \text{new}$  clustering then the cost has improved
- If  $\text{old} = \text{new}$  clustering then the cost is unchanged




**Conclusion:** Lloyd's Algorithm always converges!




# K-means - Lloyd's Algorithm




Q2: Will this always converge to the optimal solution?




























# What's the problem?




















## Farthest First Traversal









**But...**





















Random would have  
been better




# K-means++

Initialize with a combination of the two methods:


1. Start with a random center
2. Let  $\mathbf{D}(\mathbf{x})$  be the distance between  $\mathbf{x}$  and the closest of the centers picked so far. Choose the next center with probability proportional to  $\mathbf{D}(\mathbf{x})^2$

# K-means++









# K-means++



# K-means++



# K-means++



No reason to use k-means over  
k-means++




# K-means++

Suppose we are given a black box that will generate a uniform random number between 0 and any **N**. How can we use this black box to select points with probability proportional to **D(x)<sup>2</sup>**?


# K-means++

Suppose we are given a black box that will generate a uniform random number between 0 and any **N**. How can we use this black box to select points with probability proportional to  **$D(\mathbf{x})^2$** ?



# K-means++

Suppose we are given a black box that will generate a uniform random number between 0 and any **N**. How can we use this black box to select points with probability proportional to  $D(\mathbf{x})^2$ ?




$$D(\mathbf{x})^2 = 3^2 = 9$$

$$D(\mathbf{y})^2 = 2^2 = 4$$

$$D(\mathbf{z})^2 = 1^2 = 1$$

# K-means++


Suppose we are given a black box that will generate a uniform random number between 0 and any **N**. How can we use this black box to select points with probability proportional to  $D(\mathbf{x})^2$ ?



$$D(\mathbf{x})^2 = 3^2 = 9$$


$$D(\mathbf{y})^2 = 2^2 = 4$$

$$D(\mathbf{z})^2 = 1^2 = 1$$



# K-means++


Suppose we are given a black box that will generate a uniform random number between 0 and any **N**. How can we use this black box to select points with probability proportional to  $D(\mathbf{x})^2$ ?



$$D(\mathbf{x})^2 = 3^2 = 9$$

$$D(\mathbf{y})^2 = 2^2 = 4$$


$$D(\mathbf{z})^2 = 1^2 = 1$$





# K-means++


Suppose we are given a black box that will generate a uniform random number between 0 and any **N**. How can we use this black box to select points with probability proportional to  $D(\mathbf{x})^2$ ?



$$D(\mathbf{x})^2 = 3^2 = 9$$


$$D(\mathbf{y})^2 = 2^2 = 4$$

$$D(\mathbf{z})^2 = 1^2 = 1$$



# K-means++


Suppose we are given a black box that will generate a uniform random number between 0 and any **N**. How can we use this black box to select points with probability proportional to  $D(\mathbf{x})^2$ ?



$$D(\mathbf{x})^2 = 3^2 = 9$$


$$D(\mathbf{y})^2 = 2^2 = 4$$

$$D(\mathbf{z})^2 = 1^2 = 1$$



# K-means++


Suppose we are given a black box that will generate a uniform random number between 0 and any **N**. How can we use this black box to select points with probability proportional to  $D(\mathbf{x})^2$ ?



$$D(\mathbf{x})^2 = 3^2 = 9$$

$$D(\mathbf{y})^2 = 2^2 = 4$$


$$D(\mathbf{z})^2 = 1^2 = 1$$



$$\begin{aligned} &= D(\mathbf{x})^2 + D(\mathbf{y})^2 \\ &\quad + D(\mathbf{z})^2 = 14 \end{aligned}$$

# K-means++


Suppose we are given a black box that will generate a uniform random number between 0 and any **N**. How can we use this black box to select points with probability proportional to  $D(\mathbf{x})^2$ ?



$$D(\mathbf{x})^2 = 3^2 = 9$$


$$D(\mathbf{y})^2 = 2^2 = 4$$

$$D(\mathbf{z})^2 = 1^2 = 1$$




# K-means++

Q3: the black box returns "12" as the random number generated. Which point do we choose for the next center (x, y, or z) ?



# K-means++

Q4: the black box returns "4" as the random number generated. Which point do we choose for the next center (x, y, or z) ?





# K-means++


What happens if the black box can only generate numbers between 0 and 1?


# Kmeans Quizz (take 2)












































# How to choose the right $k$ ?

1. Iterate through different values of  $k$  (elbow method)



# How to choose the right $k$ ?

1. Iterate through different values of  $k$  (elbow method)
2. Use empirical / domain-specific knowledge  
Example: Is there a known approximate distribution of the data? (K-means is good for spherical gaussians)
3. Metric for evaluating a clustering output



# Evaluation

Recall our goal: Find a clustering such that

- **Similar** data points are in the **same cluster**
- **Dissimilar** data points are in **different clusters**

# Evaluation

Recall our goal: Find a clustering such that

- **Similar** data points are in the **same cluster** ✓
- **Dissimilar** data points are in **different clusters**


# Evaluation

K-means cost function tells us the within-cluster distances between points will be small overall.

But what about the intra-cluster distance? Are the clusters we created far?  
How far? Relative to what?


## Discuss - 5min

Define a metric that evaluates how spread out the clusters are from one another.




a: average within-cluster distance

b: average intra-cluster distance




a: average within-cluster distance  
b: average intra-cluster distance

What does it mean for  $(b - a)$  to be 0?




a: average within-cluster distance  
b: average intra-cluster distance

What does it mean for  $(b - a)$  to be large?




The value of  $(b-a)$  doesn't mean much by itself. Can we compare it to something so that the ratio becomes a value between 0 and 1?






$$(b - a) / \max(a, b)$$





What does it mean for  $(b - a) / \max(a, b)$  to be close to 1?



What does it mean for  $(b - a) / \max(a, b)$  to be close to 0?

# Silhouette Scores

For each data point  $i$ :  
 $a_i$ : mean distance from point  $i$  to every other point in its cluster




# Silhouette Scores

For each data point  $i$ :

$a_i$ : mean distance from point  $i$  to every other point in its cluster

$b_i$ : smallest mean distance from point  $i$  to every point in another cluster





# Silhouette Scores

For each data point  $i$ :

$a_i$ : mean distance from point  $i$  to every other point in its cluster

$b_i$ : smallest mean distance from point  $i$  to every point in another cluster

$$s_i = (b_i - a_i) / \max(a_i, b_i)$$





# Silhouette Scores







$$s_i = (b_i - a_i) / \max(a_i, b_i)$$

Silhouette score plot

OR

return the mean  $s_i$  over the entire dataset as a measure of goodness of fit







# K-means Variations

1. K-medians (uses the  $L_1$  norm / manhattan distance)
2. K-medoids (any distance function + the centers must be in the dataset)
3. Weighted K-means (each point has a different weight when computing the mean)