

Bachelor thesis

# RIGHT-ANGLED COXETER GROUPS ARE RFRS

NICLAS RIST

May 16, 2024

Prof. Dr. Claudio Llosa Isenrich

Department of Mathematics

Karlsruhe Institute of Technology

Niclas Rist: *Right-angled Coxeter groups are RFRS*, © May 16, 2024

SUPERVISOR: JProf. Dr. Claudio Llosa Isenrich

LOCATION:  
Karlsruhe, Germany

TIME FRAME:  
Novembre 11, 2023 - May 16, 2024

# CONTENTS

1	Introduction	1
I	Preliminaries	
2	Preliminaries	5
2.1	Coxeter Groups . . . . .	5
2.2	Representation of Coxeter groups . . . . .	6
2.3	The Tits cone - An example . . . . .	10
2.4	The word metric and the faithful representation . . . . .	11
2.5	The fundamental domain and Stabilizers . . . . .	14
2.6	Covering actions - A Toolbox . . . . .	18
3	The main theorem	21
3.1	The RFRS property . . . . .	21
3.2	Construction of the manifold cover . . . . .	22
3.3	Some orbifold theory . . . . .	23
3.4	The cofinal cover . . . . .	25
3.5	The cofinal manifold sequence . . . . .	28
3.6	Loops bouncing off faces . . . . .	29
	Bibliography	35



The theory of 3-dimensional manifolds is far richer and much more complex than that of the more familiar 2-dimensional case of surfaces. While a significant milestone was achieved in 2002 with the completion of the 3-manifold classification by Grigori Perelman, many questions about their more detailed structure remain open. In his work on 3-manifolds, kleinian groups and hyperbolic manifolds [7], Thurston proposed 24 open questions that could guide future research in this area. In his paper [1], Ian Agol tackled question 18 of Thurston, namely whether every hyperbolic 3-manifold virtually fibers, i.e. has a finite-sheeted cover that fibers over  $S^1$ . Many authors gave reason to hope for a positive answer to this question, indeed Thurston himself wrote that ‘this dubious-sounding question seems to have a definite chance for a positive answer’. Since the proposal of this question, many 3-manifolds have been shown to virtually fiber but only few criteria for when a manifold virtually fibers were established. In many cases, the cover that fibers over  $S^1$  was exhibited fairly explicit, for example in Walsh’s work on two-bridge knots, Montesinos knots and link complements [8].

Agol introduces a more abstract condition on the fundamental group of a 3-manifold, called the RFRS property. In the following of his work, he then proves that if an irreducible 3-manifold has a fundamental group, satisfying the RFRS property, is virtually fibered.

With this context, we want to allude to what this work is about and how it relates to the theory of 3-manifolds. We will concentrate on one theorem in Agols’ work [1, Criteria for virtual fibering], stating that every right-angled Coxeter group virtually is RFRS. This result is interesting, as the group of right-angled Coxeter groups is in some sense ‘nice enough’ to satisfy the RFRS property, yet big enough to contain, or be related to, many other classes of groups. One example are the surface groups, as they are subgroups of the reflection group in a right-angled pentagon and thus, subgroups of a right-angled Coxeter group. Another example are the right-angled Artin groups, that are by definition closely related to right-angled Coxeter groups. By a result of Michael Davis and Tadeusz Januszkiewicz [3], right-angled Artin groups are commensurable with right-angled Coxeter groups and thus, it can be shown that they are also RFRS. This is also contained as a corollary in Agols’ work.

This work consists of precisely two parts. In the first half, i.e. in Chapter 2, we will start by introducing Coxeter groups, as well as right-angled Coxeter groups and will construct the representation, which consists basically of a vector space with a special bilinear form. Then, we will see that the dual space is the natural space to look for a chamber in the representation of a Coxeter group and use it to define the fundamental chamber, as well as the Tits cone. In Section 2.3 we will give an explicit example of the Tits cone, as it can be thought of quite geometrically.

In the following sections, we will prove a row of important results about Coxeter groups, starting with the proof that the representation of Coxeter groups is indeed a faithful representation. We will further work out, that the so called fundamental chamber is a fundamental domain of the group action via the representation, as well as how stabilizers of points look like. Finally, in

the last section of Chapter 2, we will put together a toolbox, concerning Covering theory and connecting it to group actions, as this will be important in the proof of the main theorem of this work.

As some sort of guidance for the reader through Chapter 3, we want to outline the ideas of the sections in it in the following as well. First, we introduce the RFRS property on groups, we already mentioned above. In what follows, we will discuss what we have to show to verify this property for a group  $G$ .

Building onto the developed understanding of the RFRS property, clearly, we want to proceed by finding a candidate for the finite index subgroup  $W'$  of  $W$  that will satisfy the RFRS condition. This candidate will induce a manifold cover of the fundamental chamber  $C$ , corresponding to  $W$  and will be introduced in Section 3.2.

Continuing from there on, the general goal is to translate from a cofinal sequence of manifold covers of the fundamental chamber  $C$ , to a sequence of groups. To achieve this, there is no way around some orbifold theory, as the fundamental chamber  $C$  carries a natural orbifold structure. Thus, Section 3.3 will cover some aspects of the theory of orbifolds.

In Section 3.4 we will use the developed tools of the theory of orbifolds, to construct a cofinal sequence of orbifold covers, beginning in the fundamental chamber  $C$ . This sequence will in particular induce a cofinal sequence of subgroups, which we want to be a sequence of subgroups of the finite index subgroup  $W'$ .

This will be dealt with in Section 3.5, where the aforementioned sequence will somewhat be forced to be a sequence of subgroups in  $W'$ . We will see that this new cofinal sequence of groups induces a sequence of manifold covers, starting in the cover established in Section 3.2.

Up to then, we will have produced a cofinal sequence of subgroups living in  $W'$  and it remains to show that the rationally derived subgroup at step  $i$ , is a subgroup of the following subgroup at step  $i + 1$ . This will be the content of the last Section 3.6, where we will watch loops bouncing off the faces in an orbifold. We will see 'how' specifically loops behave in the orbifolds, given by polytopes  $P_i$ , which will then allow us to deduce the main theorem.

## Part I

### PRELIMINARIES





# 2 | PRELIMINARIES

## 2.1 COXETER GROUPS

First of all, we define the main object of interest we want to study.

**Definition 2.1.1.** Let  $S$  be a set consisting of elements  $s_i$  indexed by an index set  $I$  with  $|I| = n < \infty$ . Let  $(m_{ij}) = M$  be a symmetrical matrix in  $(\mathbb{N} \cup \{\infty\})^{n \times n}$ , where  $m_{ii} = 1$  for all  $i$  and  $m_{ij} \geq 2$  for  $i \neq j$ . Define a group  $W$  via the following presentation:

$$W := \langle S \mid (s_i s_j)^{m_{ij}} = 1 \text{ for all } i, j \in I \rangle,$$

where the relations with  $m_{ij} = \infty$  are usually omitted, i.e. give trivial relations. The pair  $(W, S)$  is called a Coxeter System and  $M$  is called the corresponding Coxeter Matrix. A Coxeter group is a group isomorphic to a group  $W$ , corresponding to a Coxeter System  $(W, S)$ . It is generated by the set  $S$ .

In this work we will be particularly interested in a special class of Coxeter groups that we call right-angled. They are defined as follows, by imposing significant constraints on the entries of the Coxeter matrix.


**Definition 2.1.2.** A Coxeter System  $(W, S)$  is right-angled if, for all distinct  $i, j \in I$ , the condition  $m_{ij} \in \{2, \infty\}$  is satisfied. In this context, the group  $W$  is then called a right-angled Coxeter group (RACG).

We give some important examples of Coxeter groups as well as right-angled Coxeter groups.

- Example 2.1.3.**
1. Dihedral groups,  $D_{2m} \cong \langle s_1, s_2 \mid s_1^2 = s_2^2 = 1, (s_1 s_2)^m = 1 \rangle$  are Coxeter groups for all  $m \in \mathbb{N}$ .
  2. The triangle groups,  $\Delta(l, m, n) \cong \langle r, s, t \mid r^2 = s^2 = t^2 = (rs)^l = (st)^m = (tr)^n = 1 \rangle$  with  $l, m$  and  $n$  integers greater or equal to 2 are Coxeter groups.
  3. The infinite Dihedral group,  $D_\infty \cong \langle s, t \mid s^2 = t^2 = 1 \rangle$  is a right-angled Coxeter group.
  4. The free product,  $\mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z} \cong \langle r, s, t \mid r^2 = s^2 = t^2 = 1 \rangle$  is a right-angled Coxeter group.

In Figure 2.1, small portions of the combinatorial Cayley graphs of these examples are depicted.

Figure 2.1: The combinatorial Cayley graph of ...



We define a special type of subgroups, called parabolic subgroups within a Coxeter group  $W$ . These subgroups are constructed from a subset of the index set  $I$ . The definition is as follows:

**Definition 2.1.4.** Let  $(W, S)$  be a Coxeter System as above with finite index set  $I$ , and  $J$  be a subset of the index set  $I$ . The group  $W_J := \langle \{s_j \mid j \in J\} \rangle$ , generated by the  $s_j$  in  $S$  with  $j \in J$ , is then called a parabolic subgroup of  $W$ . Moreover, we call any conjugate of  $W_J$  a parabolic subgroup as well.

Once we have constructed the representation of Coxeter groups on a vector space as well as the Tits cone in the coming section, the parabolic subgroups will be a useful tool to form a deeper understanding of these objects. We will extensively use them in Sections 2.4 and 2.5.

## 2.2 REPRESENTATION OF COXETER GROUPS

Given a Coxeter System  $(W, S)$ , let  $V$  be a real vector space with basis  $\{e_1, \dots, e_n\}$ , where  $n = |I| = |S|$ . This provides a natural identification,  $GL_n(V) \cong GL_n(\mathbb{R})$ . We define a bilinear form  $B_W$  on  $V$  as follows:

$$B_W(e_i, e_j) := \begin{cases} -\cos\left(\frac{\pi}{m_{ij}}\right) & , m_{ij} < \infty \\ -1 & , m_{ij} = \infty \end{cases}.$$

By Definition 2.1.1, it is assured that  $m_{ij} \geq 2$  for distinct  $i, j$ , ensuring the cosine term is non-positive. Consequently, we have  $B_W(e_i, e_j) \leq 0$  for distinct  $i, j$ . Furthermore, from  $m_{ii} = 1$ , it follows that  $B_W(e_i, e_i) = 1$ . Using this bilinear form, we define hyperplanes with corresponding reflections for each basis element  $e_i$  as follows:

$$H_i := \{v \in V \mid B_W(e_i, v) = 0\}, \quad \sigma_i : V \rightarrow V, \quad v \mapsto v - 2B_W(e_i, v)e_i.$$

**Theorem 2.2.1.** The map given by:

$$\rho : W \rightarrow GL_n(V) \cong GL_n(\mathbb{R}), \quad s_i \mapsto \sigma_i$$

is an injective homomorphism and therefore a faithful representation of  $W$ .

Before we prove the homomorphism property, we recall: A map  $\varphi : S \rightarrow G$  from a set  $S$  to a group  $G$  extends to a homomorphism  $\hat{\varphi} : \langle S \mid R \rangle \rightarrow G$ , if and only if the induced homomorphism  $\overline{\varphi} : F_S \rightarrow G$  from the free group over  $S$  satisfies  $\overline{\varphi}(r) = 1_G$  for every  $r \in R$ .

*Proof.* Observe that  $\sigma_i^2 = \text{id}$  in  $\text{GL}_n(V)$  and thus to prove that  $\rho$  is a homomorphism, we apply the above to our situation to see that it suffices to show that the product  $\sigma_i \sigma_j$  has order  $m_{ij}$  in  $\text{GL}_n(V)$  for distinct  $i, j \in I$ . Also note that in the case of  $m_{ij} = \infty$  there is nothing to prove as these relations are defined to be trivial in the presentation of  $W$ .

Thus, consider the two-dimensional subspace  $V_{ij}$  of  $V$  spanned by two basis vectors  $e_i, e_j$  and take a general element  $v = \lambda \cdot e_i + \mu \cdot e_j$  in  $V_{ij}$  with  $\lambda, \mu \in \mathbb{R}$  not simultaneously zero. As  $m_{ij} < \infty$ , the bilinear form  $B_W$  is positive definite by the following calculation

$$B_W(v, v) = \lambda^2 - 2\lambda\mu \cos\left(\frac{\pi}{m_{ij}}\right) + \mu^2 = \left(\lambda - \mu \cos\left(\frac{\pi}{m_{ij}}\right)\right)^2 + \mu^2 \sin^2\left(\frac{\pi}{m_{ij}}\right) > 0.$$

Identify  $(V_{ij}, B_W|_{V_{ij}})$  with the euclidean plane  $(\mathbb{R}^2, \langle \cdot, \cdot \rangle)$  up to a change of basis. Then  $\sigma_i$ , resp.  $\sigma_j$  act by orthogonal reflections in their corresponding hyperplanes  $H_i$ , resp.  $H_j$  intersected with  $V_{ij}$ . By having a look at the inner product of  $e_i$  and  $e_j$

$$B_W(e_i, e_j) = -\cos\left(\frac{\pi}{m_{ij}}\right) = \cos\left(\pi - \frac{\pi}{m_{ij}}\right),$$

we observe that the angle between the two in  $V_{ij}$  is precisely  $\pi - \frac{\pi}{m_{ij}}$ . We conclude from this that the angle between their reflecting lines in  $\frac{\pi}{m_{ij}}$  and the composition  $\sigma_i \sigma_j$  turns out to be a rotation about  $\frac{2\pi}{m_{ij}}$ . As the composition  $\sigma_i \sigma_j$  fixes the orthogonal complement  $V_{ij}^\perp$  of  $V_{ij}$  by definition of the  $\sigma_i$ , we see that it has order  $m_{ij}$  on the whole space  $V$ .

This proves that  $\rho$  extends to a homomorphism. It remains to show the injectivity of  $\rho$ , which will be a consequence of a bigger result in section 2.4, see Corollary 2.4.4.  $\square$

In contrast to the above proof, in the case of  $m_{ij} = \infty$ , we cannot expect our bilinear form to be positive definite. Let  $v \in V_{ij}$  be as in the proof, then

$$B_W(v, v) = \lambda^2 - 2\lambda\mu + \mu^2 = (\lambda - \mu)^2 \geq 0.$$

This shows that we can at least expect it to be positive semidefinite. The direct calculation

$$(\sigma_i \sigma_j)(e_i) = \sigma_i(e_i + 2e_j) = e_i + 2(e_i + e_j),$$

together with an inductive argument, shows that  $(\sigma_i \sigma_j)^n(e_i) = e_i + 2n(e_i + e_j)$ . Therefore, the composition  $\sigma_i \sigma_j$  has infinite order on  $V_{ij}$  and in particular on the whole of  $V$ . The following remark is a consequence of this, together with the observation in above proof.

**Remark 2.2.2.** *From last paragraph and the proof of Theorem 2.2.1, we conclude that two-generator subgroups of Coxeter groups are dihedral. Either of order  $2m_{ij}$  or infinite order.*

We want to extend the action of  $W$  to the dual of the vector space  $V$ . This is achieved by acting on  $V^*$  via the dual representation of  $\rho$ , which we define by

$$\rho^* : W \rightarrow \text{GL}_n(V^*), \quad w \mapsto (\rho^*(w) : V^* \rightarrow V^*, \varphi \mapsto \rho^*(w)(\varphi)), \quad w \in W, \varphi \in V^*.$$

We can evaluate the functional  $\rho^*(w)(\varphi)$  on some  $v \in V$  via  $(\rho^*(w)(\varphi))(v) := \varphi(\rho(w^{-1})(v))$ . Notation wise, we will simply write  $w(v)$ , when  $w \in W$  acts on  $v \in V$  via  $\rho(w)(v)$ . Similarly, we write  $w(\varphi)$  when we mean that  $w \in W$  acts on some element  $\varphi \in V^*$  of the dual space, via the dual representation  $\rho^*(w)(\varphi)$ . As in the case of the vector space  $V$ , we want to give a definition for the notion of a hyperplane with corresponding reflection in the dual space  $V^*$  as well. By dual hyperplane, we mean a subspace  $H_i^* := \{\varphi \in V^* \mid \varphi(e_i) = 0\}$ , and the corresponding dual reflections will be a map from  $V^*$  to  $V^*$ , given by:

$$\sigma_i^* : V^* \rightarrow V^*, \quad \varphi \mapsto \varphi \circ \sigma_i = \varphi - 2B_W(e_i, \cdot) \varphi(e_i).$$

To further explore Coxeter groups and their action via this representation, we need some more notation. In particular, we want a so-called *chamber*. This should be thought of as a cone over a polytope with finitely many faces such that the reflections in its codimension one faces correspond to the generators of  $W$  under the representation.

**Definition 2.2.3.** *The fundamental chamber  $C$  of the dual representation is the set, given by*

$$C := \{\varphi \in V^* \mid \varphi(e_i) \geq 0 \ \forall i \in I\} \subset V^*.$$

Denote by  $\{e_1^*, \dots, e_n^*\}$  the dual basis of  $V^*$  corresponding to the standard basis  $\{e_1, \dots, e_n\}$  of  $V$ . Then we calculate, using the  $\sigma_i^*$  from above:

$$\sigma_i^*(e_j^*) = e_j^* - 2B_W(e_i, \cdot) e_j^*(e_i) = \begin{cases} e_j^* & \text{for } i \neq j \\ e_j^* - 2B_W(e_j, \cdot) & \text{for } i = j \end{cases},$$

which implies that each reflection  $\sigma_i^*$  fixes all the hyperplanes  $H_j^*$ , for distinct indices  $i$  and  $j$ . Moreover, note that the fundamental chamber can be written in the form

$$C = \bigcap_{i \in I} \{\varphi \in V^* \mid \varphi(e_i) \geq 0\} = \bigcap_{i \in I} (H_i^* \cup \{\varphi \in V^* \mid \varphi(e_i) > 0\}),$$

where we observe that the sets  $H_i^* \cap C$  form the pairwise distinct codimension one faces of the chamber  $C$ . The open halfspaces  $\{\varphi \in V^* \mid \varphi(e_i) > 0\}$  in the latter term will be called  $A_i^*$  and using these, we define the open fundamental chamber to be the intersection of the open halfspaces:

$$\text{int}(C) = \mathring{C} = \bigcap_{i \in I} A_i^*.$$

As mentioned above, we want to study the action of our Coxeter group via the dual representation, acting by reflection in the faces  $H_i^*$ . However, in general the translates of the chamber under the group action won't cover the whole of  $V^*$ , which motivates the following definition:

**Definition 2.2.4.** *The Tits cone is the union of all  $W$ -translates of the chamber,  $WC := \bigcup_{w \in W} wC \subset V^*$ .*

As the name suggests, the fundamental chamber is a fundamental domain for the action of  $W$  on its Tits cone  $WC$  under the dual representation  $\rho^*$ . This will be proved in Theorem 2.5.7. While the formal definition of the Tits cone provides a rigorous foundation, it is not very insightful from a geometric perspective. As one can think about the Tits cone quite geometrically, especially in low dimensions, we will take a closer look at an explicit example in the following section. Before doing so, we end this section with the following remark.

**Remark 2.2.5.** *One may ask why we transport everything to the dual space, instead of working in the standard representation  $\rho$ . For this, consider the infinite dihedral group  $D_\infty \cong \langle s, t \mid s^2 = t^2 = 1 \rangle$ . We fix a basis  $\{e_1, e_2\}$  of  $V$  and obtain that the bilinear form in this basis is given by the matrix*

$$B_W = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}.$$

*We observe that  $H_1 = \text{span}\{e_1 + e_2\} = H_2$ , which implies that  $\sigma_1$  and  $\sigma_2$  fix the same hyperplane, despite having different  $(-1)$ -Eigenspaces (namely the span of  $e_1$ , resp.  $e_2$ ). Therefore, in general, working in the standard representation won't result in a chamber, giving rise to the existence of the Tits cone. Now, passing to the dual space  $V^*$  by fixing the dual basis  $\{e_1^*, e_2^*\}$ , consider the dual reflections  $\sigma_i^*$  as discussed before. By the more general calculation earlier, we obtain*

$$\sigma_1^*(H_2^*) = H_2^* \quad \text{and} \quad \sigma_2^*(H_1^*) = H_1^*.$$

*And furthermore, we note that for all  $i, j \in \{1, 2\}$*

$$\sigma_i^*(B_W(e_j, \cdot)) = B_W(e_j, \cdot) - 2B_W(e_i, \cdot)B_W(e_j, e_i) = -B_W(e_j, \cdot),$$

*using that  $B_W(e_j, \cdot) = -B_W(e_i, \cdot)$ . This shows that both dual reflections have the same  $(-1)$ -Eigenspace, but fix different hyperplanes (i. e., have different  $(+1)$ -Eigenspaces), resulting in a chamber as wished. We conclude with the following picture.*




Figure 2.2: The chamber of  $D_\infty$

### 2.3 THE TITS CONE - AN EXAMPLE

As an example we take a closer look at the free product  $W \cong \langle r, s, t \mid r^2 = s^2 = t^2 = 1 \rangle$ , a right-angled Coxeter group from Example 2.1.3. We fix the basis  $\{e_1, e_2, e_3\}$  and identify  $V$  with  $\mathbb{R}^3$ . In this basis, the bilinear form  $B_W$  is given by the matrix

$$B_W = \begin{pmatrix} 1 & -1 & -1 \\ -1 & 1 & -1 \\ -1 & -1 & 1 \end{pmatrix}.$$

By the spectral theorem we find a basis of orthonormal Eigenvectors, in which  $B_W$  is a diagonal matrix with its eigenvalues as entries. Using the Gram-Schmidt procedure, we get

$$\begin{pmatrix} -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \end{pmatrix} \cdot \begin{pmatrix} 1 & -1 & -1 \\ -1 & 1 & -1 \\ -1 & -1 & 1 \end{pmatrix} \cdot \begin{pmatrix} -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ 0 & \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

In other words, we have  $V^T B_W V = D$  with  $V \in O(n)$  and  $D = \text{diag}(2, 2, -1)$ . Now, since we have a diagonal matrix, we can multiply the entries by squares, since the resulting matrix will be congruent to the given one:

Let  $A = \text{diag}(\mu_1, \dots, \mu_n) \in \mathbb{R}^{n \times n}$ ,

$$S = \text{diag}(\lambda_1, \dots, \lambda_n) \in (\mathbb{R} \setminus \{0\})^{n \times n} \implies S^T A S = \text{diag}(\lambda_1^2 \mu_1, \dots, \lambda_n^2 \mu_n).$$

To apply this and further transform our matrix  $D$ , define the invertible matrix  $T$  as follows

$$T = \begin{pmatrix} \frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

which then implies  $\tilde{D} := T(V^T B_W V)T = \text{diag}(1, 1, -1)$ . The images of the basis vectors  $\{e_1, e_2, e_3\}$  are given by the three columns of the matrix  $TV^T$ , namely:

$$\tilde{e}_1 = \begin{pmatrix} -\frac{1}{2} \\ -\frac{\sqrt{3}}{6} \\ \frac{1}{\sqrt{3}} \end{pmatrix}, \tilde{e}_2 = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{pmatrix} \text{ and } \tilde{e}_3 = \begin{pmatrix} \frac{1}{2} \\ -\frac{\sqrt{3}}{6} \\ \frac{1}{\sqrt{3}} \end{pmatrix}.$$

Note that we have  $V \cong \mathbb{R}^3$ , equipped with the inner product

$$\langle x, y \rangle_{2,1} := x^T \tilde{D} y = x_1 y_1 + x_2 y_2 - x_3 y_3.$$

Since  $\langle \tilde{e}_i, \tilde{e}_i \rangle_{2,1} = 0$  for all  $i \in \{1, 2, 3\}$ , we have that these three vectors span an ideal triangle in the hyperboloid model of  $\mathbb{H}^2$ , given by  $\langle x, y \rangle_{2,1} = -1$ . One gets the ideal triangle by intersecting the hyperboloid with the hyperplanes spanned by each two of the  $\tilde{e}_i$  (they will only intersect in the surrounding cone of the hyperboloid).

Given these new coordinates under the transformation  $TV^T$ , the Tits cone will be given by  $x_1^2 + x_2^2 - x_3^2 < 0$  union the images of  $\tilde{e}_1, \tilde{e}_2$  and  $\tilde{e}_3$  under the reflection in the sides of the chamber, i.e. in the sides of the ideal triangle. Moreover, we get a subgroup of  $O(2, 1)_+$  generated by the following three matrices

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{5}{3} & -\frac{4}{3} \\ 0 & \frac{4}{3} & \frac{5}{3} \end{pmatrix}, \begin{pmatrix} -1 & \frac{2}{\sqrt{3}} & -\frac{2}{\sqrt{3}} \\ \frac{2}{\sqrt{3}} & \frac{1}{3} & \frac{2}{3} \\ \frac{2}{\sqrt{3}} & -\frac{2}{3} & \frac{5}{3} \end{pmatrix} \text{ and } \begin{pmatrix} -1 & -\frac{2}{\sqrt{3}} & \frac{2}{\sqrt{3}} \\ -\frac{2}{\sqrt{3}} & \frac{1}{3} & \frac{2}{3} \\ -\frac{2}{\sqrt{3}} & -\frac{2}{3} & \frac{5}{3} \end{pmatrix}.$$

Each of the above matrices corresponds to one of the generators  $r, s$  and  $t$  of the right angled Coxeter group  $W$  under the transformation  $TV^T$ , constructed above.




Figure 2.3: The fundamental chamber in the Poincaré disc.

## 2.4 THE WORD METRIC AND THE FAITHFUL REPRESENTATION

Recall that for any finitely generated group  $W = \langle S \rangle$ , its Cayley graph induces a metric on  $W$ , relative to the generating set  $S$ . We call it the *word metric* of  $W$  relative to  $S$  and denote it by  $d_S$ . Also note that the word metric is left-invariant, meaning that for group elements  $u, v, w \in W$ , we have the equality  $d_S(uv, uw) = d_S(v, w)$ . Now, define the *word length* of an element  $w \in W$  to be  $\ell(w) := d_S(w, 1_W)$ , the distance of an element to the neutral element of the group. Note that  $\ell(w) = 0$  if and only if  $w = 1_W$ .

**Lemma 2.4.1.** *We collect some properties of the length function we will use later on.*

1.  $\forall w \in W : \ell(w) = \ell(w^{-1})$
2.  $\forall s \in S : \ell(s) = 1 \text{ and } \ell(w) = 1 \iff w \in S^{\pm 1}$
3.  $\forall v, w \in W : \ell(vw) \leq \ell(v) + \ell(w)$
4.  $\forall v, w \in W : \ell(v) - \ell(w) \leq \ell(vw)$
5.  $\forall w \in W, s \in S^{\pm 1} : \ell(w) - 1 \leq \ell(ws) \leq \ell(w) + 1$

*Proof.* All of the above statements follow from the fact that  $d_S$  is a left-invariant metric.  $\square$

Coming back to Coxeter groups, by definition each generator  $s \in S$  has order 2 in  $W$ . Therefore, we can write every non-trivial element in  $W$  as a sequence of generators in  $S$ . Note that in this sequence there might be redundancies, so that the following definition makes sense. We call an expression  $w = s_{i_1} \cdots s_{i_r}$  for  $i_1, \dots, i_r \in I$  and  $r \in \mathbb{N}$  *reduced*, if  $\ell(w) = r$ , i.e.  $w$  cannot be represented by a shorter word. These reduced expressions have the caveat of not being unique by any means.

Given a parabolic subgroup  $W_J$  of a Coxeter group  $W$ , it admits its own word metric with respect to  $J \subset I$ . Therefore each parabolic subgroup admits its own length function, which we denote by  $\ell_J(w)$  for words  $w$  in  $W_J$ . In the following we will make use of the general fact that we have  $\ell(w) \leq \ell_J(w)$  for all  $w \in W_J$ .

The length function turns out to be an important and powerful tool in studying Coxeter groups. Indeed, its role will be demonstrated in several of the forthcoming proofs, beginning with the following theorem which is a key step in proving faithfulness of our previously defined representation. But before doing so, we want to motivate the next theorem. We consider the set  $\{w(e_i) \mid i \in I, w \in W\}$ . Note that we fixed a basis in our representation vector space and therefore each of these elements can be written as a linear combination of the basis vectors. We will use this to define ‘positive’ and ‘negative’ elements. An element  $w(e_i)$  is said to be

- positive (denoted  $w(e_j) > 0$ ), if  $w(e_j) = \sum_{i \in I} \lambda_i e_i$  with  $\lambda_i \geq 0$  for all  $i \in I$ ,
- negative (denoted  $w(e_j) < 0$ ), if  $w(e_j) = \sum_{i \in I} \lambda_i e_i$  with  $\lambda_i \leq 0$  for all  $i \in I$ .

**Theorem 2.4.2.** *Let  $w \in W$  and  $s_i \in S$  for  $i \in I$ . Then  $\ell(ws_i) > \ell(w)$  implies that  $w(e_i) > 0$ .*

*Proof.* The base case is trivial, as  $\ell(w) = 0$  implies  $w = 1_W$  and thus  $w$  fixes every basis element. Therefore, assume  $w$  is non-trivial and in reduced form. We state the induction hypothesis.

**(IH)** Let  $v \in W$ , so that  $\ell(v) < \ell(w)$  and  $\ell(vs_i) > \ell(v)$ , then we have  $v(e_i) > 0$ .

**Claim 1:** There is a  $j \in I$  such that  $s_i \neq s_j$  and  $\ell(ws_j) = \ell(w) - 1$ .

*Proof of Claim 1.* Since  $w$  is in reduced form, write  $w = s_{i_1} \cdots s_{i_k}$  and set  $s_j = s_{i_k}$ . This implies

$$\ell(ws_i) > \ell(w) > \ell(w) - 1 = \ell(ws_j),$$

and in particular  $s_j \neq s_i$ . This proves the first Claim. ■

Consider the parabolic subgroup  $\langle s_i, s_j \rangle \leq W$ , generated by these two distinct elements in  $S$ . Set  $J := \{i, j\}$  in  $I$  and denote the subgroup by  $W_J$ . We use the decomposition  $W = W/W_J \cdot W_J$  to decompose  $w$  into two parts, each living in one component. For this, consider a specific subset of the coset  $wW_J$ , given by

$$A := \{v \in W \mid vW_J = wW_J \text{ and } \ell(v) + \ell_J(v^{-1}w) = \ell(w)\}.$$

By definition,  $w \in A$ . With regard to the following, we choose  $v \in A$  such that its length  $\ell(v)$  is minimal. Now set  $v_J = v^{-1}w$ , which is equivalent to writing  $w = v v_J$ , giving us a decomposition of  $w$  as desired. Due to this decomposition, our analysis of  $w$  now boils down



to studying the separate actions of  $v$  and  $v_J$  on  $V$ . First note that  $ws_j$  is contained in  $A$  as well. Clearly,  $s_j^{-1} = s_j$  and thus we have that

$$s_j w^{-1} w = s_j \in W_J \text{ and } \ell(ws_j) + \ell_J(s_j) = \ell(w) - 1 + 1 = \ell(w).$$

By the choice of  $\ell(v)$  to be minimal, we have  $\ell(v) \leq \ell(ws_j) = \ell(w) - 1$ , implying  $\ell(v) < \ell(w)$ . Hence, we are almost set up to apply the induction hypothesis to  $v$  and  $s_i$ . The last ingredient missing to do so is the following Claim.

**Claim 2:** For the lengths of  $vs_i$  and  $v$ , we have the relation:  $\ell(vs_i) \geq \ell(v)$ .

*Proof of Claim 2.* Assume towards contradiction:  $\ell(vs_i) < \ell(v)$ , equivalently  $\ell(vs_i) = \ell(v) - 1$ . Then:

$$\begin{aligned} \ell(w) &= \ell(vv_J) = \ell(vs_i s_i v^{-1} w) \leq \ell(vs_i) + \ell(s_i v^{-1} w) \\ &\leq \ell(v) - 1 + \ell_J(v^{-1} w) + 1 = \ell(v) + \ell_J(v^{-1} w) = \ell(w). \end{aligned}$$

This means equality holds throughout and in particular  $\ell(w) = \ell(vs_i) + \ell_J(s_i v^{-1} w)$ . But this implies  $vs_i$  belongs to  $A$ , contradicting the minimality of  $\ell(v)$ . Thus, the claim holds. ■

Applying the induction hypothesis **(IH)** to  $v$  and  $s_i$  leaves us with  $v(e_i) > 0$ . The exact same argument applied to  $v$  and  $s_j$  shows  $v(e_j) > 0$ , so that we omit this here. Due to the decomposition and the fact that  $w$  lies in  $W_J$ , we are now reduced to the rank two situation. As we have seen that all two generator subgroups are dihedral, we work in the flat plane. But first, we have to observe the following claim.

**Claim 3:** For  $v_J s_i$  and  $v_J$ , we have the relation:  $\ell_J(v_J s_i) \geq \ell(v)$ .

*Proof of Claim 3.* Assume towards contradiction, that  $\ell_J(v_J s_i) < \ell(v_J)$  holds. Then:

$$\ell(ws_i) = \ell(vv_J s_i) \leq \ell(v) + \ell(v_J s_i) \leq \ell(v) + \ell_J(v_J s_i) < \ell(v) + \ell_J(v_J) = \ell(w),$$

which contradicts the assumption of the theorem, that  $\ell(ws_i) > \ell(w)$ . ■

Moreover, this shows that every reduced expression of  $v_J$  in the parabolic subgroup  $W_J$  has to end in  $s_j$ . Otherwise, we would have  $\ell(v_J) > \ell(v_J s_i)$ , which contradicts *Claim 3*. To deduce the theorem, it suffices to show the following claim.

**Claim 4:**  $(s_i s_j)(e_i) = \lambda_i e_i + \lambda_j e_j$  with  $\lambda_i, \lambda_j \geq 0$ , implying  $(s_i s_j)(e_i) > 0$ .

*Proof of Claim 4.* We already know that  $W_J$  is dihedral, either of order  $2m_{ij}$  or infinite order. As pointed out above, every reduced expression of  $v_J$  - an alternating product of  $s_i$  and  $s_j$  - in  $W_J$  has to end in  $s_j$ . Using this, we distinguish the two cases.

- 1)  $m_{ij} < \infty$ :  $W_J$  is dihedral of order  $2m_{ij}$ . Note that the maximum length of  $w \in W_J$  is precisely  $m_{ij}$  (as the Cayley graph is a cycle of length  $2m_{ij}$ ) and the element of maximal length is represented by the reduced expressions  $s_i s_j \cdots s_i s_j$  and  $s_j s_i \cdots s_j s_i$ . This implies that  $v_J$  has to have length strictly smaller  $m_{ij}$ , else it would have a reduced expression ending in  $s_i$ , contradicting the above. Then  $v_J$  is a product of less than  $\frac{m_{ij}}{2}$  terms  $s_i s_j$ , and each of the products  $s_i s_j$  is a rotation about  $\frac{2\pi}{m_{ij}}$  towards  $e_j$ .

Using that  $e_i$  and  $e_j$  are at angle  $\pi - \frac{\pi}{m_{ij}}$ , we see that  $v_J$  rotates  $e_i$  at most about  $\pi - \frac{2\pi}{m_{ij}}$

towards  $e_j$ . Thus,  $v_j(e_i)$  still lies inside the cone spanned by  $e_i$  and  $e_j$ . It could very likely be that we have another reflection  $s_j$ , but as the angle between  $e_i$  and the reflecting line is  $\frac{\pi}{2} - \frac{\pi}{m_{ij}}$ , the resulting vector still lies in the cone.

- 2)  $m_{ij} = \infty$ : In the case of the infinite dihedral group, we have previously seen that  $(s_i s_j)^n(e_i) = e_i + 2n(e_i + e_j) = (2n + 1)e_i + 2ne_j$ . Thus,  $\lambda_i, \lambda_j \geq 0$  with  $|\lambda_i - \lambda_j| = 1$ .

This proves the last Claim. ■

Now that we have established that  $w = vv_J$  holds, for the element  $w(e_i)$  we have

$$w(e_i) = (vv_J)(e_i) = v(\lambda_i e_i + \lambda_j e_j) = \lambda_i \underbrace{v(e_i)}_{>0} + \lambda_j \underbrace{v(e_j)}_{>0}.$$

Claim 4 implies  $\lambda_i, \lambda_j \geq 0$ , allowing us to conclude  $w(e_i) > 0$ . □

The result of the previous theorem readily extends to its converse, offering valuable insights as well. We shall formally record this implication as a corollary.

**Corollary 2.4.3.** *In the previous theorem, set  $w = \tilde{w}s_i$  for  $i \in I$ . Then  $\ell(\tilde{w}s_i) < \ell(w)$  implies  $\ell(\tilde{w}s_i s_i) > \ell(\tilde{w}s_i)$ . Thus we have  $(ws_i)(e_i) = -w(e_i) > 0$ , or equivalently  $w(e_i) < 0$ .*

With Corollary 2.4.3 and Theorem 2.4.2 in hand, we will finally be able to deduce the injectivity and thus faithfulness of our representation  $\rho$ .

**Corollary 2.4.4.** *The homomorphism of Theorem 2.2.1 is injective and thus a faithful representation.*

*Proof.* Assume  $w \in \ker\{\rho\}$  non-trivial. Then there is an  $i \in I$  with corresponding  $s_i \in S$ , such that  $\ell(ws_i) < \ell(w)$ . Now Corollary 2.4.3 implies  $w(e_i) < 0$ , but  $w(e_i) = e_i > 0$  as  $\rho(w) = \text{id}_V$ , which is a contradiction and thus, the statement holds. □

## 2.5 THE FUNDAMENTAL DOMAIN AND STABILIZERS

In this section, we will prove that the fundamental chamber  $C \subset V^*$  is indeed a fundamental domain for the action of  $W$  on its Tits cone  $WC$ . Moreover, we will work out how stabilizers of points look like and then show that the Tits cone is really a convex cone in the dual space  $V^*$ .

We start by translating some results of the previous section to the language of chambers and halfspaces. The following two lemmas capture this essence.

**Lemma 2.5.1.** *Let  $w \in W$  and  $i \in I$ . The relation  $\ell(s_i w) > \ell(w)$  is equivalent to saying that  $w$  leaves the open chamber  $\mathring{C}$  inside the open halfspace  $A_i^*$ , corresponding to the generator  $s_i$ . Put more formally, this states that the set  $w(\mathring{C})$  is a subset of the open halfspace  $A_i^*$ .*

*Proof.* Observe that  $\ell(s_i w) > \ell(w)$  is equivalent to  $\ell(w^{-1} s_i) > \ell(w^{-1})$ . Then by Theorem 2.4.2, we have that  $w^{-1}(e_i) > 0$ . Take an arbitrary point  $\varphi \in \mathring{C}$  in the open fundamental chamber, then we have that  $w(\varphi)(e_i) = \varphi(w^{-1}(e_i))$ . Since the chamber  $C$  lies in  $A_i^*$  for all  $i \in I$  and  $\varphi \in \mathring{C}$ , we see that  $w(\varphi) > 0$  is equivalent to  $w^{-1}(e_i) > 0$ . But this is true by assumption and since  $\varphi$  is arbitrary, we conclude that  $w(\mathring{C}) \subset A_i^*$ .  $\square$

Of course this lemma also has a corresponding converse result as in the previous section.

**Lemma 2.5.2.** *Let  $w \in W$  and  $i \in I$ . The relation  $\ell(s_i w) < \ell(w)$  is equivalent to saying that  $w$  now moves the open chamber  $\mathring{C}$  inside the open halfspace, complementary to  $A_i^*$ . More formally, this states that the set  $w(\mathring{C})$  is a subset of the open halfspace  $s_i(A_i^*)$ , as the  $s_i$  permute  $A_i^*$  and its corresponding complementary halfspace.*

*Proof.* Take  $w' = s_i w$  for some  $i \in I$  and  $w \in W$  so that  $\ell(s_i w) > \ell(w)$ . Then we have that the length of  $s_i w' = w$  is strictly smaller than the length of  $w'$ . By applying above lemma we get that  $w(\mathring{C}) \subset A_i^*$ , which is equivalent to  $s_i w'(\mathring{C}) \subset A_i^*$ . Apply  $s_i$  to both sides leaves us with  $w'(\mathring{C}) \subset s_i(A_i^*)$ , which proves the lemma.  $\square$

Building onto this insight, we now want to study the action of parabolic subgroups on the Tits cone to get an understanding of the stabilizers of points. To do so, decompose the fundamental chamber into subsets corresponding to the parabolic subgroups of  $W$  as follows.

**Definition 2.5.3.** *Given a parabolic subgroup  $W_J$  corresponding to  $J \subset I$ , set*

$$C_J := \bigcap_{j \in J} H_j^* \cap \bigcap_{k \notin J} A_k^*.$$

*We call these the corresponding parabolic subsets (of the fundamental chamber).*

**Example 2.5.4.** 1. *When the set  $J$  is empty, the corresponding parabolic subset  $C_\emptyset$  coincides with the entire chamber  $C$ . Conversely, when  $J$  contains all indices,  $C_J$  reduces to the singleton  $\{0\}$ .*

2. *If  $J$  is a proper subset of  $I$  with cardinality one, then the corresponding subset  $C_J$  coincides with a codimension-one face of the chamber  $C$ .*

**Theorem 2.5.5.** *Let  $w \in W$  and  $J, K \subset I$  be subsets. Then  $w(C_J) \cap C_K \neq \emptyset$  implies  $J = K$ ,  $w \in W_J$  and thus  $w(C_J) = C_J$ . In particular, the isotropy groups of the sets  $C_J$  are the parabolic subgroups  $W_J$ .*

*Proof.* Let  $w \in W$  and  $J, K \subset I$  be subsets, such that  $w(C_J) \cap C_K \neq \emptyset$ . The proof is by induction on the length of  $w$ . The base case  $\ell(w) = 0$  is trivial, as then  $w$  is trivial.

Assume that  $\ell(w) > 0$  and choose  $i \in I$ , such that  $\ell(s_i w) < \ell(w)$ . Writing  $w = s_i(s_i w)$ , by Lemma 2.5.2 we know that  $w$  moves the open chamber  $\mathring{C}$  into the open halfspace  $s_i(A_i^*)$ , i.e.  $w(\mathring{C}) \subset s_i(A_i^*)$ . Now using the continuity of the group action, we note that  $w(C) \subset \overline{s_i(A_i^*)}$ . Recall that by definition, the fundamental chamber  $C$  lies in the halfspaces  $\overline{A_i^*}$  for all  $i \in I$ . Thus, we record that  $w(C) \cap C \subset H_i^*$  and since  $s_i$  fixes the corresponding  $H_i^*$  by definition, it

fixes every point in the intersection of  $C$  and its translate  $w(C)$ . Note that the sets  $C_J$  and  $C_K$  are subsets of the fundamental chamber  $C$  and therefore,  $s_i$  fixes every point in the non-empty set  $w(C_J) \cap C_K$ . But if  $s_i$  fixes some point  $\varphi$  in  $C_K$ , we calculate

$$\varphi(e_i) = s_i(\varphi)(e_i) = \varphi(s_i(e_i)) = -\varphi(e_i) \implies \varphi(e_i) = 0 \iff \varphi \in H_i^*.$$

From what we can deduce that  $i \in K$ , respectively  $s_i \in W_K$ . Using this together with the assumption, we get that  $s_i w(C_J) \cap C_K = s_i(w(C_J) \cap C_K)$  is non-empty. We apply the induction hypothesis to the element  $s_i w$ , to see that  $J = K$  and  $s_i w \in W_J$ . Finally, since  $s_i \in W_J = W_K$ , we have that  $s_i w \in W_J$  implies  $w \in W_J$ , proving the theorem.  $\square$

Before proceeding with proving that the fundamental chamber lives up to its name, we want to clearly state what is meant by a fundamental domain.

**Definition 2.5.6.** Let  $G$  be a group, acting on a topological space  $X$ . We call a closed subset  $F \subset X$  a fundamental domain, if for each  $x \in F$  its orbit  $\text{Orb}(x)$  intersects  $F$  in exactly one point.

Note that, by definition of the Tits cone  $WC$ , every  $W$ -orbit of a point  $\varphi \in C$  meets the fundamental chamber  $C$  in at least one point, namely  $\varphi$ . Thus, it suffices to prove that each  $W$ -orbit meets  $C$  in at most one point, to prove the following theorem.

**Theorem 2.5.7.** The fundamental chamber is a fundamental domain for the action of the Coxeter group  $W$  on its Tits cone  $WC$ , justifying its name.

*Proof.* Assume that  $\varphi, \psi \in C$  lie in the same  $W$ -orbit, but in different parabolic subsets  $C_J$ , respectively  $C_K$  of the fundamental chamber. Since they lie in the same orbit, there is a  $w \in W$  with  $\varphi = w(\psi)$ . Thus, the intersection  $w(C_J) \cap C_K$  is non-empty and Theorem 2.5.5 implies equality of  $J$  and  $K$ , as well as  $w \in W_J$ . We deduce  $\varphi = w(\psi) = \psi$ . Thus, every  $W$ -orbit of a point  $\varphi \in C$  meets the fundamental chamber  $C$  at most in  $\varphi$ , proving the theorem.  $\square$

Define a set  $\mathcal{C}$  as the union of all translates of possible parabolic subsets  $C_J$  i. e., define  $\mathcal{C}$  by

$$\mathcal{C} := \bigcup_{J \subset I} \bigcup_{w \in W/W_J} w(C_J).$$

We want to emphasize here that by Theorem 2.5.5 the sets of the form  $w(C_J)$  in the Tits cone  $WC$  are all disjoint for different  $J \subset I$  and  $w$  ranging over the coset  $W/W_J$ . Thus, the sets of  $\mathcal{C}$  form a partition of the Tits cone. This decomposition (although not into chambers) is a key component in the following theorem.

**Theorem 2.5.8.** The Tits cone  $WC$  is a convex cone, and every closed line segment in the Tits cone meets only finitely many of the sets in  $\mathcal{C}$ .

*Proof.* First note that the fundamental chamber is a convex cone as the intersection of the finitely many closed halfspaces  $\overline{A_i^*}$ . This implies that the Tits cone is a cone as well. We will prove the convexity by showing that every closed segment between any two points in the Tits

cone is contained in it. Furthermore, we will prove that these segments can be covered by finitely many of the sets in the above defined union  $\mathcal{C}$ , implying the latter statement.

Consider the closed segment  $[\varphi, \psi]$  with  $\varphi, \psi \in WC$  and without loss of generality, assume the endpoints lie in different chambers  $C$  and  $w(C)$  for  $w \in W$ . Proceed by induction on the word length  $\ell(w)$ . The base case  $\ell(w) = 0$  reduces to  $\varphi, \psi \in C$ . Since  $C$  is convex and can trivially be covered by finitely many of the  $C_J$ , this case has been dealt with.

Therefore, we now assume  $\ell(w) > 0$ . Intersect the segment  $[\varphi, \psi]$  with the fundamental chamber  $C$ , to receive two new segments  $[\varphi, \xi]$  and  $[\xi, \psi]$  (cf. Figure 2.4a). The first segment can be covered by finitely many of the sets in  $\mathcal{C}$ , as it lies inside the fundamental chamber  $C$ . Thus, we need to show that we can cover the second segment  $[\xi, \psi]$  by finitely many of these sets. Assume further, we have a  $J \subset I$ , such that  $\psi \in s_i(A_i^*)$  for an  $i \in J$  and  $\psi \in \overline{A_i^*}$  for all  $i \notin J$ . Then we have that  $\psi \notin C$ .

**Claim:**  $\xi$  has to lie in one of the codimension-one faces  $H_i^*$ ,  $i \in J$ .

*Proof of Claim.* Assume that  $\xi$  lies in the open halfspace  $A_i^*$  for some  $i \in J$ . Then every point  $\zeta$  in the intersection of a neighborhood of  $\xi$ , contained in  $A_i^*$ , with the segment  $[\xi, \psi]$  has to also lie in  $A_i^*$ . Clearly,  $\zeta \in \overline{A_i^*}$  for  $i \notin J$  holds as well, implying that  $\zeta \in C$ . But this is a contradiction to the decomposition of the initial segment  $[\varphi, \psi]$ . Thus,  $\xi$  has to lie in one of the  $H_i^*$ . ■

Using the assumptions  $\psi \in s_i(A_i^*)$  and  $\psi \in w(C)$ , we deduce  $w(C) \subset \overline{s_i(A_i^*)}$ , hence by continuity of the action  $w(\overset{\circ}{C}) \subset s_i(A_i^*)$ . By Lemma 2.5.2 this is equivalent to  $\ell(s_i w) < \ell(w)$  and we are set up to apply the induction hypothesis to  $\xi$  and  $s_i(\psi) \in s_i w(C)$ . This produces a cover of  $[\xi, s_i(\psi)]$  by finitely many sets in  $\mathcal{C}$ . But since we established that  $\xi$  has to lie in  $H_i^*$ , translation by  $s_i$  gives  $[s_i(\xi), s_i^2(\psi)] = [\xi, \psi]$ , and thus we can cover this segment by finitely many sets as well. The result follows. □




Figure 2.4: A schematic picture to the proof.

To summarize, the essence of this section is that first of all the fundamental chamber is indeed a fundamental domain of the action of a Coxeter group on its Tits cone. This was shown in Theorem 2.5.7. Furthermore, we want to emphasize that each point in the interior of the fundamental chamber has trivial stabilizer, which is a direct consequence of Theorem 2.5.5. Moreover, every point in a parabolic subset  $C_J$  of  $C$ , for  $J$  a subset of  $I$ , is stabilized by the corresponding parabolic subgroup  $W_J$ .

## 2.6 COVERING ACTIONS – A TOOLBOX

In this section we will assemble a set of tools, mostly coming from algebraic topology, connecting group actions to the notion of coverings. They will be somewhat essential in the proof of the main theorem in the next chapter. We start by saying what is meant by a covering.

**Definition 2.6.1.** *Let  $X, Y$  be topological spaces. A continuous map  $p : Y \rightarrow X$  is a covering map and  $Y$  a covering space for  $X$ , if every point  $x$  in  $X$  has an open neighborhood  $U$ , such that the preimage of  $U$  under  $p$  is a disjoint union of open sets  $U_i$  in  $Y$  for an index set  $I$ . Furthermore, the map  $p$  has to be a local homeomorphism, meaning the restriction of  $p$  to the  $U_i$  is a homeomorphism onto its image  $p(U_i)$ . Then,  $U$  is called evenly covered and  $|I|$  the degree of the covering, while the open sets  $U_i$  are called the sheets over  $U$  and the preimage of an  $x$  in  $X$  is called a fiber of  $x$ .*

It turns out that restricting the action of a group on a topological space in the right way will give coverings by quotienting out the action. We make this precise in the following.

**Definition 2.6.2.** *Let  $G$  be a group, acting on a space  $X$ . The action is said to be properly discontinuous, if every point in  $X$  has a neighborhood  $U$  such that the set  $\{g \in G \mid g(U) \cap U \neq \emptyset\}$  is finite.*

**Definition 2.6.3.** *Let  $G$  be a group, acting on a space  $X$ . The action is said to be a covering (space) action if every point has a neighborhood  $U$  such that the set  $\{g \in G \mid g(U) \cap U \neq \emptyset\}$  consists only of the neutral element.*

Obviously, this last condition is even more restrictive. Yet, at least for Hausdorff spaces, there is a connection between the two. In addition to acting properly discontinuously, we have to demand the action to be free.

**Lemma 2.6.4.** *Let  $G$  be a group that acts freely and properly discontinuously on a Hausdorff space  $X$ . Then the action of  $G$  is a covering action in the above sense.*

*Proof.* Let  $X$  be a Hausdorff space and  $G$  be a group acting freely, properly discontinuously on  $X$ . Then for an open neighborhood  $U$  of  $x \in X$ , the set  $M := \{g \in G \mid gU \cap U \neq \emptyset\}$  is finite. For these  $g \in M$  we pick pairwise disjoint Neighborhoods  $V_g$  of  $gx$ , which is possible since  $X$  is Hausdorff and  $G$  acts freely (thus  $gx \neq x$ ). Finally, set  $V = \left(\bigcap_{g \in M} g^{-1}V_g\right) \cap U$ , which is open as finite intersection of open sets and by definition a neighborhood of  $x$ .  $\square$

To finally construct coverings from group actions and connect them in some sense, we need a last condition on the space the group acts on.

**Definition 2.6.5.** *A path-connected topological space  $X$  is called simply connected, if  $\pi_1(X) \cong \{1\}$ .*

Given a group  $G$  acting on a space  $X$ , we will denote by  $\text{Orb}_G(x)$  the orbit of a point  $x \in X$  under the action of the group  $G$ , i.e.  $\text{Orb}_G(x) := \{gx \mid g \in G\} \subset X$ .

**Lemma 2.6.6.** *Let  $G$  be a group, acting by a covering action on a simply connected topological space  $X$ . Then the quotient map  $p_G : (X, x_0) \rightarrow (G \backslash X, \text{Orb}(x_0))$  is a covering map and  $\pi_1(G \backslash X, \text{Orb}(x_0)) \cong G$ .*

*Proof.* Let  $U$  be an open neighborhood of  $x \in X$ , such that  $\{g \in G \mid gU \cap U \neq \emptyset\} = \{1\}$ .

**Claim 1:** The map  $p_G$  restricted to  $U$  is a continuous bijection onto its image  $p_G(U)$ .

*Proof of Claim 1.*

- Continuity:  $V \subset G \backslash X$  is open if and only if  $p_G^{-1}(V)$  is open and thus  $p_G$  is continuous.
- Surjectivity: Since orbits of points are non-empty,  $p_G$  is surjective.
- Injectivity: Assume  $x, y \in X$  distinct with  $p_G(x) = p_G(y)$ . But then there is a non-trivial element  $g \in G$ ,  $g \neq 1_G$  with  $gx = y$ , implying  $g(U) \cap U \neq \emptyset$ . This is a contradiction to  $G$  acting by a covering action. ■

**Claim 2:** Every point  $\text{Orb}(x) \in G \backslash X$  has a neighborhood  $V \subset G \backslash X$  with  $p_G^{-1}(V) = \bigsqcup_{g \in G} gU$ .

*Proof of Claim 2.* By assumption the sets  $gU$  are all disjoint neighborhoods of  $gx \in X$  for all  $g \in G$ . Moreover, for  $V := p_G(U) \subset G \backslash X$ , we have  $p_G^{-1}(V) = \bigsqcup_{g \in G} gU$ , proving the Claim. ■

**Claim 3:** The map  $p_G|_{gU} \rightarrow V = p_G(U)$  is a homeomorphism for all  $g \in G$ .

*Proof of Claim 3.* Note that the action of  $G$  on  $X$  is by homeomorphisms and  $U \subset X$  is open. Then the union  $\bigsqcup_{g \in G} gU$  is open as well and since  $p_G^{-1}(V) = \bigsqcup_{g \in G} gU$ , the set  $V$  is open in the quotient topology of  $G \backslash X$ . Therefore, the map  $p_G$  restricted to  $gU$  for  $g \in G$  is an open map and hence, by Claim 1 a homeomorphism. ■

Now all conditions in Definition 2.6.1 are satisfied, so  $p_G$  is a covering map. For the last part of the statement, note that  $X$  is simply connected, implying that it is the universal cover of  $G \backslash X$ . By definition,  $G$  is the group of Deck transformations, thus  $\pi_1(G \backslash X, \text{Orb}(x_0)) \cong G$ . □





# 3

## THE MAIN THEOREM

Let us state and recall the main theorem we want to prove.

**Theorem.** *A finitely generated right-angled Coxeter group  $W$  has a finite index subgroup  $W'$ , such that  $W'$  is residually finite and rationally solvable (RFRS).*

The goal of this whole chapter is to explain Agol's proof of this theorem, using the preliminaries and tools we have developed. We will break the proof down into smaller pieces and will discuss in each section a specific aspect of the proof.

### 3.1 THE RFRS PROPERTY

We should state clearly what we have to show. This is essentially the construction of a finite index subgroup, that is RFRS. But what does it mean for a group  $G$  to be RFRS? We shall start by giving the definition of the RFRS condition to shed some light onto our goal.

**Definition 3.1.1.** *Let  $G$  be a group. Then,  $G$  satisfies the RFRS condition, if there is a sequence of subgroups  $G = G_0 > G_1 > \dots$ , satisfying the following conditions.*

1. *for each  $i$ ,  $G_i \triangleleft G$  is a normal subgroup of  $G$ ,*
2. *for each  $i$ , the index of  $G_i$  in  $G$ ,  $[G : G_i]$  is finite,*
3. *the intersection  $\bigcap_i G_i = \{1\}$  is the trivial group, meaning the sequence is cofinal, and*
4. *for each  $i$ , the group  $(G_i)_r^{(1)} := \ker\{G_i \rightarrow \mathbb{Q} \otimes_{\mathbb{Z}} (G_i)_{ab}\}$  is a subgroup of  $G_{i+1}$ .*

Observe that in practice, it suffices to show that conditions 2 to 4 hold. To justify this, let  $G = G_0 > G_1 > \dots$  be a cofinal sequence of finite index subgroups  $G_i < G$  with  $(G_i)_r^{(1)} \leq G_{i+1}$ . Then, for each  $i$ , we pass to the core of  $G_i$  in  $G$ , which is defined by  $\text{Core}(G_i) := \bigcap_{g \in G} gG_i g^{-1}$ . By construction, the core of a subgroup  $G_i$  is normal in  $G$  and furthermore, we claim:

**Claim:** If  $G_i$  has finite index in  $G$ , its core will also have finite index in  $G$ .

*Proof of Claim.* Let  $G_i$  be a subgroup in  $G$  with  $[G : G_i] < \infty$ . Then,  $G$  acts (by left multiplication) on the left cosets of  $G_i$  in  $G$  and the kernel of this action consists of the elements  $h \in G$  with

$$hgG_i = gG_i \iff g^{-1}hgG_i = G_i \iff g^{-1}hg \in G_i \iff h \in gG_i g^{-1} \quad \forall g \in G.$$

We note that the kernel is exactly the core of  $G_i$ . Moreover, the quotient  $G/\text{Core}(G_i)$  embeds into the symmetric group  $\text{Sym}(G/G_i)$ , as the left action on the cosets permutes them. As the order of this symmetric group is  $[G : G_i]!$ , we see that if  $G_i$  has finite index in  $G$ , the core has finite index as well. ■

Thus, to see that this new sequence satisfies the RFRS condition, it remains to check that the rational derived subgroup  $(\text{Core}(G_i))_r^{(1)}$  is a subgroup of the group  $\text{Core}(G_{i+1})$ .

Clearly, for  $g \in G$  we have the equality  $g(G_i)_r^{(1)}g^{-1} = (gG_i g^{-1})_r^{(1)}$ , from which we obtain

$$(\text{Core}(G_i))_r^{(1)} \leq \bigcap_{g \in G} (gG_i g^{-1})_r^{(1)} = \bigcap_{g \in G} g(G_i)_r^{(1)}g^{-1} \leq \bigcap_{g \in G} gG_{i+1}g^{-1} = \text{Core}(G_{i+1}).$$

To summarize, by accepting to pass to the sequence of core subgroups in  $G$ , we can drop the first condition on the sequence of subgroups in the RFRS condition.

**Remark 3.1.2.** Note that an abelian group  $A$  has a natural structure as a  $\mathbb{Z}$ -module. Thus, the tensor product with the rationals is well-defined and tensoring  $A$  with  $\mathbb{Q}$  kills the torsion part of the abelian group  $A$ . Whence, we have that  $\mathbb{Q} \otimes_{\mathbb{Z}} A \cong \mathbb{Q} \otimes_{\mathbb{Z}} A/\text{Torsion}$ . To see this, take an elementary tensor  $r \otimes a$  in  $\mathbb{Q} \otimes_{\mathbb{Z}} A$  so that  $a$  is a torsion-element of order  $n$ . Then we get the following equalities,

$$r \otimes a = r \cdot \frac{n}{n} \otimes a = r \cdot \frac{1}{n} \otimes n \cdot a = 0.$$

In particular, for the abelianization  $G_{ab}$ , the group  $\mathbb{Q} \otimes_{\mathbb{Z}} G_{ab}$  is isomorphic to  $\mathbb{Q} \otimes_{\mathbb{Z}} G_{ab}/\text{Torsion}$ . Moreover, the torsion-free abelianization embeds into  $\mathbb{Q} \otimes_{\mathbb{Z}} G_{ab}$  and the so called rational derived subgroup  $(G)_r^{(1)}$  is the kernel of the homomorphism from  $G$  to its torsion-free abelianization tensored with the rationals.

### 3.2 CONSTRUCTION OF THE MANIFOLD COVER

Consider the abelianization  $W_{ab}$  of  $W$ , which is isomorphic to  $(\mathbb{Z}/2\mathbb{Z})^n$ . Now, the abelianization yields a homomorphism  $\alpha : W \rightarrow W_{ab}$ . In the following we will focus on its kernel  $\ker \alpha$ . Note that by the first homomorphism theorem,  $\ker \alpha$  has finite index in  $W$ , since

$$|W/\ker \alpha| = |\mathbb{Z}/2\mathbb{Z}|^n = 2^n < \infty.$$

Here we used the fact that  $W$  is finitely generated, whence  $n = |I| = |S| < \infty$ .

Furthermore, note that for each  $J \subset I$  with  $W_J$  finite, we have an isomorphism between  $W_J$  and  $(\mathbb{Z}/2\mathbb{Z})^{|J|}$ . Thus, the restriction of  $\alpha$  to each such subgroup  $\alpha|_{W_J}$  is an injective homomorphism. We now use that in the right-angled case, the isotropy subgroups of codimension- $k$  faces are all of this form. This can be seen, as the isotropy subgroup of a codimension- $k$  face  $F$  is generated by the reflections in the  $k$  codimension-1 faces, whose intersection forms  $F$ . As all these codimension-1 faces meet at right-angles, the generators commute pairwise. Therefore, all isotropy subgroups inject into the abelianization of  $W$ . This implies that the intersection

of an isotropy subgroup with the kernel of  $\alpha$  is trivial and consequently no isotropy group is contained in the kernel  $\ker \alpha$ . Since finite subgroups are contained in isotropy subgroups, the kernel  $\ker \alpha$  acts freely on the Tits cone  $WC$  corresponding to  $W$ .

In particular, by Theorem 2.5.5 the action of  $W$  on the interior of its Tits cone  $\text{int}(WC)$  is properly discontinuous. By Theorem 2.5.8 the Tits cone is also a convex cone, implying that it has trivial fundamental group, whence is simply-connected. Having all this information, we are able to apply Lemma 2.6.6 to obtain that the following map is a covering

$$\text{int}(WC) \longrightarrow \text{int}(WC)_{/\ker \alpha} \quad x \mapsto \text{Orb}_{\ker \alpha}(x).$$

We conclude the results of this section in the following proposition.

**Proposition 3.2.1.** *Let  $W$  be a right-angled Coxeter group with corresponding Tits cone  $WC$ . Then, there is a finite index subgroup  $W' \leq W$ , acting by covering action on the interior of the Tits cone, implying that the quotient  $\text{int}(WC)_{/W'}$  is a manifold. It is given by  $W' = \ker\{W \rightarrow W_{ab}\} = \ker \alpha$ .*

### 3.3 SOME ORBIFOLD THEORY

In this section, we want to elaborate more on the natural orbifold structure of the fundamental chamber  $C$ . We start by giving a formal definition of an orbifold. We break the definition down into smaller pieces, starting with local models, sometimes called (orbifold) charts.

**Definition 3.3.1.** *A local model is a pair  $(\tilde{U}, \Gamma)$ , where  $\tilde{U} \subset \mathbb{R}^n$  is open and  $\Gamma$  is a finite subgroup of the group of diffeomorphisms of  $\tilde{U}$ , denoted  $\text{diffeo}(\tilde{U})$ , acting on  $\tilde{U}$ . By abusing notation, we will sometimes say that the quotient  $U = \tilde{U}_{/\Gamma}$  is the local model.*

Now that we have defined the local structure of an orbifold, we want to translate between these local models. This is being made precise by orbifold maps.

**Definition 3.3.2.** *An orbifold map between local models  $(\tilde{U}_i, \Gamma_i), (\tilde{U}_j, \Gamma_j)$  is a pair of maps  $(\tilde{\psi}, \varphi)$ , consisting of a smooth map  $\tilde{\psi} : \tilde{U}_i \rightarrow \tilde{U}_j$  and a homomorphism of groups  $\varphi : \Gamma_i \rightarrow \Gamma_j$ . We enforce the map  $\tilde{\psi}$  to be  $\varphi$ -equivariant, meaning that for all  $g \in \Gamma_i$  and all  $\tilde{x} \in \tilde{U}_i$ ,  $\tilde{\psi}(g\tilde{x}) = \varphi(g)\tilde{\psi}(\tilde{x})$  holds. Then  $\tilde{\psi}$  induces a map  $\psi : \tilde{U}_i_{/\Gamma_i} \rightarrow \tilde{U}_j_{/\Gamma_j}$ , between the local models. When all three of these maps are injective, we call  $\psi$  a local isomorphism.*

Now that we have these local definitions, we ‘glue’ them together, to obtain an orbifold. Before we do so, we recall some notions from topology.

Suppose, we are given an open cover  $\{U_i\}$  of a topological space  $X$ . It is said to be *locally finite*, if every  $x \in X$  admits a neighborhood  $N$  such that  $N \cap U_i$  is empty for all but finitely many of the indices  $i$ . The open cover  $\{U_i\}$  is a *refinement* of an open cover  $\{V_j\}$  of  $X$ , if for every  $V_j$  there is a  $U_i$  with  $U_i \subseteq V_j$ . Now, the topological space  $X$  is said to be *paracompact*, if every open cover of  $X$  admits such a locally finite refinement.

**Definition 3.3.3.** An  $n$ -dimensional (smooth) orbifold  $Q$  is a pair  $(X_Q, \mathcal{A})$ . The space  $X_Q$  is a paracompact Hausdorff space, called the underlying space. The set  $\mathcal{A}$  is called an orbifold atlas, consisting of charts  $(U_i, \phi_i)$ , indexed by some set  $I$  and satisfying the following conditions:

- the  $U_i$  form an open cover of the underlying space  $X_Q$ ,
- for each  $U_i$  there exists a local model  $\tilde{U}_i/\Gamma_i$  with a homeomorphism  $\phi_i : U_i \rightarrow \tilde{U}_i/\Gamma_i$  and
- charts have to be compatible, meaning that for  $U_i \subset U_j$  the inclusion is a local isomorphism.

To sketch the connection between manifolds and orbifolds, let us mention one more thing.


**Definition 3.3.4.** The local group  $\text{loc}(x)$  of some  $x$  in a local model  $\tilde{U}/\Gamma$  is the isotropy group of any  $\tilde{x}$  living in  $\tilde{U}$ , getting projected onto  $x$ . The singular locus  $\Sigma(Q)$  of an orbifold  $Q$  consists of all points in the underlying space  $X_Q$  with non-trivial local group, i.e.  $\Sigma(Q) = \{x \in X_Q \mid \text{loc}(x) \neq \{1\}\}$ .

By this definition, we see that an orbifold with empty singular locus is just a manifold. Furthermore, when thinking about an orbifold, we can just think about the underlying space and label each element in the singular locus by its local group.


Many basic examples arise by taking the quotient relative to a properly discontinuous group action on  $\mathbb{R}^n$ . We want to mention at least some examples.

**Example 3.3.5.** 1. Consider the flat plane  $\mathbb{R}^2$  with the action of a cyclic group  $\mathbb{Z}/n\mathbb{Z}$  by rotation about the origin. The arising orbifold is a cone with singular point the origin and cone angle  $\frac{2\pi}{n}$ .

2. Consider the sphere  $S^2$  with the action of a cyclic group  $\mathbb{Z}/n\mathbb{Z}$  by rotation about the north pole  $N$ . The arising orbifold now is called a teardrop with singular point, the north pole  $N$ .



(a) The cone orbifold.



(b) The teardrop orbifold.

The concept of orbifold coverings translates almost one to one from the ordinary case.

**Definition 3.3.6.** An orbifold covering  $p : Q' \rightarrow Q$  is a continuous map on the underlying spaces  $X_{Q'} \rightarrow X_Q$ , such that for each point  $x \in X_Q$  there is a local model  $U = \tilde{U}/\Gamma$  around  $x$  and each component  $V_i$  of  $p^{-1}(U)$  is homeomorphic to  $\tilde{U}/\Gamma_i$  for a subgroup  $\Gamma_i$  of  $\Gamma$ . Furthermore, the restriction  $p|_{V_i} : V_i \rightarrow U$  corresponds to the natural projection  $\tilde{U}/\Gamma_i \rightarrow \tilde{U}/\Gamma$ .

**Remark 3.3.7.** We want to mention here, that the universal cover of an orbifold is exactly what we think of, i.e. the initial object in the category of orbifold coverings. Moreover, there is a notion of orbifold fundamental groups, which we won't define here. For us it is only important to know that they behave to orbifold covers the same as fundamental groups to coverings in the ordinary case. In particular, subgroups (of index  $n$ ) of the orbifold fundamental group correspond to ( $n$ -sheeted) orbifold coverings.

Having all this in mind, we have two ways of approaching the fundamental chamber  $C$ . One way, it obtains its natural orbifold structure as the quotient  $\text{int}(WC)/W$ . As the Tits cone is simply connected, it is the universal cover of  $C$  and we observe that the orbifold fundamental group of  $C$  is precisely  $W$ . On the other hand, the fundamental chamber is the quotient of the manifold  $\text{int}(WC)/W'$  by a finite group action.

Since  $W'$  is a finite index subgroup of  $W$ , Remark 3.3.7 implies that we get a covering map  $p$  so that the diagram on the right commutes. Thus, both of the approaches to the fundamental chamber  $C$  agree in the sense that the projection map  $\text{int}(WC) \rightarrow \text{int}(WC)/W$  factors through the manifold cover, constructed in Section 3.1.

$$\begin{array}{ccc} \text{int}(WC) & \xrightarrow{\quad} & \text{int}(WC)/W \\ \downarrow & \nearrow p & \\ \text{int}(WC)/W' & & \end{array}$$

### 3.4 THE COFINAL COVER

In this section, we *sketch* the proof of the existence of a cofinal sequence of two-fold covers, arising by iteratively reflecting in faces. Some more results from the theory of orbifolds would be needed for the whole proof. Since they go beyond the scope of this work, we will state and use them without proof.

In the following, doubling along faces means the construction of a sequence of right-angled polytopes  $P_k = P_{k-1} \cup g_{k-1}(P_{k-1})$ , where  $g_{k-1}$  is the reflection in one of the faces of the previous polytope  $P_{k-1}$ , denoted by  $F_{k-1}$ . The following proposition will tell us, that in the case of a right-angled Coxeter group, we are able to double the fundamental chamber  $C$  in one of its faces and are still able to cover the whole Tits cone  $WC$  by reflecting in the faces of the resulting polytope.

**Proposition 3.4.1.** Let  $P_k \subset \mathbb{R}^n$  be a right-angled polytope with orbifold structure, coming from a doubling sequence starting in  $P_0 = C$ . Denote its orbifold fundamental group by  $\pi_1^{\text{orb}}(P_k)$  and its universal cover by  $\tilde{P}$ . Then, the fundamental group  $\pi_1^{\text{orb}}(P_{k+1})$  is isomorphic to the group generated by reflections in the faces of the double  $P_{k+1}$ . Furthermore, the fundamental group  $\pi_1^{\text{orb}}(P_{k+1})$  of the double is an index-2 subgroup of  $\pi_1^{\text{orb}}(P_k)$  and the following two conditions are satisfied.

1.  $\forall k \in \mathbb{N} : \forall g \in \pi_1^{\text{orb}}(P_k) : g(\tilde{P}_k) \cap \tilde{P}_k \neq \emptyset \implies g = 1$ , and
2.  $\forall k \in \mathbb{N} : \bigcup_{g \in \pi_1^{\text{orb}}(P_k)} g(P_k) = \tilde{P}$ .

*Proof.* We will omit this proof here, as it goes beyond the scope of this work. □

To state the other auxiliary lemma we will need, we first want to show that each chamber in the Tits cone can uniquely be labeled by elements of its corresponding group.

**Lemma 3.4.2.** *Let  $W$  be a Coxeter group and  $WC$  its Tits cone. The chambers of the form  $w(C)$  in the Tits cone can be labeled uniquely by elements in  $W$ .*

*Proof.* Let  $v(C), w(C)$  be chambers in  $WC$  for  $v, w \in W$ , such that  $v \neq w$  and  $w(C) = v(C)$ . But then, by Theorem 2.5.5 we have that

$$w(\mathring{C}) \cap v(\mathring{C}) = w(\mathring{C} \cap w^{-1}v(\mathring{C})) \neq \emptyset \implies w^{-1}v = 1 \iff w = v.$$

Thus, the chambers can be labeled in a unique way.  $\square$

**Remark 3.4.3.** *We remark that each of the right-angled polytopes  $P_k$  can be decomposed into a union of translates of the fundamental chamber  $C$ . This means for each polytope  $P_k$  there is a  $n_k \in \mathbb{N}$  and elements  $w_{i,k} \in W$ , so that  $P_k = \bigcup_{i=1}^{n_k} w_{i,k}(C)$ .*

We proceed with another helpful result, whose proof relies on the aforementioned results.

**Lemma 3.4.4.** *Let  $C \subset \mathbb{R}^n$  be the fundamental chamber of a Coxeter group  $W$ . The non-trivial labels  $w_{i,k} \neq 1_W$  of the chambers, defining the polytope  $P_k = \bigcup_{i=1}^{n_k} w_{i,k}(C)$ , are not contained in the reflection group, generated by reflecting in its faces.*

*Proof.* Note that by Lemma 3.4.2, the chambers of the form  $w(C)$  are uniquely labeled by the elements  $w \in W$ . Thus, let  $P_k := \bigcup_{i=1}^{n_k} w_{i,k}(C)$  be as in the lemma and  $w_{i,k}$  one of its defining labels. Assume,  $w_{i,k} \neq 1_W$  is contained in the reflection group generated by reflecting in the faces of  $P_k$ . But then we have that  $w_{i,k}(\mathring{P}_k) \cap \mathring{P}_k \neq \emptyset$ , contradicting Proposition 3.4.1.  $\square$

Using this insight, we define a graph  $G$  with vertices  $V(G) := \{w(C) \mid w \in W\}$  and edges  $E(G) := W \times S$ . The endpoint map is given by  $\delta : W \times S \rightarrow 2^{V(G)}$ ,  $(w, s) \mapsto (w(C), ws(C))$ . Thus, two vertices  $w(C)$  and  $v(C)$  have an edge, if there is an  $s \in S$ , such that  $ws = v$ . Then, clearly we have  $ws w^{-1}w(C) = ws(C) = v(C)$ . Note that the element  $ws w^{-1}$  is the only element, that flips the edge  $\{w(C), ws(C)\}$ . Thus, the map  $W \times S \rightarrow W$ ,  $(w, s) \mapsto ws w^{-1}$  defines a labeling of the edges by elements in  $W$ . We will call the combinatorial graph of  $G$  (meaning that every double edge gets collapsed) the *chamber graph*  $\text{Cham}(W, S)$  of  $W$ .

This graph is canonically isomorphic to the combinatorial Cayley graph of the group  $W$  and we thus see that the Cayley graph ‘embeds’ into the Tits cone  $WC$ .

**Theorem 3.4.5.** *There exists a cofinal doubling sequence  $P_k$ , coming from the fundamental chamber  $P_0 = C$ . This means that we have  $WC = \bigcup_{n \in \mathbb{N}} P_n$ .*

*Proof.* The proof is by induction on the word length  $\ell(w)$ . For  $\ell(w) = 0$ , the corresponding chamber is the fundamental chamber  $C$ , which is covered by itself.

(IH) The chambers with labels  $v$  of length  $\ell(v) = k$  are covered by a polytope  $P_k = \bigcup_{i=1}^{n_k} w_{i,k}(C)$  in the doubling sequence  $\{P_k\}_{k \in \mathbb{N}}$ .

We need to show that the chambers  $w(C)$  with labels  $w \in W$  of length  $\ell(w) = k + 1$  can be covered by doubling  $P_k$  finitely many times in faces. First observe that there is a label  $v$ , whose chamber  $v(C)$  is contained in  $P_k$  and that is connected to  $w$  by an edge in the Cayles graph. Since the Cayley graph sits inside the Tits cone and is isomorphic to the chamber graph, the chambers  $w(C)$  and  $v(C)$  have to be adjacent, sharing a face that is contained in a face of  $P_k$ . Then, we can distinguish two cases. Either,  $w(C)$  is already contained in  $P_k$  and we are done, or we double  $P_k$  in the face shared by  $v(C)$  and  $w(C)$ , so that  $w(C)$  is contained in  $P_{k+1}$ . Since  $W$  is finitely generated, there are finitely many more labels  $w' \in W$  of length  $\ell(w') = k + 1$  and we repeat the above procedure for each of them. Thus, every chamber in the Tits cone can be covered by doubling finitely many times in faces of the fundamental chamber  $C$ .  $\square$

**Remark 3.4.6.** Note that by Proposition 3.4.1 the elements corresponding to labels of chambers inside a right-angled polytope  $P_k$  are not contained in  $\pi_1^{orb}(P_k)$  and  $\pi_1^{orb}(P_{k+1})$  is a subgroup in  $\pi_1^{orb}(P_k)$  of index two. By Theorem 3.4.5 we have a doubling sequence  $\{P_k\}_{k \in \mathbb{N}}$  with  $\bigcup_{n \in \mathbb{N}} P_n = WC$ . As the fundamental group of  $WC$  is trivial (it is a convex cone), the intersection of the fundamental groups of the  $P_k$  is trivial as well. Thus, the cofinality of the doubling sequence translates to the group side.

**Example 3.4.7.** As an example, transporting the Tits cone of the free product from Example 2.1.3 to the poincaré disc, a portion of the chamber graph is depicted below.




Figure 3.1: The chamber graph of  $\mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z}$ .

### 3.5 THE COFINAL MANIFOLD SEQUENCE

We finally have developed enough tools, to construct the desired sequence of subgroups in  $W'$ . Following Theorem 3.4.5, we fix a cofinal doubling sequence  $P_k$ , starting in the fundamental chamber  $P_0 = C$  corresponding to the dual representation of  $W$ , i.e. we fix a cofinal sequence of two-fold orbifold covers  $C \leftarrow P_1 \leftarrow P_2 \leftarrow \dots$ . Then, we define the groups  $W_i := \pi_1^{\text{orb}}(P_i) \cap W'$  for  $i \in \{0\} \cup \mathbb{N}$ , each inducing a manifold via the quotient  $\text{int}(WC)/_{W_i}$ . Clearly, each  $W_i$  is a subgroup of  $W'$  as well as it is a subgroup of the orbifold fundamental group of the polytope (with orbifold structure)  $P_i$ . Using the theory of orbifold coverings, we observe that the construction of these manifolds implies that they are coverings of the polytopes  $P_i$  for each  $i$ . By Remark 3.4.6, the cofinal two-fold tower induces a descending series of fundamental groups of the polytopes  $P_i$ . Intersecting them with the finite index subgroup  $W'$  does not affect the cofinality and thus, the sequence of subgroups formed by the groups  $W_i$  is again cofinal.

We observe that to show that  $W'$  is RFRS, it remains to show the following two conditions:

1. each  $W_i := \pi_1^{\text{orb}}(P_i) \cap W'$  has finite index in the group  $W'$  and
2.  $(W_i)_r^{(1)} = \ker\{W_i \rightarrow \mathbb{Q} \otimes_{\mathbb{Z}} (W_i)_{\text{ab}}\}$  is a subgroup of  $W_{i+1}$ .

The first condition is essentially a consequence of the fact that the sequence of orbifold fundamental groups has index precisely two in each step. To conclude, we prove the following lemma.

**Lemma 3.5.1.** *The homomorphism  $\phi : W_i/W_{i+1} \rightarrow \pi_1^{\text{orb}}(P_i)/\pi_1^{\text{orb}}(P_{i+1})$  is injective.*

*Proof.* Suppose,  $g, h \in W_i/W_{i+1}$  with  $\phi(g) = \phi(h)$ , which is equivalent to

$$g \cdot \pi_1^{\text{orb}}(P_{i+1}) = h \cdot \pi_1^{\text{orb}}(P_{i+1}).$$

Since  $g$  and  $h$  lie in  $W_i/W_{i+1}$ , we have that  $gW_{i+1} = hW_{i+1}$ , equivalently  $g^{-1}h \in W_{i+1}$  from what we conclude that  $g = h$  in the quotient  $W_i/W_{i+1}$  and the result follows.  $\square$

An immediate consequence of this is the following result, we record here.

**Corollary 3.5.2.** *The index of  $W_{i+1}$  in  $W_i$  is less than or equal to 2.*

Based on this observation, we can conclude that the first of the above conditions holds for the constructed sequence of groups. We state this here as a corollary.

**Corollary 3.5.3.** *The index of  $W_i$  in  $W'$  is finite.*

*Proof.* By the above corollary, we have that the index of  $W_{i+1}$  in  $W_i$  is less than or equal to 2. Thus, for the index of  $W_i$  in  $W'$  we have the upper bound  $[W' : W_i] \leq 2^i < \infty$  for  $i \in \mathbb{N}$ .  $\square$



### 3.6 LOOPS BOUNCING OFF FACES

As discussed in the previous section, we need to show that the rational derived group  $(W_i)_r^{(1)}$  is a subgroup of  $W_{i+1}$ . First, recall the following result, addressing abelian quotients.

**Lemma 3.6.1.** *Let  $G$  be a group with normal subgroup  $N \triangleleft G$ . The quotient group  $G/N$  is abelian if and only if the derived subgroup  $G^{(1)} = [G, G]$  of  $G$  is contained in the normal subgroup  $N$ .*

*Proof.* ‘ $\Rightarrow$ ’ As  $N$  is normal in  $G$  and the quotient is abelian, for any  $g, h \in G$  we have that

$$ghN = gN \cdot hN = hN \cdot gN = hgN \iff g^{-1}h^{-1}gh \in N.$$

This is equivalent to the commutator  $[g, h]$  being in  $N$  for all  $g, h \in G$ , therefore  $[G, G] \subset N$ .

‘ $\Leftarrow$ ’ As now,  $[G, G] \subset N$  for any  $g \in G$  and  $n \in N$ , we have

$$gng^{-1} = gng^{-1}n^{-1}n = [g, n]n.$$

Since  $[G, G] \subset N$ , this element is again in  $N$  so that  $N$  is normal in  $G$ . To see that the quotient by  $N$  is abelian, we take  $g, h \in G/N$  and calculate

$$gN \cdot hN = ghN = hgg^{-1}h^{-1}ghN = hg[g^{-1}, h^{-1}]N = hgN = hN \cdot gN.$$

The result follows directly. □

**Lemma 3.6.2.** *Let  $G$  be a group and  $N \leq G$  a subgroup of index two. Then,  $N$  is normal in  $G$ .*

*Proof.* Let  $g$  be an element in  $G$ . As  $N$  has index two in  $G$ , we have that either  $g \in N$  or  $g \in G \setminus N$ . The first case is trivial, as  $gN = N = Ng$  and in the second case we have that  $gN = G \setminus N = Ng$ , since cosets partition the group  $G$ . This implies that  $N$  is normal in  $G$ . □

Using these two auxiliary lemmas, we can prove the following intermediate result.

**Proposition 3.6.3.** *The quotient group  $W_i/W_{i+1}$  is either isomorphic to  $\mathbb{Z}/2\mathbb{Z}$  or the trivial group. Furthermore, the quotient map  $W_i \rightarrow W_i/W_{i+1}$  factors through the abelianization and the derived subgroup  $(W_i)^{(1)}$  of  $W_i$  is a subgroup of  $W_{i+1}$ .*

*Proof.* The first part is a direct consequence of Corollary 3.5.2 and Lemma 3.6.2. Thus, the quotient  $W_i/W_{i+1}$  is in particular abelian and we are able to apply Lemma 3.6.1. This implies that the derived subgroup  $(W_i)^{(1)} = [W_i, W_i]$  is a subgroup of  $W_{i+1}$ . Stated differently, we have a factorization of  $W_i \rightarrow W_i/W_{i+1}$  through the abelianization  $(W_i)_{ab}$ . □

**Remark 3.6.4.** Recall that by Remark 3.1.2 torsion elements will not survive in the rational derived subgroup  $(W_i)^{(1)}_r$  of  $W_i$ , and by Proposition 3.6.3 we already have that  $(W_i)^{(1)}$  is a subgroup of  $W_{i+1}$ . We claim that it suffices to show that every element  $w \in W_i$  with non-trivial image in the quotient group  $W_i/W_{i+1}$  is not torsion in the abelianization  $(W_i)_{ab}$ . This is justified, as by Proposition 3.6.3, we have a factorization through the abelianization of  $W_i$ . Then, since  $w$  is not torsion in  $(W_i)_{ab}$ , it will admit a factorization through the torsion-free abelianization.

$$\begin{array}{ccc} W_i & \xrightarrow{\quad} & W_i/W_{i+1} \cong \mathbb{Z}/2\mathbb{Z} \\ & \searrow & \nearrow \text{dashed} \\ & (W_i)_{ab}/\text{Torsion} & \end{array}$$

In particular, the group  $W_i/W_{i+1}$  then is a quotient of the torsion-free abelianization  $(W_i)_{ab}/\text{Torsion}$ .


In the following we will denote the manifolds  $\text{int}(WC)/W_i$  by  $M_i$  for  $i \in \{0\} \cup \mathbb{N}$  and the projections to the orbifolds  $P_i$  by  $p_i : M_i \rightarrow P_i$ . Furthermore, we fix an element  $w$  that is contained in  $W_i$  but not in  $W_{i+1}$ , together with a loop  $\gamma$ , which represents the element  $w$  in the manifold  $M_i$ .

**Lemma 3.6.5.** The loop  $\gamma$  intersects the preimage  $p_i^{-1}(F_i)$  of the face  $F_i$ , in which  $P_i$  is doubled to obtain the polytope  $P_{i+1}$ , an odd number of times.

*Proof.* Project  $\gamma$  down to the polytope  $P_i$  generically, using the projection  $p_i$ . As the polytopes  $P_i$  have cyclic local group of order two everywhere on their codimension-one faces, the projection of  $\gamma$  gets reflected in the faces of the polytopes  $P_i$ . Now, consider the face  $F_i$  in which  $P_i$  gets doubled. Then,  $p_i(\gamma)$  hits this face an odd number of times, else  $p_i(\gamma)$  would lift to a loop in the double  $P_{i+1}$ , and therefore, the element  $w$  represented by  $\gamma$  would lift to the group  $W_{i+1}$ . This contradicts our choice of  $w$ , not being contained in  $W_{i+1}$ . In particular, observing the loop  $\gamma$  directly up in the manifold  $M_i$ , we see it intersecting the preimage  $p_i^{-1}(F_i)$  of the face  $F_i$  an odd number of times, which proves the lemma.  $\square$

One might think of the orbifolds  $P_i$  as polytopes that are formed by mirrors, and the loop  $\gamma$  as a closed path formed by a ray of light, being reflected in these mirrors. This interpretation also motivated the name of this section. Before proceeding, we want to provide a more explicit picture of the above argument. We record this in the following remark.


**Remark 3.6.6.** We want to explain what is happening here in a more explicit way. To do so, we look at the following picture, which one should think of in a schematic way. We see a loop  $\gamma$  (which, strictly speaking, is the projection of a loop in the manifold  $M_i$ ) in the polytope  $P_i$ , being reflected in the codimension-one faces of  $P_i$ . The face of interest  $F_i$  is colored in blue and we assume  $\gamma$  to be the orange path, walked around exactly once, starting in the basepoint  $x_0$ .



By doubling the polytope  $P_i$  along the face  $F_i$ , we land on the right side of this picture. Observe that the basepoint  $x_0$  now has a corresponding double, denoted by  $\tilde{x}_0$ . The more interesting part is, that now the loop  $\gamma$  gets unwound to a path going from  $x_0$  to  $\tilde{x}_0$ . If, on the other hand, we consider the loop  $\gamma^2$ , i.e. the loop  $\gamma$  gets run through twice, consequently hitting the face  $F_i$  twice, by our reasoning above, we should see a loop coming from  $\gamma^2$  in the two-fold cover  $P_{i+1}$ . Indeed, the loop  $\gamma^2$  lifts to a loop  $P_{i+1}$  and is pictured by the concatenated loop  $\tilde{\gamma} \cdot \gamma$  in  $P_{i+1}$  on the right.

Now that we have a more intuitive view on our reasoning, we have to deal with one more minor issue. We will mention it in the following remark, but will omit the details.

**Remark 3.6.7.** Note that we could also have a loop  $\gamma$  like in the left of the following image.



The difficulty with such loops is that the codimension-2 faces of a polytope also consist of singular points. Their local group may very well be different from the local groups of the points in codimension-1 faces. Nevertheless, we fortunately do not have to deal with them. In fact, we can always find a loop  $\gamma'$ , homotopy equivalent to  $\gamma$  and that is only hitting codimension-1 faces.

In the following, our goal will be to obtain a homomorphism of fundamental groups, from the fundamental group of the manifold  $\pi_1(M_i)$  to the fundamental group of the circle  $\pi_1(S^1)$ . This morphism will be induced by a continuous quotient map. To make this precise, we have to introduce one more concept, we have not yet mentioned. This is the fact that the preimage of a face  $F_i$  in the manifold  $M_i$  is two-sided (moreover it is oriented and embedded). This is a technical condition on the preimage  $p_i^{-1}(F_i) \subset M_i$ . As it will play an important role in the following, we give the definition of what it means to be two-sided.

**Definition 3.6.8.** Let  $M$  be a manifold. A submanifold  $F \subset M$  is called two-sided in  $M$ , if it locally looks like the product with an interval. More precisely, there is a neighborhood  $N_F$  with  $N_F \cong F \times (-\varepsilon, \varepsilon)$  for suitable  $\varepsilon > 0$ , where the submanifold  $F$  is identified with the fiber  $F \times \{0\}$ .


Denote by  $N_{F_i} \subset M_i$  the neighborhood coming from the two-sidedness of the preimage  $p_i^{-1}(F_i) \subset M_i$ . Using this neighborhood, we can construct the morphism in the next lemma.

**Lemma 3.6.9.** *There exists a continuous quotient map  $q : M_i \rightarrow S^1$  that induces a homomorphism of groups  $q_* : \pi_1(M_i) \rightarrow \pi_1(S^1)$ , between the fundamental groups.*

*Proof.* We will give an constructive proof. To do so, introduce an equivalence relation on the points in the manifold  $M_i$  as follows:

$$x \sim y \quad : \Longleftrightarrow \quad x, y \in M_i \setminus N_{F_i} \quad \text{or} \quad x = y.$$


Equip the quotient  $M_i / \sim$  with the quotient topology. Then, by passing to this quotient, we see a double cone with identified cone points.



Continuing from this space on, we have to pass to another quotient. Namely, we have to collapse each of the fibers  $p_i^{-1}(F_i) \times \{z\}$  for every  $z \in (-\varepsilon, \varepsilon)$  to a single point. This can be done by introducing another equivalence relation on the quotient space  $M_i / \sim$  by

$$x \sim_1 y \quad : \Longleftrightarrow \quad \exists z \in (-\varepsilon, \varepsilon) : x, y \in p_i^{-1}(F_i) \times \{z\}.$$


Equip this quotient again with the quotient topology and by passing to this further quotient, we are left with the interval  $(-\varepsilon, \varepsilon)$ , whose endpoints are identified.



As all the quotient spaces carry the quotient topology that is defined just right to make such constructions work, we only have to think about the red point. Consider the composed quotient map  $q : M_i \rightarrow S^1$  and a neighborhood  $U$  around the red point. Then,  $U$  is of the form  $[-\varepsilon, -\varepsilon + \delta] \cup (\varepsilon - \delta, \varepsilon]$  for some  $\delta \in (0, \varepsilon)$  and it is open in the quotient if and only if its preimage  $p_i^{-1}(U)$  is open in  $M_i$ . The preimage of  $U$  in  $M_i$  is of the form

$$q^{-1}(U) = M_i \setminus N_{F_i} \cup p_i^{-1}(F_i) \times [-\varepsilon, -\varepsilon + \delta] \cup p_i^{-1}(F_i) \times (\varepsilon - \delta, \varepsilon],$$

which is open, as its complement  $M_i \setminus q^{-1}(U) = p_i^{-1}(F_i) \times [-\varepsilon + \delta, \varepsilon - \delta]$  is closed in  $M_i$ .



Thus, the quotient map  $q : M_i \rightarrow S^1$  is continuous and therefore, it induces the desired homomorphism  $q_* : \pi_1(M_i) \rightarrow \pi_1(S^1)$  between the corresponding fundamental groups.  $\square$

We will now use the developed results to prove the following key-lemma. But before we do so, we want to add one more remark.

**Remark 3.6.10.** Note that in the proof of Lemma 3.6.5, we used the term ‘projecting generically’. We implicitly already used there that the preimage  $p_i^{-1}(F_i)$  is two-sided. The addendum ‘generically’ means that  $p_i : M_i \rightarrow P_i$  composed with the projection  $P_i \rightarrow (-\varepsilon, \varepsilon)$ , to the interval coming from the neighborhood  $p_i^{-1}(N_{F_i})$  in  $M_i$ , is actually injective in every neighborhood of an intersection of  $\gamma$  and  $F_i$ .

**Lemma 3.6.11.** The loop  $\gamma$  represents an element of infinite order in the abelianization  $(W_i)_{ab}$ .

*Proof.* By Lemma 3.6.5, we know that  $\gamma$  intersects the preimage of  $F_i$  in  $M_i$  an odd number of times. Thus, projecting  $\gamma$  down to  $S^1$ , using the continuous map  $q$  from Lemma 3.6.9, we see that the projection has to wind around the circle an odd number of times. For that reason, using the induced morphism  $q_*$ , we see that  $\gamma$  has to represent a non-trivial element in  $\pi_1(S^1)$ . But since  $\pi_1(S^1) \cong \mathbb{Z}$ , this immediately implies that  $\gamma$  represents an element of infinite order and moreover, since  $\mathbb{Z}$  is abelian, this implies that  $\gamma$  represents an element of infinite order in  $\pi_1(M_i)_{ab}$ , i.e.  $w$  has infinite order in  $\pi_1(M_i)_{ab} \cong (W_i)_{ab}$  as claimed.  $\square$

**Corollary 3.6.12.** The loop  $2\gamma$  in  $M_i$  represents a non-trivial element in the group  $\mathbb{Q} \otimes_{\mathbb{Z}} W_i / (W_i)_r^{(1)}$ .

*Proof.* By Lemma 3.6.11,  $\gamma$  represents an element of infinite order in the abelianization  $(W_i)_{ab}$ . In particular, this element cannot be contained in the rationally derived subgroup  $(W_i)_r^{(1)}$  and the loop  $2\gamma$  has to also represent an element of infinite order with this property. This implies that  $2\gamma$  represents an element that survives in the group  $\mathbb{Q} \otimes_{\mathbb{Z}} W_i / (W_i)_r^{(1)}$ , as claimed.  $\square$

**Corollary 3.6.13.** The finite index subgroup  $W'$  of the right-angled Coxeter group  $W$  is RFRS.

*Proof.* Since  $\gamma$  is a representative of the element  $w$ , we have that  $2\gamma$  is a representative of the element  $w^2$ . Thus, by Corollary 3.6.12  $w^2$  is not contained in the rational derived subgroup  $(W_i)_r^{(1)}$ . But by Corollary 3.5.2, the index of  $W_{i+1}$  in the group  $W_i$  is less than or equal to two, which implies that  $w^2$  has to be contained in  $W_{i+1}$ . Considering the factorization from Remark 3.6.4, we conclude that  $W_i / W_{i+1} \cong \mathbb{Z} / 2\mathbb{Z}$  is a quotient of the torsion-free abelianization  $(W_i)_{ab} / \text{Torsion}$ . In particular, we have finally established that every element mapping non-trivially to the group  $W_i / W_{i+1}$  is not torsion in the abelianization  $(W_i)_{ab}$ , and by the discussion in Remark 3.6.4, we can conclude that  $(W_i)_r^{(1)}$  is a subgroup of  $W_{i+1}$ . That was the only part missing for  $W'$  to be RFRS and therefore, we have finally proven the main theorem.  $\square$



## BIBLIOGRAPHY

- [1] Ian Agol. "Criteria for virtual fibering." In: *Journal of Topology* (2008).
- [2] Michael W. Davis. *The Geometry and Topology of Coxeter Groups*. (LMS-32) -. Kassel: Princeton University Press, 2012. ISBN: 978-1-400-84594-1.
- [3] Michael W. Davis and Tadeusz Januszkiewicz. "Right-angled Artin groups are commensurable with right-angled Coxeter groups." In: *Journal of Pure and Applied Algebra* 153.3 (2000), pp. 229–235. ISSN: 0022-4049. DOI: [https://doi.org/10.1016/S0022-4049\(99\)00175-9](https://doi.org/10.1016/S0022-4049(99)00175-9).
- [4] Allen Hatcher. *Algebraic topology*. Cambridge: Cambridge Univ. Press, 2002. ISBN: 0-521-79540-0.
- [5] James E. Humphreys. *Reflection Groups and Coxeter Groups* -. Cambridge: Cambridge University Press, 1992. ISBN: 978-0-521-43613-7.
- [6] Anne Thomas. *Geometric and Topological Aspects of Coxeter Groups and Buildings* -. European Mathematical Society Publishing House, 2018. ISBN: 978-3-037-19189-7.
- [7] William P. Thurston. "Three dimensional manifolds, Kleinian groups and hyperbolic geometry." In: *Bulletin (New Series) of the American Mathematical Society* 6.3 (1982), pp. 357–381.
- [8] Genevieve Walsh. "Great circle links and virtually fibered knots." In: *Topology* 44 (2004), pp. 947–958.