

# Counting on Thorsten: Representing Ordinals in Homotopy Type Theory

Fredrik Nordvall Forsberg

joint work with **Nicolai Kraus** and **Chuangjie Xu**

Thorstenfest 60

Nottingham, 12 October 2022



Working with Thorsten





# A Categorical Semantics for Inductive-Inductive Definitions

Thorsten Altenkirch<sup>1\*,\*\*</sup>, Peter Morris<sup>1\*\*</sup>, Fredrik Nordvall Forsberg<sup>2\*</sup>, and  
Anton Setzer<sup>2\*</sup>

<sup>1</sup> School of Computer Science, University of Nottingham, UK

<sup>2</sup> Department of Computer Science, Swansea University, UK

**Abstract.** Induction-induction is a principle for defining data types in Martin-Löf Type Theory. An inductive-inductive definition consists of a set  $A$ , together with an  $A$ -indexed family  $B : A \rightarrow \text{Set}$ , where both  $A$  and  $B$  are inductively defined in such a way that the constructors for  $A$  can refer to  $B$  and vice versa. In addition, the constructors for  $B$  can refer to the constructors for  $A$ . We extend the usual initial algebra semantics for ordinary inductive data types to the inductive-inductive setting by considering di-algebras instead of ordinary algebras. This gives a new and compact formalisation of inductive-inductive definitions, which we prove is equivalent to the usual formulation with elimination rules.

## 1 Introduction

Induction is an important principle of definition and reasoning, especially so in constructive mathematics and computer science, where the concept of inductively defined set and data type coincide. There are two well-established approaches to model the semantics of such data types: in Martin-Löf Type Theory [14], each set  $A$  comes equipped with an eliminator which at the same time represents reasoning by induction over  $A$  and the definition of recursive functions out of  $A$ . A more categorical approach [30] models data types as initial  $T$ -algebras for a suitable endofunctor  $T$ .

At first, it would seem that the eliminator approach is stronger, as it allows us to define dependent functions  $(x : A) \rightarrow P(x)$ , in contrast with the non-dependent arrows  $A \rightarrow B$  given by the initiality of the algebra. However, Hermida and Jacobs [22] showed that an eliminator can be defined for every initial  $T$ -algebra, where  $T$  is a polynomial functor. Ghani et al. [8] then extended this to arbitrary endofunctors. This covers many forms of induction and data type definitions such as indexed inductive definitions [5] and induction-recursion [7] (Dybjer and Setzer [8] also give a direct proof for induction-recursion).

There are, however, other meaningful forms of data type which are not covered by these results. One such example are inductive-inductive definitions [16], where a set  $A$  and a family  $B : A \rightarrow \text{Set}$  are simultaneously inductively defined (compare


\* Supported by EPSRC grant EP/G033374/1.

\*\* Supported by EPSRC grant EP/G033399/1.









# Representing ordinals in homotopy type theory

# Ordinals in computer science

**Bold claim:** ordinals are one of the most important structures in logic and computer science.



# Ordinals in computer science

**Bold claim:** ordinals are one of the most important structures in logic and computer science.

**For example:** Ordinal analysis in proof theory, termination of programs, free constructions via transfinite iteration, ...

# Ordinals in computer science

**Bold claim:** ordinals are one of the most important structures in logic and computer science.

**For example:** Ordinal analysis in proof theory, termination of programs, free constructions via transfinite iteration, ...

Unintuitive but amazing that classically all these uses can be reduced to the set-theoretic axioms for ordinals.

# Set-theoretical ordinals

In classical set theory, an ordinal is a set with an order  $\prec$  which is

- ▶ **transitive:**  $(a \prec b) \rightarrow (b \prec c) \rightarrow (a \prec c)$
- ▶ **wellfounded:** every sequence  $a_0 \succ a_1 \succ a_2 \succ a_3 \succ \dots$  terminates
- ▶ **trichotomous:**  $(a \prec b) \vee (a = b) \vee (b \prec a)$

# Set-theoretical ordinals

In set theory, an ordinal is a set with an order  $\prec$  which is

- ▶ **transitive:**  $(a \prec b) \rightarrow (b \prec c) \rightarrow (a \prec c)$
- ▶ **wellfounded:** transfinite induction along  $\prec$  is valid
- ▶ **trichotomous:**  $(a \prec b) \vee (a = b) \vee (b \prec a)$



# Set-theoretical ordinals

In set theory, an ordinal is a set with an order  $\prec$  which is

- ▶ **transitive:**  $(a \prec b) \rightarrow (b \prec c) \rightarrow (a \prec c)$
- ▶ **wellfounded:** transfinite induction along  $\prec$  is valid
- ~~▶ **trichotomous:**  $(a \prec b) \vee (a = b) \vee (b \prec a)$~~
- ▶ **extensional:**  $(\forall a. a \prec b \leftrightarrow a \prec c) \rightarrow b = c$


# Set-theoretical ordinals


In set theory, an ordinal is a set with an order  $\prec$  which is

- ▶ **transitive:**  $(a \prec b) \rightarrow (b \prec c) \rightarrow (a \prec c)$
- ▶ **wellfounded:** transfinite induction along  $\prec$  is valid
- ~~▶ **trichotomous:**  $(a \prec b) \vee (a = b) \vee (b \prec a)$~~
- ▶ **extensional:**  $(\forall a. a \prec b \leftrightarrow a \prec c) \rightarrow b = c$

This definition can be adopted in type theory, but it is not the only candidate!


# Different notions of ordinals in type theory




 N. Kraus, F. N-F., and C. Xu.  
Connecting constructive notions of ordinals in homotopy type theory  
[MFCS 2021](#).

 N. Kraus, F. N-F., and C. Xu.  
Type-Theoretic Approaches to Ordinals  
[arXiv:2208.03844](#)

# Different notions of ordinals in type theory



 N. Kraus, F. N-F., and C. Xu.  
Connecting constructive notions of ordinals in homotopy type theory  
*MFCS 2021*.

 N. Kraus, F. N-F., and C. Xu.  
Type-Theoretic Approaches to Ordinals  
[arXiv:2208.03844](https://arxiv.org/abs/2208.03844)



# Cantor Normal Forms as an inductive-inductive definition

Motivation:  $\alpha = \omega^{\beta_1} + \omega^{\beta_2} + \cdots + \omega^{\beta_n}$  with  $\beta_1 \geq \beta_2 \geq \cdots \geq \beta_n$

# Cantor Normal Forms as an inductive-inductive definition

Motivation:  $\alpha = \omega^{\beta_1} + \omega^{\beta_2} + \dots + \omega^{\beta_n}$  with  $\beta_1 \geq \beta_2 \geq \dots \geq \beta_n$

Let **Cnf** be the type of *hereditary descending lists*:

$0 : \text{Cnf}$

$\omega^- + - : (a : \text{Cnf}) \rightarrow (b : \text{Cnf}) \rightarrow \{p : a \geq \text{fst } b\} \rightarrow \text{Cnf}$

where  $\text{fst } (\omega^a + b) = a$ .

# Cantor Normal Forms as an inductive-inductive definition

Motivation:  $\alpha = \omega^{\beta_1} + \omega^{\beta_2} + \dots + \omega^{\beta_n}$  with  $\beta_1 \geq \beta_2 \geq \dots \geq \beta_n$

Let **Cnf** be the type of *hereditary descending lists*:

$0 : \text{Cnf}$

$\omega^- + - : (a : \text{Cnf}) \rightarrow (b : \text{Cnf}) \rightarrow \{p : a \geq \text{fst } b\} \rightarrow \text{Cnf}$

where  $\text{fst } (\omega^a + b) = a$ .

Order  $\leq$  defined simultaneously (inductively) as lexicographical order.

# Cantor Normal Forms as an inductive-inductive definition

Motivation:  $\alpha = \omega^{\beta_1} + \omega^{\beta_2} + \dots + \omega^{\beta_n}$  with  $\beta_1 \geq \beta_2 \geq \dots \geq \beta_n$

Let **Cnf** be the type of *hereditary descending lists*:

**0** : **Cnf**

$\omega^- + -$  :  $(a : \text{Cnf}) \rightarrow (b : \text{Cnf}) \rightarrow \{p : a \geq \text{fst } b\} \rightarrow \text{Cnf}$

where  $\text{fst } (\omega^a + b) = a$ .

Order  $\leq$  defined simultaneously (inductively) as lexicographical order.

**Equivalent implementations:** (i) Sigma type of normal forms, and (ii) finite hereditary multisets (as a quotient inductive type).



# Brouwer ordinal trees in constructive type theory

Inductive type  $\mathcal{B}$  of Brouwer trees [Brouwer 1926; Martin-Löf 1970]:

data  $\mathcal{B}$  where

zero :  $\mathcal{B}$

succ :  $\mathcal{B} \rightarrow \mathcal{B}$

limit :  $(\mathbb{N} \rightarrow \mathcal{B}) \rightarrow \mathcal{B}$

# Brouwer ordinal trees in constructive type theory

Inductive type  $\mathcal{B}$  of Brouwer trees [Brouwer 1926; Martin-Löf 1970]:

data  $\mathcal{B}$  where

zero :  $\mathcal{B}$

succ :  $\mathcal{B} \rightarrow \mathcal{B}$

limit :  $(\mathbb{N} \rightarrow \mathcal{B}) \rightarrow \mathcal{B}$

Problems:

limit  $(0, 1, 2, 3, \dots) \neq$  limit  $(2, 3, \dots)$

# Brouwer ordinal trees in constructive type theory

Inductive type  $\mathcal{B}$  of Brouwer trees [Brouwer 1926; Martin-Löf 1970]:

data  $\mathcal{B}$  where

zero :  $\mathcal{B}$

succ :  $\mathcal{B} \rightarrow \mathcal{B}$

limit :  $(\mathbb{N} \rightarrow \mathcal{B}) \rightarrow \mathcal{B}$

Problems:

limit (0, 1, 2, 3, ...)  $\neq$  limit (2, 3, ...)

# Brouwer ordinal trees in constructive type theory

Inductive type  $\mathcal{B}$  of Brouwer trees [Brouwer 1926; Martin-Löf 1970]:

data  $\mathcal{B}$  where

zero :  $\mathcal{B}$

succ :  $\mathcal{B} \rightarrow \mathcal{B}$

limit :  $(\mathbb{N} \rightarrow \mathcal{B}) \rightarrow \mathcal{B}$

Problems:

limit (0, 1, 2, 3, ...)  $\neq$  limit (2, 3, ...)

limit (0, 1, 2, 3, ...)  $\neq$  limit (0, 2, 1, 3, ...)

# Brouwer ordinal trees in constructive type theory

Inductive type  $\mathcal{B}$  of Brouwer trees [Brouwer 1926; Martin-Löf 1970]:

data  $\mathcal{B}$  where

zero :  $\mathcal{B}$

succ :  $\mathcal{B} \rightarrow \mathcal{B}$

limit :  $(\mathbb{N} \rightarrow \mathcal{B}) \rightarrow \mathcal{B}$

Problems:

limit (0, 1, 2, 3, ...)  $\neq$  limit (2, 3, ...)

limit (0, 1, 2, 3, ...)  $\neq$  limit (0, 2, 1, 3, ...)

# Brouwer ordinal trees in constructive type theory

Inductive type  $\mathcal{B}$  of Brouwer trees [Brouwer 1926; Martin-Löf 1970]:

data  $\mathcal{B}$  where

zero :  $\mathcal{B}$

succ :  $\mathcal{B} \rightarrow \mathcal{B}$

limit :  $(\mathbb{N} \rightarrow \mathcal{B}) \rightarrow \mathcal{B}$

Problems:

limit (0, 1, 2, 3, ...)  $\neq$  limit (2, 3, ...)

limit (0, 1, 2, 3, ...)  $\neq$  limit (0, 2, 1, 3, ...)

Our solution: induction-induction and path constructors, ensuring:

- ▶ Limits can only be taken of strictly increasing sequences;
- ▶ Bisimilar sequences have equal limits.

# A refined type of Brouwer tree ordinals

```
data Brw where
  zero  : Brw
  succ  : Brw → Brw
  limit : (f : ℕ → Brw) → {f↑ : increasing f} → Brw
  bisim : ∀ f {f↑} g {g↑} →
    f ≈ g →
    limit f {f↑} ≡ limit g {g↑}
  trunc : isSet Brw
```

```
data _≤_ where
  ≤-zero      : ∀ {x} → zero ≤ x
  ≤-trans     : ∀ {x y z} → x ≤ y → y ≤ z → x ≤ z
  ≤-succ-mono : ∀ {x y} → x ≤ y → succ x ≤ succ y
  ≤-cocone    : ∀ {x} f {f↑ k} → (x ≤ f k) → (x ≤ limit f {f↑})
  ≤-limiting  : ∀ f {f↑ x} → ((k : ℕ) → f k ≤ x) → limit f {f↑} ≤ x
  ≤-trunc     : ∀ {x y} → isProp (x ≤ y)
```

note:  $x < y$

means  $\text{succ } x \leq y$

$f \approx g$  means

$\forall k. \exists n. f(k) \leq g(n)$

and vice versa

# A refined type of Brouwer tree ordinals

```
data Brw where
  zero   : Brw
  succ   : Brw → Brw
  limit  : (f : ℕ → Brw) → {f↑ : increasing f} → Brw
  bisim  : ∀ f {f↑} g {g↑} →
    f ≈ g →
    limit f {f↑} ≡ limit g {g↑}
  trunc  : isSet Brw
```

```
data _≤_ where
  ≤-zero      : ∀ {x} → zero ≤ x
  ≤-trans     : ∀ {x y z} → x ≤ y → y ≤ z → x ≤ z
  ≤-succ-mono : ∀ {x y} → x ≤ y → succ x ≤ succ y
  ≤-cocone    : ∀ {x} f {f↑ k} → (x ≤ f k) → (x ≤ limit f {f↑})
  ≤-limiting  : ∀ f {f↑ x} → ((k : ℕ) → f k ≤ x) → limit f {f↑} ≤ x
  ≤-trunc     : ∀ {x y} → isProp (x ≤ y)
```

note:  $x < y$

means  $\text{succ } x \leq y$

$f \approx g$  means

$\forall k. \exists n. f(k) \leq g(n)$

and vice versa

Implemented in **Cubical Agda** (along with everything else).



# Transitive, wellfounded, and extensional orders as a large type

The type `Ord` consists of pairs  $(X : \mathcal{U}, \prec : X \rightarrow X \rightarrow \text{Prop})$  such that  $\prec$  is transitive, extensional, wellfounded [HoTT book].

# Transitive, wellfounded, and extensional orders as a large type

The type `Ord` consists of pairs  $(X : \mathcal{U}, \prec : X \rightarrow X \rightarrow \text{Prop})$  such that  $\prec$  is transitive, extensional, wellfounded [HoTT book].

Order on `Ord` given by monotone functions with *simulation* property.

# Transitive, wellfounded, and extensional orders as a large type

The type **Ord** consists of pairs  $(X : \mathcal{U}, \prec : X \rightarrow X \rightarrow \text{Prop})$  such that  $\prec$  is transitive, extensional, wellfounded [HoTT book].

Order on **Ord** given by monotone functions with *simulation* property.

Theorem (Escardo [2022])

*The type **Ord** has a non-trivial decidable property if and only if weak excluded middle  $\neg P \uplus \neg\neg P$  holds.*

If it quacks like an ordinal

What do these notions have in common?

# If it quacks like an ordinal

What do these notions have in common?

## Theorem

*For each notion of ordinals  $(A, <, \leq)$  considered:*

- ▶  *$A$  is a set and  $<$  and  $\leq$  are Prop-valued.*
- ▶  *$<$  is transitive, extensional and wellfounded.*
- ▶  *$x < y \rightarrow x \leq y$ , and  $x < y \leq z \rightarrow x < z$   
(note:  $x \leq y < z \rightarrow x < z$  equivalent to LEM for *Ord*).*
- ▶ *Can define  $+$ ,  $\times$  and prove they uniquely satisfy relational specification  
(for *Cnf*, *Brw* also exponentiation).*
- ▶ *Can compute limits of *Brw* and *Ord*, but not *Cnf* (implies WLPO).*
- ▶ **Cnf* has decidable equality and  $<$ , whereas *Brw* and *Ord* do not (this is  
equivalent to LPO and LEM respectively).*
- ▶ *There are order-preserving embeddings  $\text{Cnf} \hookrightarrow \text{Brw} \hookrightarrow \text{Ord}$ .*
- ▶ *...*

# If it quacks like an ordinal

What do these notions have in common?

## Theorem

For each notion of ordinals  $(A, <, \leq)$  considered:

►  $A$  is a set and  $<$  and  $\leq$  are Prop-valued

### ► Summary


- • All notions support transfinite induction;
- All notions support ordinal arithmetic (in different ways!);
- • **Cnf**: decidable  $\leftrightarrow$  True; **Brw**: decidable  $\leftrightarrow$  LPO; **Ord**: decidable  $\leftrightarrow$  LEM.

► Can compute limits of **Brw** and **Ord**, but not **Cnf** (implies WLPO).

► **Cnf** has decidable equality and  $<$ , whereas **Brw** and **Ord** do not (this is equivalent to LPO and LEM respectively).

► There are order-preserving embeddings **Cnf**  $\hookrightarrow$  **Brw**  $\hookrightarrow$  **Ord**.

►

A scenic landscape photograph showing a large, turquoise lake nestled between green, forested hills. In the background, there are blue, hazy mountain ranges under a sky filled with white and grey clouds. A winding road is visible on the right side of the image, and a semi-transparent white box with black text is centered over the middle of the scene.

One more notion of ordinal

# Another notion of ordinal

Thorsten suggested the following variation `coOrd`:

## Definition

- ▶ An ordinal is given by a colist of ordinals.
- ▶  $\alpha \leq \beta$  if  $a_i < \beta$  for all  $a_i \in \alpha$ .
- ▶  $\alpha < \beta$  if  $\alpha \leq b_j$  for some  $b_j \in \beta$ .



# Another notion of ordinal

Thorsten suggested the following variation **coOrd**:

## Definition

- ▶ An ordinal is given by a colist of ordinals.
- ▶  $\alpha \leq \beta$  if  $a_i < \beta$  for all  $a_i \in \alpha$ . (Coinductive definition!)
- ▶  $\alpha < \beta$  if  $\alpha \leq b_j$  for some  $b_j \in \beta$ .

# Another notion of ordinal

Thorsten suggested the following variation `coOrd`:

## Definition

- ▶ An ordinal is given by a colist of ordinals.
- ▶  $\alpha \leq \beta$  if  $a_i < \beta$  for all  $a_i \in \alpha$ . (Coinductive definition!)
- ▶  $\alpha < \beta$  if  $\alpha \leq b_j$  for some  $b_j \in \beta$ . (Inductive definition!)

# Another notion of ordinal

Thorsten suggested the following variation **coOrd**:

## Definition

- ▶ An ordinal is given by a colist of ordinals.
  - ▶ We quotient by antisymmetry: if  $\alpha \leq \beta$  and  $\beta \leq \alpha$  then  $\alpha = \beta$ .
- ▶  $\alpha \leq \beta$  if  $a_i < \beta$  for all  $a_i \in \alpha$ . (Coinductive definition!)
- ▶  $\alpha < \beta$  if  $\alpha \leq b_j$  for some  $b_j \in \beta$ . (Inductive definition!)

# Another notion of ordinal

Thorsten suggested the following variation **coOrd**:

## Definition

- ▶ An ordinal is given by a colist of ordinals.
  - ▶ We quotient by antisymmetry: if  $\alpha \leq \beta$  and  $\beta \leq \alpha$  then  $\alpha = \beta$ .
- ▶  $\alpha \leq \beta$  if  $a_i < \beta$  for all  $a_i \in \alpha$ . (Coinductive definition!)
- ▶  $\alpha < \beta$  if  $\alpha \leq b_j$  for some  $b_j \in \beta$ . (Inductive definition!)

**Intuition:**  $[a_1, a_2, \dots]$  is supremum of the successors of all  $a_i$ .

Similar to ideas by Coquand, Lombardi, and Neuwirth [2022].

## Some simple ordinals

0 is represented by the empty colist  $[]$ .

The successor of  $\alpha$  is represented by the colist  $[\alpha]$ .

The limit of  $f : \mathbb{N} \rightarrow \text{coOrd}$  is represented by the infinite colist  $[f(0), f(1), \dots]$ .

# Some simple ordinals

0 is represented by the empty colist  $[]$ .

The successor of  $\alpha$  is represented by the colist  $[\alpha]$ .

The limit of  $f : \mathbb{N} \rightarrow \text{coOrd}$  is represented by the infinite colist  $[f(0), f(1), \dots]$ .

**Proposition:** There is an embedding  $\text{Brw} \hookrightarrow \text{coOrd}$ .

# Some simple ordinals

0 is represented by the empty colist  $[]$ .

The successor of  $\alpha$  is represented by the colist  $[\alpha]$ .

The limit of  $f : \mathbb{N} \rightarrow \text{coOrd}$  is represented by the infinite colist  $[f(0), f(1), \dots]$ .

**Proposition:** There is an embedding  $\text{Brw} \hookrightarrow \text{coOrd}$ .

**Proposition:** There is an embedding  $\alpha \mapsto (\Sigma \beta : \text{coOrd}) (\beta < \alpha) : \text{coOrd} \hookrightarrow \text{Ord}$ .

## Some simple properties

It is decidable if an ordinal is 0, but not if it is a successor or a limit.



## Some simple properties

It is decidable if an ordinal is 0, but not if it is a successor or a limit.

Arithmetic operations can be defined “pointwise”:

$$\alpha + [] = \alpha$$

$$\alpha + (b_1 :: \beta) = (\alpha + b_1) :: (\alpha + \beta)$$

## Some simple properties

It is decidable if an ordinal is 0, but not if it is a successor or a limit.

Arithmetic operations can be defined “pointwise”:

$$\alpha + [] = \alpha$$

$$\alpha + (b_1 :: \beta) = (\alpha + b_1) :: (\alpha + \beta)$$

$<$  and  $\leq$  should have expected properties (e.g.  $<$  wellfounded, extensional, transitive,  $\alpha < \beta \rightarrow \beta \leq \gamma \rightarrow \alpha < \gamma$ , etc).

# Is it more expressive?

One could hope that constructions on `coOrd` are simpler than on `Brw`, since the definition is more compact.

# Is it more expressive?

One could hope that constructions on `coOrd` are simpler than on `Brw`, since the definition is more compact.

## Is it more expressive?

One could hope that constructions on `coOrd` are simpler than on `Brw`, since the definition is more compact.

However in terms of raw power: Classically each colist is either finite or infinite. Hence classically  $\text{coOrd} \hookrightarrow \text{Brw}$ , i.e.  $\text{coOrd} \cong \text{Brw}$  is consistent.

## Is it more expressive?

One could hope that constructions on `coOrd` are simpler than on `Brw`, since the definition is more compact.

However in terms of raw power: Classically each colist is either finite or infinite. Hence classically  $\text{coOrd} \hookrightarrow \text{Brw}$ , i.e.  $\text{coOrd} \cong \text{Brw}$  is consistent.

So our interest in `coOrd` should be constructive. It might be in good case study in combining coinduction and quotient inductive types?

# Conclusions

There are many ways to represent ordinals in type theory; concepts from homotopy type theory help.

Meaningful distinctions deserve to be preserved: use the right representation for the job.

New representation of ordinals based on **colists**.

Happy birthday Thorsten!

