

Non-commutative gauge symmetries

Nicolas Gilliers

NTNU, Trondheim, Norway

07.06.2020

Free probability online seminar

Yang–Mills fields and master fields

Symmetries \rightarrow Interactions

(circle) $U(1)$ \rightsquigarrow Maxwell electromagn.

(quaternion) $SU(2)$ \rightsquigarrow electroweak interaction.

$SU(3)$ \rightsquigarrow Strong interaction
(quark)

(classical) Yang-Mills theory

Space time geometry (Lorentzian geometry)

Holonomy

- $\Sigma =$ plane or a riemannian manifold, P is a G fiber bundle,

Holonomy

- Σ = plane or a riemannian manifold, P is a G fiber bundle,
- $P = \Sigma \times G$, $w \in \Lambda^1(P, \mathfrak{g})$ a *connection*, $m \in \Sigma$, $\gamma \in \text{Loops}(m)$

Holonomy

- Σ = plane or a riemannian manifold, P is a G fiber bundle,
- $P = \Sigma \times G$, $w \in \Lambda^1(P, \mathfrak{g})$ a *connection*, $m \in \Sigma$, $\gamma \in \text{Loops}(m)$
- Parallel transport :

$$//(\gamma, \omega) : P_m \rightarrow P_m,$$

Holonomy

- Σ = plane or a riemannian manifold, P is a G fiber bundle,
- $P = \Sigma \times G$, $w \in \Lambda^1(P, \mathfrak{g})$ a *connection*, $m \in \Sigma$, $\gamma \in \text{Loops}(m)$
- Parallel transport :

$$//(\gamma, \omega) : P_m \rightarrow P_m,$$

$$\omega(\dot{q}_t) = 0, \quad q_t \in P_{\gamma_t}, \quad //(\gamma, \omega)(q_0) = q_1.$$

Holonomy

- Σ = plane or a riemannian manifold, P is a G fiber bundle,
- $P = \Sigma \times G$, $w \in \Lambda^1(P, \mathfrak{g})$ a *connection*, $m \in \Sigma$, $\gamma \in \text{Loops}(m)$
- Parallel transport :

$$//(\gamma, \omega) : P_m \rightarrow P_m,$$

$$\omega(\dot{q}_t) = 0, \quad q_t \in P_{\gamma_t}, \quad //(\gamma, \omega)(q_0) = q_1.$$

$$q_t = (\gamma_t, h_t),$$

$$\text{Hol}^\omega(\gamma) = h_1 \in G,$$

$$\frac{d}{dt} h_t(x) = -\omega_{\gamma_t}(\dot{\gamma}_t) \cdot h_t(x)$$

Holonomy

- Σ = plane or a riemannian manifold, P is a G fiber bundle,
- $P = \Sigma \times G$, $w \in \Lambda^1(P, \mathfrak{g})$ a *connection*, $m \in \Sigma$, $\gamma \in \text{Loops}(m)$
- Parallel transport :


$$//(\gamma, \omega) : P_m \rightarrow P_m,$$

$$\omega(\dot{q}_t) = 0, \quad q_t \in P_{\gamma_t}, \quad //(\gamma, \omega)(q_0) = q_1.$$

$$q_t = (\gamma_t, h_t),$$


$$\text{Hol}^\omega(\gamma) = h_1 \in G,$$

$$\frac{d}{dt} h_t(x) = -\omega_{\gamma_t}(\dot{\gamma}_t) \cdot h_t(x)$$

 Kobayashi–Nomizu

 Bleecker

Holonomy



Loop on the plane \mapsto An element of the group (Holonomy)

What would a random connexion be?

- Affine space of all connections on Σ = configuration space of the theory,

What would a random connexion be?

- Affine space of all connections on Σ = configuration space of the theory,
- Quantify \rightarrow Choose an amplitude for each of these states.

$$\mu^{YM}(d\omega) = \exp(S^{YM}(\omega)) d\omega = \exp\left(\int_{\Sigma} \|\Omega^{\omega}\|_m^2 dm\right) d\omega$$


What would a random connexion be?


- Affine space of all connections on Σ = configuration space of the theory,
- Quantify \rightarrow Choose an amplitude for each of these states.

$$\mu^{YM}(d\omega) = \exp(S^{YM}(\omega)) d\omega = \exp\left(\int_{\Sigma} \|\Omega^{\omega}\|_m^2 dm\right) d\omega$$

Curvature :

$$\Omega^{\omega} = d\omega + \frac{1}{2}[\omega \wedge \omega] \in \Lambda^2(P, \mathfrak{g})$$





- The process $(\text{Hol}(\gamma_t))_{t>0}$ is a **Brownian motion** on the group, **whose law is conjugacy-invariant**.
- If $\overset{\circ}{\ell}_1 \cap \overset{\circ}{\ell}_2 = \emptyset$, $\text{Hol}(\ell_1)$ are $\text{Hol}(\ell_2)$ independent.

Holonomy fields

An holonomy field with structure group G is a collection $\{h_\ell, \ell \in \text{Loops}\}$ of G valued random variables.

- *Multiplicativity* : $h_{\ell_1 \ell_2} = h_{\ell_1} h_{\ell_2}$

Holonomy fields

An holonomy field with structure group G is a collection $\{h_\ell, \ell \in \text{Loops}\}$ of G valued random variables.

- *Multiplicativity* : $h_{\ell_1 \ell_2} = h_{\ell_1} h_{\ell_2}$
- *Gauge invariance* : for all $g \in G$, $gh_\ell g^{-1} \stackrel{d}{=} h_\ell$

Holonomy fields

An holonomy field with structure group G is a collection $\{h_\ell, \ell \in \text{Loops}\}$ of G valued random variables.

- *Multiplicativity* : $h_{\ell_1 \ell_2} = h_{\ell_1} h_{\ell_2}$
- *Gauge invariance* : for all $g \in G$, $gh_\ell g^{-1} \stackrel{d}{=} h_\ell$
- *Invariance for the action of area preserving morphisms* :

$h_{\phi(\ell)} \stackrel{d}{=} h_\ell$ if $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is area preserving .

Holonomy fields

An holonomy field with structure group G is a collection $\{h_\ell, \ell \in \text{Loops}\}$ of G valued random variables.

- *Multiplicativity* : $h_{\ell_1 \ell_2} = h_{\ell_1} h_{\ell_2}$
- *Gauge invariance* : for all $g \in G$, $gh_\ell g^{-1} \stackrel{d}{=} h_\ell$
- *Invariance for the action of area preserving morphisms* :

$$h_{\phi(\ell)} \stackrel{d}{=} h_\ell \text{ if } \phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \text{ is area preserving .}$$

- *Holonomy of simple loops with non-intersecting domains are independent.*

Holonomy fields

⟨⟨ Lévy process indexed by loops ⟩⟩


An holonomy field with structure group G is a collection $\{h_\ell, \ell \in \text{Loops}\}$ of G valued random variables.

- *Multiplicativity* : $h_{\ell_1 \ell_2} = h_{\ell_1} h_{\ell_2}$
- *Gauge invariance* : for all $g \in G$, $gh_\ell g^{-1} \stackrel{d}{=} h_\ell$
- *Invariance for the action of area preserving morphisms* :

$$h_{\phi(\ell)} \stackrel{d}{=} h_\ell \text{ if } \phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \text{ is area preserving .}$$

- *Holonomy of simple loops with non-intersecting domains are independent.*

Which generalization ?



Arrow reversing

$$H_\ell(f) = f(h_\ell), \quad H_{\ell_1, \dots, \ell_n}(f) = f(h_{\ell_1}, \dots, h_{\ell_n})$$

Arrow reversing

$$H_\ell(f) = f(h_\ell), \quad H_{\ell_1, \dots, \ell_n}(f) = f(h_{\ell_1}, \dots, h_{\ell_n})$$

- H_ℓ is an (algebra) **homomorphism** from $\mathcal{F}(G)$ to a space of (almost...) bounded random variables.

Arrow reversing

$$H_\ell(f) = f(h_\ell), \quad H_{\ell_1, \dots, \ell_n}(f) = f(h_{\ell_1}, \dots, h_{\ell_n})$$

- H_ℓ is an (algebra) **homomorphism** from $\mathcal{F}(G)$ to a space of (almost...) bounded random variables.
- **Multiplicativity** : $f(h_{\ell_1 \ell_2}) = f(h_{\ell_1} h_{\ell_2})$, $H_{\ell_1 \ell_2} = H_{\ell_1} \times H_{\ell_2}$

Arrow reversing

$$H_\ell(f) = f(h_\ell), \quad H_{\ell_1, \dots, \ell_n}(f) = f(h_{\ell_1}, \dots, h_{\ell_n})$$

- H_ℓ is an (algebra) **homomorphism** from $\mathcal{F}(G)$ to a space of (almost...) bounded random variables.
- **Multiplicativity** : $f(h_{\ell_1 \ell_2}) = f(h_{\ell_1} h_{\ell_2})$, $H_{\ell_1 \ell_2} = H_{\ell_1} \times H_{\ell_2}$
- **Gauge invariance** :

$$f(Zh_{\ell_1}Z^{-1}, \dots, Zh_{\ell_n}Z^{-1}) \stackrel{(d)}{=} f(h_{\ell_1}, \dots, h_{\ell_n})$$

$$(\tau_Z \otimes \mathbb{E}) \circ (\text{id}_{\mathcal{F}(G)} \sqcup H_{\ell_1, \dots, \ell_n})) \circ \Omega_c^n = \mathbb{E} \circ H_{\ell_1, \dots, \ell_n}.$$

Arrow reversing

$$H_\ell(f) = f(h_\ell), \quad H_{\ell_1, \dots, \ell_n}(f) = f(h_{\ell_1}, \dots, h_{\ell_n})$$

- H_ℓ is an (algebra) **homomorphism** from $\mathcal{F}(G)$ to a space of (almost...) bounded random variables.
- **Multiplicativity** : $f(h_{\ell_1 \ell_2}) = f(h_{\ell_1} h_{\ell_2})$, $H_{\ell_1 \ell_2} = H_{\ell_1} \times H_{\ell_2}$
- **Gauge invariance** :

$$f(Zh_{\ell_1}Z^{-1}, \dots, Zh_{\ell_n}Z^{-1}) \stackrel{(d)}{=} f(h_{\ell_1}, \dots, h_{\ell_n})$$

$$(\tau_Z \otimes \mathbb{E}) \circ (\text{id}_{\mathcal{F}(G)} \sqcup H_{\ell_1, \dots, \ell_n})) \circ \Omega_c^n = \mathbb{E} \circ H_{\ell_1, \dots, \ell_n}.$$

- **Independence** : If ℓ_1 and ℓ_2 have disjoint domains,

H_{ℓ_1} et H_{ℓ_2} are independent.

Arrow reversing

$$H_\ell(f) = f(h_\ell), \quad H_{\ell_1, \dots, \ell_n}(f) = f(h_{\ell_1}, \dots, h_{\ell_n})$$

- H_ℓ is an (algebra) **homomorphism** from $\mathcal{F}(G)$ to a space of (almost...) bounded random variables.
- **Multiplicativity** : $f(h_{\ell_1 \ell_2}) = f(h_{\ell_1} h_{\ell_2}), \quad H_{\ell_1 \ell_2} = H_{\ell_1} \times H_{\ell_2}$
- **Gauge invariance** :

$$f(Zh_{\ell_1}Z^{-1}, \dots, Zh_{\ell_n}Z^{-1}) \stackrel{(d)}{=} f(h_{\ell_1}, \dots, h_{\ell_n})$$

$$(\tau_Z \otimes \mathbb{E}) \circ (\text{id}_{\mathcal{F}(G)} \sqcup H_{\ell_1, \dots, \ell_n})) \circ \Omega_C^n = \mathbb{E} \circ H_{\ell_1, \dots, \ell_n}.$$

- **Independence** : If ℓ_1 and ℓ_2 have disjoint domains,


H_{ℓ_1} et H_{ℓ_2} are independent.

- **Invariance for the action of area preserving homeomorphisms**

$$\mathbb{E} \circ H_{\ell_1, \dots, \ell_n} = \mathbb{E} \circ H_{\phi(\ell_1), \dots, \phi(\ell_n)}.$$

Gauge symmetry and Zhang algebras

Zhang algebra vs Hopf algebra



- Alexeev, Gross & Schomerus (1993)
- Buffenoir, Roche (1995)
- Heusinger (2015)



Algebraic Category

An algebraic category \mathcal{C} has an **initial object** k :


for all $A \in \mathcal{C}$, $\exists!$ $\eta : k \rightarrow A$.

Algebraic Category

An algebraic category \mathcal{C} has an **initial object** k :

for all $A \in \mathcal{C}$, $\exists!$ $\eta : k \rightarrow A$.

and a **coproduct** satisfying the universal property :




Algebraic Category

An algebraic category \mathcal{C} has an **initial object** k :

for all $A \in \mathcal{C}$, $\exists!$ $\eta : k \rightarrow A$.

and a **coproduct** satisfying the universal property :



- Set with disjoint union, $k = \emptyset$,
- ComAlg with (the usual) tensor product, $k = \mathbb{C}$,
- Alg with the free product.
- $\text{Alg}(B)$ with the free product over B the unit is B .

Zhang algebra or H -algebra

A Zhang algebra in \mathcal{C} is a tuple $(H, \Delta, \varepsilon, S)$ with

Zhang algebra or H -algebra

A Zhang algebra in \mathcal{C} is a tuple $(H, \Delta, \varepsilon, S)$ with

1. H is an object in \mathcal{C} ,

Zhang algebra or H -algebra

A Zhang algebra in \mathcal{C} is a tuple $(H, \Delta, \varepsilon, S)$ with

1. H is an object in \mathcal{C} ,
2. (Coproduct) $\Delta : H \rightarrow H \sqcup H$, $(\Delta \sqcup \text{id}_H) \circ \Delta = (\text{id}_H \sqcup \Delta) \circ \Delta$

Zhang algebra or H -algebra

A Zhang algebra in \mathcal{C} is a tuple $(H, \Delta, \varepsilon, S)$ with

1. H is an object in \mathcal{C} ,
2. (Coproduct) $\Delta : H \rightarrow H \sqcup H$, $(\Delta \sqcup \text{id}_H) \circ \Delta = (\text{id}_H \sqcup \Delta) \circ \Delta$
3. (Counit) $\varepsilon : H \rightarrow k$, $(\varepsilon \sqcup \text{id}_H) \circ \Delta = \text{id}_H = (\text{id}_H \sqcup \varepsilon) \circ \Delta$

Zhang algebra or H -algebra

A Zhang algebra in \mathcal{C} is a tuple $(H, \Delta, \varepsilon, S)$ with

1. H is an object in \mathcal{C} ,
2. (Coproduct) $\Delta : H \rightarrow H \sqcup H$, $(\Delta \sqcup \text{id}_H) \circ \Delta = (\text{id}_H \sqcup \Delta) \circ \Delta$
3. (Counit) $\varepsilon : H \rightarrow k$, $(\varepsilon \dot{\sqcup} \text{id}_H) \circ \Delta = \text{id}_H = (\text{id}_H \dot{\sqcup} \varepsilon) \circ \Delta$
4. (Antipode) $S : H \rightarrow H$, $(S \dot{\sqcup} \text{id}_H) \circ \Delta = \eta \circ \varepsilon = (\text{id}_H \dot{\sqcup} S) \circ \Delta$,

Zhang algebra or H -algebra

A Zhang algebra in \mathcal{C} is a tuple $(H, \Delta, \varepsilon, S)$ with

1. H is an object in \mathcal{C} ,
2. (Coproduct) $\Delta : H \rightarrow H \sqcup H$, $(\Delta \sqcup \text{id}_H) \circ \Delta = (\text{id}_H \sqcup \Delta) \circ \Delta$
3. (Counit) $\varepsilon : H \rightarrow k$, $(\varepsilon \dot{\sqcup} \text{id}_H) \circ \Delta = \text{id}_H = (\text{id}_H \dot{\sqcup} \varepsilon) \circ \Delta$
4. (Antipode) $S : H \rightarrow H$, $(S \dot{\sqcup} \text{id}_H) \circ \Delta = \eta \circ \varepsilon = (\text{id}_H \dot{\sqcup} S) \circ \Delta$,

» A commutative Hopf algebra is a Zhang algebra in the category ComAlg .

- Algebra of functions on a group in ComAlg^\star

$$f_1 f_2(g) = f_1(g) f_2(g), \quad \Delta(f)(g_1, g_2) = f(g_1 g_2), \quad S(f)(g) = f(g^{-1}).$$

- Voiculescu's dual groups, in Alg^\star ,

$$\mathcal{O}\langle n \rangle = \langle u_{ij}, u_{ij}^\star \mid \sum_k u_{ik} u_{jk}^\star = 1 \rangle \quad \Delta(u_{ij}) = u_{ik} u_{ki}, \quad S(u_{ij}) = u_{ji}^\star.$$

- Set $B = \langle p_1, \dots, p_n \mid p_i p_j = p_j p_i = 0, p_i^2 = p_i, i \neq j \rangle$ in $\text{Alg}(B)$.

$$\mathcal{O}_B\langle n \rangle = \langle u, u^\star \mid uu^\star = 1 \rangle_B \quad \Delta(u) = u_{|1} u_{|2}, \quad S(u) = u^\star$$

Graphical calculus à la Hopf

$$\Delta = \begin{array}{c} \diagup \quad \diagdown \\ | \end{array} \quad \mu = \begin{array}{c} | \\ \diagdown \quad \diagup \end{array} \quad S = \begin{array}{c} | \\ \hline | \end{array} \quad \varepsilon = \begin{array}{c} \bullet \\ | \end{array} \quad \eta = \begin{array}{c} | \\ \bullet \end{array}$$

Graphical calculus à la Hopf

$$\Delta = \begin{array}{c} \diagup \quad \diagdown \\ | \end{array} \quad \mu = \begin{array}{c} | \\ \diagdown \quad \diagup \end{array} \quad S = \begin{array}{c} | \\ \hline \end{array} \quad \varepsilon = \begin{array}{c} \bullet \\ | \end{array} \quad \eta = \begin{array}{c} | \\ \bullet \end{array}$$

$\alpha, \beta : H \rightarrow C$, morphisms in \mathcal{C} .

$\alpha \times \beta = (\alpha \sqcup \beta) \circ \Delta$ is a morphism in the category

$\alpha, \beta : H \rightarrow C$, morphisms in \mathcal{C} .

» $\alpha \times \beta = (\alpha \sqcup \beta) \circ \Delta$ is a morphism in the category

$\alpha, \beta : H \rightarrow C$, morphisms in \mathcal{C} .

» $\alpha \times \beta = (\alpha \sqcup \beta) \circ \Delta$ is a morphism in the category

$(\text{Hom}_{\mathcal{C}}(H, A), \times)$ is a group.

$\alpha, \beta : H \rightarrow C$, morphisms in \mathcal{C} .


» $\alpha \times \beta = (\alpha \sqcup \beta) \circ \Delta$ is a morphism in the category

$(\text{Hom}_{\mathcal{C}}(H, A), \times)$ is a group.

J. Zhang

For any functor $F : \text{Alg}(\mathcal{C}) \rightarrow \mathcal{G}rp$, there exists an Zhang algebra

$$F(\cdot) = \text{Hom}_{\mathcal{C}}(H, \cdot)$$

 J. Zhang, H-algebra, 1991.

Comodules and Conjugation coaction

Let (\mathcal{C}, \sqcup, k) be an algebraic category A comodule (M, ρ) over a Zhang algebra $(H, \Delta, \varepsilon, S)$ is :

- An object in \mathcal{C}
- A morphism $\rho \in \text{Hom}_{\mathcal{C}}(M, M \sqcup H)$,

$$(\rho \sqcup \text{id}_H) \circ \rho = (\text{id}_M \dot{\sqcup} \Delta) \circ \rho, \quad (\text{id}_M \dot{\sqcup} \varepsilon) \circ \rho = \text{id}_M.$$

Comodules and Conjugation coaction

Let (\mathcal{C}, \sqcup, k) be an algebraic category A comodule (M, ρ) over a Zhang algebra $(H, \Delta, \varepsilon, S)$ is :

- An object in \mathcal{C}
- A morphism $\rho \in \text{Hom}_{\mathcal{C}}(M, M \sqcup H)$,

$$(\rho \sqcup \text{id}_H) \circ \rho = (\text{id}_M \dot{\sqcup} \Delta) \circ \rho, \quad (\text{id}_M \dot{\sqcup} \varepsilon) \circ \rho = \text{id}_M.$$

Proposition

The category $\text{ICoMod}\mathcal{C}(H)$ is an algebraic category.

Proposition (G. 2019)

A Zhang algebra H is a left comodule over H :



$$\Omega_c = (\iota_1 \dot{\sqcup} \iota_2 \dot{\sqcup} \iota_1) \circ (\text{id}_{H \sqcup H} \sqcup S) \circ (\Delta \sqcup \text{id}_H) \circ \Delta.$$

Proposition (G. 2019)

A Zhang algebra H is a left comodule over H :



$$\Omega_c = (\iota_1 \dot{\sqcup} \iota_2 \dot{\sqcup} \iota_1) \circ (\mathrm{id}_{H \sqcup H} \sqcup S) \circ (\Delta \sqcup \mathrm{id}_H) \circ \Delta.$$

Conjugation coaction on the algebra of functions on a group

$$\Omega^c(f)(h, g) = f(hgh^{-1}), \quad f \in \mathcal{F}(G), \quad g, h \in G.$$

...on the dual Voiculescu group $\mathcal{O}\langle n \rangle$:

$$\Omega^c(o_{ij}) = \sum_{k,q=1}^n o_{ik}|_1 o_{kq}|_2 o_{kq}^*|_1$$

Categorical independence and monoidal structures

Classical independence and commutative diagrams

X, Y two random variables. They are independent if and only if :

$$\begin{array}{ccc} & (L^\infty(\Omega, \mathcal{F}), \mathbb{E}) & \\ j_X \nearrow & & \nwarrow j_Y \\ (L^\infty(S, \mathcal{S}), \tau_X) & & (L^\infty(T, \mathcal{T}), \tau_Y) \end{array}$$

avec :

$$\begin{aligned} j_X : \quad L^\infty(S, \mathcal{S}) &\rightarrow L(\Omega, \mathcal{F}, \mathbb{P}) \\ f &\mapsto f(X) \end{aligned},$$
$$\tau_X = \mathbb{E} \circ j_X.$$

Classical independence and commutative diagrams

X, Y two random variables. They are independent if and only if :

$$\begin{array}{ccccc}
 & & (L^\infty(\Omega, \mathcal{F}), \mathbb{E}) & & \\
 & \nearrow j_X & & \nwarrow j_Y & \\
 (L^\infty(S, \mathcal{S}), \tau_X) & \xrightarrow{\iota} & (L^\infty(S, \mathcal{S}), \tau_X) \otimes (L^\infty(T, \mathcal{T}), \tau_Y) & \xleftarrow{\iota} & (L^\infty(T, \mathcal{T}), \tau_Y)
 \end{array}$$

avec :

$$\begin{aligned}
 j_X : \quad L^\infty(S, \mathcal{S}) &\rightarrow L(\Omega, \mathcal{F}, \mathbb{P}) \\
 f &\mapsto f(X) \\
 \tau_X &= \mathbb{E} \circ j_X.
 \end{aligned}$$

Classical independence and random variables

X, Y two random variables. They are independent if and only if :

$$\begin{array}{ccccc}
 & & (L^\infty(\Omega, \mathcal{F}), \mathbb{E}) & & \\
 & \nearrow j_X & \uparrow j & \nwarrow j_Y & \\
 (L^\infty(S, \mathcal{S}), \tau_X) & \xrightarrow{\iota} & (L^\infty(S, \mathcal{S}), \tau_X) \otimes (L^\infty(T, \mathcal{T}), \tau_Y) & \xleftarrow{\iota} & (L^\infty(T, \mathcal{T}), \tau_Y)
 \end{array}$$

with :

$$\begin{aligned}
 j_X : \quad L^\infty(S, \mathcal{S}) &\rightarrow L(\Omega, \mathcal{F}, \mathbb{P}) \\
 f &\mapsto f(X) \quad , \\
 \tau_X &= \mathbb{E} \circ j_X.
 \end{aligned}$$

Monoidal structures

- tensor product of vector spaces, $V, W \in \mathbf{Vect}$,

$$V \otimes W = \{vw, v \in V, w \in W\}, \quad v(\lambda w) = (\lambda v)w = \lambda(vw).$$

- tensor product of unital algebras, $A, B \in \mathbf{ComAlg}$

$$a_1 \otimes b_1 \cdot a_2 \otimes b_2 = a_1 a_2 \otimes b_1 b_2$$

- tensor product of non-commutative unital algebras, $A, B \in \mathbf{Alg}$

Monoidal structures

- tensor product of vector spaces, $V, W \in \mathbf{Vect}$,

$$V \otimes W = \{vw, v \in V, w \in W\}, \quad v(\lambda w) = (\lambda v)w = \lambda(vw).$$

- tensor product of unital algebras, $A, B \in \mathbf{ComAlg}$

$$a_1 \otimes b_1 \cdot a_2 \otimes b_2 = a_1 a_2 \otimes b_1 b_2$$

- tensor product of non-commutative unital algebras, $A, B \in \mathbf{Alg}$

\otimes is...


- ...associative :

$$\alpha_{A,B,C} : A \otimes (B \otimes C) \xrightarrow{\cong} (A \otimes B) \otimes C,$$

- ...has a unit object $E \in \mathbf{Obj}(\mathcal{C})$:

$$\ell_A : E \otimes A \xrightarrow{\cong} A, \quad r_A : A \otimes E \xrightarrow{\cong} A$$

Categorical independence



Categorical independence

$$\begin{array}{ccccc} & & A & & \\ & \nearrow f_1 & & \nwarrow f_2 & \\ C_1 & \xrightarrow{\iota_{C_1}} & C_1 \otimes C_2 & \xleftarrow{\iota_{C_2}} & C_2 \end{array}$$

Categorical independence

$$\begin{array}{ccccc} & & A & & \\ & \nearrow f_1 & \uparrow f & \nwarrow f_2 & \\ C_1 & \xrightarrow{\iota_{C_1}} & C_1 \otimes C_2 & \xleftarrow{\iota_{C_2}} & C_2 \end{array}$$

Categorical independence

$$\begin{array}{ccccc} & & A & & \\ & \nearrow f_1 & \uparrow f & \nwarrow f_2 & \\ C_1 & \xrightarrow{\iota_{C_1}} & C_1 \otimes C_2 & \xleftarrow{\iota_{C_2}} & C_2 \end{array}$$


- f is not unique,

Categorical independence

$$\begin{array}{ccccc} & & A & & \\ & \nearrow f_1 & \uparrow f & \nwarrow f_2 & \\ C_1 & \xrightarrow{\iota_{C_1}} & C_1 \otimes C_2 & \xleftarrow{\iota_{C_2}} & C_2 \end{array}$$

- f is not unique,
- (f_1, f_2) independent $\not\Rightarrow (f_2, f_1)$ independent,

Categorical independence



- f is not unique,
- (f_1, f_2) independent $\not\Rightarrow (f_2, f_1)$ independent,
- This does not define " (f_1, \dots, f_n) independent",

Independence in $\text{Prob}(\mathbb{C})$

Pick $(\mathcal{A}_1, \phi_{\mathcal{A}_1})$ and $(\mathcal{A}_2, \phi_{\mathcal{A}_2})$ two probability spaces.

Universal classical tensor product $\hat{\otimes}$

$$\mathcal{A}_1 \hat{\otimes} \mathcal{A}_2 = \mathcal{A}_1 \sqcup \mathcal{A}_2,$$

$$(\phi_{\mathcal{A}_1} \hat{\otimes} \phi_{\mathcal{A}_2})(a_1^{i_1} a_2^{i_2} \cdot \dots \cdot a_p^{i_p}) = \phi_{\mathcal{A}_1}(a_{j_1^1}^1 \cdots a_{j_{q_1}^1}^1) \phi_{\mathcal{A}_2}(a_{j_1^2}^2 \cdots a_{j_{q_2}^2}^2)$$

Independence in Prob(\mathbb{C})

Pick $(\mathcal{A}_1, \phi_{\mathcal{A}_1})$ and $(\mathcal{A}_2, \phi_{\mathcal{A}_2})$ two probability spaces.

Universal classical tensor product $\hat{\otimes}$

$$\mathcal{A}_1 \hat{\otimes} \mathcal{A}_2 = \mathcal{A}_1 \sqcup \mathcal{A}_2,$$

$$(\phi_{\mathcal{A}_1} \hat{\otimes} \phi_{\mathcal{A}_2})(a_1^{i_1} a_2^{i_2} \cdots a_p^{i_p}) = \phi_{\mathcal{A}_1}(a_{j_1}^1 \cdots a_{j_{q_1}}^1) \phi_{\mathcal{A}_2}(a_{j_1}^2 \cdots a_{j_{q_2}}^2)$$

Free tensor product \star

$$\mathcal{A}_1 \star \mathcal{A}_2 = \mathcal{A}_1 \sqcup \mathcal{A}_2,$$

$$(\phi_{\mathcal{A}_1} \star \phi_{\mathcal{A}_2})(s_1 s_2 \cdots s_p) = 0, \quad s_j \in \mathcal{A}_{i_j}, \quad i_j \neq i_{j+1}, \quad \phi_{\mathcal{A}_{i_j}}(s_j) = 0.$$

Independence in $\text{Prob}(\mathbb{C})$

Pick $(\mathcal{A}_1, \phi_{\mathcal{A}_1})$ and $(\mathcal{A}_2, \phi_{\mathcal{A}_2})$ two probability spaces.

Universal classical tensor product $\hat{\otimes}$

$$\mathcal{A}_1 \hat{\otimes} \mathcal{A}_2 = \mathcal{A}_1 \sqcup \mathcal{A}_2,$$

$$(\phi_{\mathcal{A}_1} \hat{\otimes} \phi_{\mathcal{A}_2})(a_1^{i_1} a_2^{i_2} \cdots a_p^{i_p}) = \phi_{\mathcal{A}_1}(a_{j_1}^1 \cdots a_{j_{q_1}}^1) \phi_{\mathcal{A}_2}(a_{j_1}^2 \cdots a_{j_{q_2}}^2)$$

Free tensor product \star

$$\mathcal{A}_1 \star \mathcal{A}_2 = \mathcal{A}_1 \sqcup \mathcal{A}_2,$$

$$(\phi_{\mathcal{A}_1} \star \phi_{\mathcal{A}_2})(s_1 s_2 \cdots s_p) = 0, \quad s_j \in \mathcal{A}_{i_j}, \quad i_j \neq i_{j+1}, \quad \phi_{\mathcal{A}_{i_j}}(s_j) = 0.$$

Boolean tensor product \diamond

$$\mathcal{A}_1 \diamond \mathcal{A}_2 = \mathcal{A}_1 \sqcup \mathcal{A}_2,$$

$$(\phi_{\mathcal{A}_1} \diamond \phi_{\mathcal{A}_2})(a_1^{i_1} a_2^{i_2} \cdots a_p^{i_p}) = \phi_{\mathcal{A}_{i_1}}(a_1^{i_1}) \cdots \phi_{\mathcal{A}_{i_p}}(a_p^{i_p}).$$

Non-commutative holonomy fields

Quantum Levy processes

$$\mathcal{C} = \text{Alg}^*, (\text{Prob}, \otimes) \ni (\mathcal{A}, \tau)$$

H is a Zhang algebra in Alg^* .

Quantum Levy processes

$$\mathcal{C} = \text{Alg}^*, (\text{Prob}, \otimes) \ni (\mathcal{A}, \tau)$$

H is a Zhang algebra in Alg^* .

A *Lévy process* is a family of morphisms of Alg^* :

$$j_s : H \rightarrow \mathcal{A}$$

Quantum Lévy processes

$$\mathcal{C} = \text{Alg}^*, (\text{Prob}, \otimes) \ni (\mathcal{A}, \tau)$$

H is a Zhang algebra in Alg^* .

A *Lévy process* is a family of morphisms of Alg^* :

$$j_s : H \rightarrow \mathcal{A}$$

The increments $j_{s,t} = j_s^{-1} \times j_t = ((j_s \circ S) \sqcup j_t) \circ \Delta$ satisfy

- for any tuple $u < s < t$, $j_{u,s} \times j_{s,t} = j_{u,t}$, $j_{s,s} = \varepsilon$,

Quantum Lévy processes

$$\mathcal{C} = \text{Alg}^*, (\text{Prob}, \otimes) \ni (\mathcal{A}, \tau)$$

H is a Zhang algebra in Alg^* .

A *Lévy process* is a family of morphisms of Alg^* :

$$j_s : H \rightarrow \mathcal{A}$$

The increments $j_{s,t} = j_s^{-1} \times j_t = ((j_s \circ S) \dot{\sqcup} j_t) \circ \Delta$ satisfy

- for any tuple $u < s < t$, $j_{u,s} \times j_{s,t} = j_{u,t}$, $j_{s,s} = \varepsilon$,
- For times $(s_1 < t_1 \leq s_2 < t_2 \leq \dots \leq s_p < t_p)$,

$$\tau \circ (j_{s_1,1} \dot{\sqcup} \dots \dot{\sqcup} j_{s_p,t_p}) = \tau \circ j_{s_1,t_1} \otimes \dots \otimes \tau \circ j_{s_p,t_p}$$

Quantum Lévy processes

$$\mathcal{C} = \text{Alg}^*, (\text{Prob}, \otimes) \ni (\mathcal{A}, \tau)$$

H is a Zhang algebra in Alg^* .

A *Lévy process* is a family of morphisms of Alg^* :

$$j_s : H \rightarrow \mathcal{A}$$

The increments $j_{s,t} = j_s^{-1} \times j_t = ((j_s \circ S) \dot{\sqcup} j_t) \circ \Delta$ satisfy

- for any tuple $u < s < t$, $j_{u,s} \times j_{s,t} = j_{u,t}$, $j_{s,s} = \varepsilon$,
- For times $(s_1 < t_1 \leq s_2 < t_2 \leq \dots \leq s_p < t_p)$,

$$\tau \circ (j_{s_1,1} \dot{\sqcup} \dots \dot{\sqcup} j_{s_p,t_p}) = \tau \circ j_{s_1,t_1} \otimes \dots \otimes \tau \circ j_{s_p,t_p}$$

- $\tau \circ j_{s,t} = \tau \circ j_{t-s}$

Holonomy fields

$$\circ H_\ell, \ell \in \text{Loops}_0(\mathbb{R}^2), H_\ell \in \text{Hom}_{\text{Alg}^*}(H, (\mathcal{A}, \tau))$$

Holonomy fields

- $H_\ell, \ell \in \text{Loops}_0(\mathbb{R}^2)$, $H_\ell \in \text{Hom}_{\text{Alg}^*}(H, (\mathcal{A}, \tau))$
- **Multiplicativity** : $H_{\ell_1 \ell_2} = H_{\ell_1} \times H_{\ell_2}$

Holonomy fields

- $H_\ell, \ell \in \text{Loops}_0(\mathbb{R}^2)$, $H_\ell \in \text{Hom}_{\text{Alg}^*}(H, (\mathcal{A}, \tau))$
- **Multiplicativity** : $H_{\ell_1 \ell_2} = H_{\ell_1} \times H_{\ell_2}$
- **Gauge invariance** :

$$(\phi_H \otimes \phi_{\mathcal{A}}) \circ (\text{id}_B \sqcup H_{\ell_1, \dots, \ell_n})) \circ \Omega_c^n = \phi_{\mathcal{A}} \circ H_{\ell_1, \dots, \ell_n}.$$

Holonomy fields

- $H_\ell, \ell \in \text{Loops}_0(\mathbb{R}^2)$, $H_\ell \in \text{Hom}_{\text{Alg}^*}(H, (\mathcal{A}, \tau))$
- **Multiplicativity** : $H_{\ell_1 \ell_2} = H_{\ell_1} \times H_{\ell_2}$
- **Gauge invariance** :

$$(\phi_H \otimes \phi_{\mathcal{A}}) \circ (\text{id}_B \sqcup H_{\ell_1, \dots, \ell_n})) \circ \Omega_c^n = \phi_{\mathcal{A}} \circ H_{\ell_1, \dots, \ell_n}.$$

- Si (ℓ_1, \dots, ℓ_n) and $(\ell'_1, \dots, \ell'_m)$ are two family of loops contained in two simple loops with non-intersecting domains,

$H_{\ell_1, \dots, \ell_n}$ et $H_{\ell'_1, \dots, \ell'_m}$ are \otimes -independent.

Holonomy fields

- $H_\ell, \ell \in \text{Loops}_0(\mathbb{R}^2)$, $H_\ell \in \text{Hom}_{\text{Alg}^*}(H, (\mathcal{A}, \tau))$
- **Multiplicativity** : $H_{\ell_1 \ell_2} = H_{\ell_1} \times H_{\ell_2}$
- **Gauge invariance** :

$$(\phi_H \otimes \phi_{\mathcal{A}}) \circ (\text{id}_B \sqcup H_{\ell_1, \dots, \ell_n})) \circ \Omega_c^n = \phi_{\mathcal{A}} \circ H_{\ell_1, \dots, \ell_n}.$$

- Si (ℓ_1, \dots, ℓ_n) and $(\ell'_1, \dots, \ell'_m)$ are two family of loops contained in two simple loops with non-intersecting domains,

$H_{\ell_1, \dots, \ell_n}$ et $H_{\ell'_1, \dots, \ell'_m}$ are \otimes -independent.

- **Invariance by area preserving morphisms** :


$$\phi_{\mathcal{A}} \circ H_{\ell_1} \dot{\sqcup} \dots \dot{\sqcup} H_{\ell_n} = \phi_{\mathcal{A}} \circ H_{\phi(\ell_1)} \dot{\sqcup} \dots \dot{\sqcup} H_{\phi(\ell_n)}.$$

Theorem

To any quantum Lévy process satisfying two properties one can associate an (algebraic) non-commutative holonomy field.


How to construct such a field ?

Affine loops, direct limit



How to construct such a field ?


Affine loops, direct limit



$$\mathrm{RL}_0(\mathbb{R}^2) = \varinjlim_{\mathbb{G}} \mathrm{RL}_0(\mathbb{G})$$

How to construct such a field ?

Affine loops, direct limit




$$\mathrm{RL}_0(\mathbb{R}^2) = \varinjlim_{\mathbb{G}} \mathrm{RL}_0(\mathbb{G})$$

$$\mathrm{Hom}(\mathrm{RL}_0(\mathbb{R}^2), G) = \varprojlim_{\mathbb{G}} \mathrm{Hom}(\mathrm{RL}_0(\mathbb{G}), G).$$

How to construct such a field?

To a sequence of loops there is an associated graph :



How to construct such a field ?


We construct (\mathcal{A}, τ) as a direct limit,

$$(\mathcal{A}, \tau) = \varinjlim_{\mathbb{G}} (\mathcal{A}(\mathbb{G}), \tau_{\mathbb{G}})$$

How to construct such a field ?

We construct (\mathcal{A}, τ) as a direct limit,

$$(\mathcal{A}, \tau) = \varinjlim_{\mathbb{G}} (\mathcal{A}(\mathbb{G}), \tau_{\mathbb{G}})$$



How to construct such a field?


Basis of counter clockwise lassos

We can find a **basis of lassos** $(c_1, \dots, c_{|F|})$ for the previous graph...

How to construct such a field?

Basis of counter clockwise lassos


We can find a **basis of lassos** $(c_1, \dots, c_{|F|})$ for the previous graph...



How to construct such a field?

Basis of counter clockwise lassos

We can find a **basis of lassos** $(c_1, \dots, c_{|\mathbb{F}|})$ for the previous graph...




$$(\mathcal{A}(\mathbb{G}), \tau_{\mathbb{G}}) = (H^{|\mathbb{F}|}, \tau_{|c_1|} \otimes \dots \otimes \tau_{|c_{|\mathbb{F}|}|})$$

How to construct such a field ?

Basis of counter clockwise lassos

We can find a **basis of lassos** $(c_1, \dots, c_{|\mathbb{F}|})$ for the previous graph...




$$(\mathcal{A}(\mathbb{G}), \tau_{\mathbb{G}}) = (H^{|\mathbb{F}|}, \tau_{|c_1|} \otimes \dots \otimes \tau_{|c_{|\mathbb{F}|}|})$$

$$\text{RL}(\mathbb{G}) = \langle c_1, \dots, c_{|\mathbb{F}|} \rangle$$

How to construct such a field ?

Basis of counter clockwise lassos

We can find a **basis of lassos** $(c_1, \dots, c_{|\mathbb{F}|})$ for the previous graph...



$$(\mathcal{A}(\mathbb{G}), \tau_{\mathbb{G}}) = (H^{|\mathbb{F}|}, \tau_{|c_1|} \otimes \dots \otimes \tau_{|c_{|\mathbb{F}|}|})$$


$$\text{RL}(\mathbb{G}) = \langle c_1, \dots, c_{|\mathbb{F}|} \rangle$$

$$\begin{array}{ccc} \text{Hol} : & \text{RL}(\mathbb{G}) & \rightarrow H \rightarrow H^{\mathbb{F}} \\ & c_j & \mapsto \iota_j \end{array}$$

How to construct such a field ?

Braids and Artin's theorem


... but wait, there are plenty of them !



How to construct such a field ?

Braids and Artin's theorem

... but wait, there are plenty of them !




Proposition

Two basis of counter clockwise lassos are related by mean of the action of a *braid*.

Braids & Lassos

Braids & Lassos



What is a non-commutative Hopf algebras?

Commutative Hopf algebras represent certain functors

$$F : \text{ComAlg} \rightarrow \text{Grp}$$

What is a non-commutative Hopf algebras?

Commutative Hopf algebras represent certain functors

$$F : \text{ComAlg} \rightarrow \text{Grp}$$

...Zhang algebras represent the same kind of functors but

$$F : \text{Alg} \rightarrow \text{Grp}$$

What is a non-commutative Hopf algebras?

Commutative Hopf algebras represent certain functors

$$F : \text{ComAlg} \rightarrow \text{Grp}$$

...Zhang algebras represent the same kind of functors but

$$F : \text{Alg} \rightarrow \text{Grp}$$

But there is another (**higher categorical** flavoured way) to see commutative Hopf algebras!

What is a non-commutative Hopf algebras ?

Commutative Hopf algebras represent certain functors

$$F : \text{ComAlg} \rightarrow \text{Grp}$$

...Zhang algebras represent the same kind of functors but

$$F : \text{Alg} \rightarrow \text{Grp}$$

But there is another (**higher categorical** flavoured way) to see commutative Hopf algebras !

» Zhang algebra are also useful for the shuffle approach to (operator-valued) non-commutative proba !

Two monoidal categories (after Aguiar and Mahajan)

$$(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$$

Two monoidal categories (after Aguiar and Mahajan)

$$(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$$

Proposition

Complex unital algebras in $(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$ with **equal units** are **commutative** algebras.

Two monoidal categories (after Aguiar and Mahajan)

$$(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$$

Proposition

Complex unital algebras in $(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$ with **equal units** are **commutative** algebras.

$$m^1 : A \otimes A \rightarrow A, \quad m^2 : (A \otimes A, m^1_{A \otimes A}) \rightarrow (A, m^1), \quad \eta^1 = \eta^2.$$

Two monoidal categories (after Aguiar and Mahajan)

$$(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$$

Proposition

Complex unital algebras in $(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$ with **equal units** are **commutative** algebras.

$$m^1 : A \otimes A \rightarrow A, \quad m^2 : (A \otimes A, m^1_{A \otimes A}) \rightarrow (A, m^1), \quad \eta^1 = \eta^2.$$

$$(\mathrm{Vect}_{\mathbb{C}}, \otimes, \mathbb{C}, \otimes, \mathbb{C})$$

Two monoidal categories (after Aguiar and Mahajan)

$$(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$$

Proposition

Complex unital algebras in $(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$ with **equal units** are **commutative** algebras.

$$m^1 : A \otimes A \rightarrow A, \quad m^2 : (A \otimes A, m^1_{A \otimes A}) \rightarrow (A, m^1), \quad \eta^1 = \eta^2.$$

$$(\mathrm{Vect}_{\mathbb{C}}, \otimes, \mathbb{C}, \otimes, \mathbb{C})$$

... What if ...

$$(\mathrm{Vect}_{\mathbb{C}}, \otimes, \mathbb{C}, \boxtimes, \mathbb{C}),$$

Two monoidal categories (after Aguiar and Mahajan)

$$(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$$

Proposition

Complex unital algebras in $(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$ with **equal units** are **commutative** algebras.

$$m^1 : A \otimes A \rightarrow A, \quad m^2 : (A \otimes A, m^1_{A \otimes A}) \rightarrow (A, m^1), \quad \eta^1 = \eta^2.$$

$$(\mathrm{Vect}_{\mathbb{C}}, \otimes, \mathbb{C}, \otimes, \mathbb{C})$$

... What if ...

$$(\mathrm{Vect}_{\mathbb{C}}, \otimes, \mathbb{C}_{\otimes}, \boxtimes, \mathbb{C}_{\boxtimes}),$$

Two monoidal categories (after Aguiar and Mahajan)

$$(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$$

Proposition

Complex unital algebras in $(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$ with **equal units** are **commutative** algebras.

$$m^1 : A \otimes A \rightarrow A, \quad m^2 : (A \otimes A, m^1_{A \otimes A}) \rightarrow (A, m^1), \quad \eta^1 = \eta^2.$$

$$(\mathrm{Vect}_{\mathbb{C}}, \otimes, \mathbb{C}, \otimes, \mathbb{C})$$

... What if ...

$$(\mathrm{Vect}_{\mathbb{C}}, \otimes, \mathbb{C}_{\otimes}, \boxtimes, \mathbb{C}_{\boxtimes}), \quad i : \mathbb{C}_{\otimes} \rightarrow \mathbb{C}_{\boxtimes}$$

Two monoidal categories (after Aguiar and Mahajan)

$$(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$$

Proposition

Complex unital algebras in $(\mathrm{Alg}_{\mathbb{C}}, \otimes, \mathbb{C})$ with **equal units** are **commutative** algebras.

$$m^1 : A \otimes A \rightarrow A, \quad m^2 : (A \otimes A, m^1_{A \otimes A}) \rightarrow (A, m^1), \quad \eta^1 = \eta^2.$$

$$(\mathrm{Vect}_{\mathbb{C}}, \otimes, \mathbb{C}, \otimes, \mathbb{C})$$

... What if ...

$$(\mathrm{Vect}_{\mathbb{C}}, \otimes, \mathbb{C}_{\otimes}, \boxtimes, \mathbb{C}_{\boxtimes}), \quad i : \mathbb{C}_{\otimes} \rightarrow \mathbb{C}_{\boxtimes}$$

$$R : (A \boxtimes B) \otimes (C \boxtimes D) \rightarrow (A \otimes C) \boxtimes (B \otimes D)$$

PROS & Hopf PROS

$$m_1 : A \otimes A \rightarrow A, \quad m^2 : A \boxtimes A \rightarrow A, \quad \eta^{\otimes} \circ i = \eta^{\boxtimes}.$$

$$m^2 : (A \boxtimes A, (m^1 \boxtimes m^1) \circ R) \rightarrow (A, m^1)$$

This is what we call **a PROS**

PROS & Hopf PROS

$$m_1 : A \otimes A \rightarrow A, \quad m^2 : A \boxtimes A \rightarrow A, \quad \eta^{\otimes} \circ i = \eta^{\boxtimes}.$$

$$m^2 : (A \boxtimes A, (m^1 \boxtimes m^1) \circ R) \rightarrow (A, m^1)$$

This is what we call **a PROS**

... We can look for PROS that are Hopf algebras...

PROS & Hopf PROS

$$m_1 : A \otimes A \rightarrow A, \quad m^2 : A \boxtimes A \rightarrow A, \quad \eta^{\otimes} \circ i = \eta^{\boxtimes}.$$

$$m^2 : (A \boxtimes A, (m^1 \boxtimes m^1) \circ R) \rightarrow (A, m^1)$$

This is what we call a **PROS**

... We can look for PROS that are Hopf algebras...

$$\Delta : (A, m^1) \rightarrow (A \boxtimes A, (m^1 \boxtimes m^1) \circ R), \quad \varepsilon : A \rightarrow \mathbb{C}_{\boxtimes}.$$

» Applications to pre-Lie theoretical perspective on amalgamated non-commutative probability.