

## Uneven balance power between hypothalamic peptidergic neurons in the control of feeding

Wei, Q., Krolewski, D. M., Moore, S., Kumar, V., Li, F., Martin, B., Tomer, R., Murphy, G. G., Deisseroth, K., Watson, S. J., & Akil, H. (2018). Uneven balance of power between hypothalamic peptidergic neurons in the control of feeding. *Proceedings of the National Academy of Sciences*, 115(40), E9489–E9498.  
<https://doi.org/10.1073/pnas.1802237115>

Background


'Hypothalamus plays an essential role in the regulation of feeding behavior'

- Ventromedial hypothalamus -> 'satiety center'
- Lateral hypothalamus -> 'feeding center'

## Author's focus on the arcuate nucleus

- Agouticortin related protein (AgRP) -> orexigenic
- POMC -> anorexigenic

# Proposed mechanism




## Models

To test the previous mechanism experimental manipulation of both neuronal system is required

- POMC and NPY neurons share a common cellular origin
  - If origin is shared, then traditional transgenic manipulation
  - Virally mediated instead allows for selective POMC ne


## Model 1: viral POMC-ChR2



- ▶ Adeno-associated virus carries blue-light activation + fluorescent protein
- ▶ Not expressed because of inversion
- ▶ POMC-CRE inverts -> expression only in POMC



## Model 2: embryonic POMC-expressing progenitor control (sub-set AgRP/NPY + POMC)



- ▶ Ended up in AgRP/NPY blue light activated, green fluorescent protein model
- ▶ neuron depolarization -> cFos

With the models at hand, they tested intake dependent on differential activation


- ▶ 'Rapid inhibition of feeding behavior in fasted viral POMC-ChR2 mice'
- ▶ 'Rapid and robust increase in food intake in Tg POMC-ChR2 mice'

## Results

## Experimental setup

- ▶ ad libitum access to standard laboratory pellet before and after tests
- ▶ testes at same time of the day
- ▶ when food deprived: 4 hours + ad libitum water

# 'Rapid inhibition of feeding behavior in fasted viral POMC-ChR2 mice'



**Fig. 3.** Selective activation of POMC neurons in the Arc suppresses food intake in fasted viral POMC-ChR2 mice. (A) Blue light stimulation of POMC neurons induced a reduction of food intake in 4-h fasted viral POMC-ChR2 mice,  $F(2, 40) = 18.99$ ,  $P < 0.0001$ ; \*\*\*\* $P < 0.0001$ ;  $n = 21$  mice. (B) Light stimulation of POMC neurons led to an increased latency for the mice to lick their paws during the hot-plate test,  $F(2, 30) = 8.82$ ,  $P < 0.01$ ; \*\* $P < 0.01$ ;  $n = 16$  mice. (C) Increased latency induced by light stimulation was blunted by pretreatment with opioid antagonist naloxone (10 mg/kg) in the hot-plate test,  $F(2, 28) = 6.797$ ,  $P < 0.01$ ; \* $P < 0.05$ ; \*\* $P < 0.01$ ;  $n = 15$  mice. Repeated-measures one-way ANOVA followed by Turkey's test were used for all above statistics. Box plots show median, mean (+), lower and upper quartiles (boxes), and minima and maxima (whiskers).

- ▶ If activation of POMC-neurons is successful, then beta-endorphin + alpha-MSH should be active
- ▶ They previously showed that global POMC-neuron activation was successful via cFos marker
- ▶ We should expect (1) reduced appetite + (2) analgesic effect
- ▶ If we block opioid receptor agonist with naloxone, analgesic effect of beta-endorphin should go down

# 'Rapid and robust increase in food intake in Tg POMC-ChR2 mice'




Fig. 4. Activation of Arc neurons derived from POMC-expressing lineage increases food intake in Tg POMC-ChR2 mice. (A) Repeated-measures two-way ANOVA revealed a significant genotype  $\times$  day interaction,  $F(2, 82) = 9.44$ ,  $P < 0.001$ , for the hot-plate test. Blue light stimulation of Arc neurons in Tg POMC-ChR2 mice led to an increased latency for the animals to lick their paws,  $F(2, 52) = 21.91$ ,  $P < 0.0001$ ; \*\*\*\* $P < 0.0001$ ;  $n = 27$  mice. Latency for single transgenic littermate control mice without ChR2-eYFP expression to lick their paws was unchanged,  $F(2, 30) = 1.06$ ,  $P = 0.36$ ;  $n = 16$  mice, 8 mice per single transgenic mouse line. (B) Increased latency induced by light stimulation in Tg POMC-ChR2 mice was reduced by pretreatment with opioid antagonist naloxone,  $F(2, 24) = 15.2$ ,  $P < 0.0001$ ; \*\*\* $P < 0.001$ , \*\*\*\* $P < 0.0001$ ;  $n = 13$  mice. (C) Repeated-measures two-way ANOVA revealed a significant genotype  $\times$  day interaction,  $F(2, 54) = 76.59$ ,  $P < 0.0001$ , for the food intake study. Light stimulation of Arc neurons evoked a robust increase in food intake in Tg POMC-ChR2 mice,  $F(2, 24) = 68.28$ ,  $P < 0.0001$ ; \*\*\*\* $P < 0.0001$ ;  $n = 13$  mice. Food intake for control mice was unchanged following light stimulation of Arc,  $F(2, 30) = 1.19$ ,  $P = 0.32$ ;  $n = 16$  mice. Box plots show median, mean (+), lower and upper quartiles (boxes), and minima and maxima (whiskers).