

# Impacts of streambed dynamics on nutrient and fine sediment transport in mountain rivers

Bowyn Yager<sup>1</sup>, **Nicole Hucke<sup>1\*</sup>**, Rachel Watts<sup>2</sup>, Andrew Tranmer<sup>1</sup>, Janice Brahney<sup>2</sup>, Joel Rowland<sup>3</sup>, George Perkins<sup>3</sup> and Rose Harris<sup>3</sup>

<sup>1</sup>University of Idaho, Boise, ID

<sup>2</sup>Utah State University, Logan, UT

<sup>3</sup>Los Alamos National Laboratory, Los Alamos, NM

## Summary

- The armor layer protects the finer bed subsurface from erosion, but when dislodged during high flow events it can release fine sediment enriched in Phosphorus (P) and Organic Carbon (OC).
- Hysteresis and seasonal variations** in particulate and soluble reactive phosphorus (PP and SRP) and in particulate and dissolved organic carbon (POC and DOC) could be **controlled by armor layer motion**
- Particulate constituents such as POC and suspended sediment (SS) often show clockwise hysteresis, whereas DOC showed different hysteresis for different seasons, suggesting they are **coming from different sources**.
- We are currently investigating these sources and constraining the exact timing of armor layer motion in each event.

## Methods

### 1. Capturing Hysteresis:

Study Site: La Jara Creek,  
Valles Caldera National Preserve, NM


- Turbidity and fDOM\***  
Collected using a YSI EXO2 Sonde
- Water samples (SS, SRP, PP, DOC & POC)**  
Stage-triggered portable ISCO samplers
- Flow depth and Discharge**  
Pressure transducers in stilling wells

\* Fluorescent dissolved organic matter – a reliable proxy for DOC

**Laboratory Procedures**  
**SS** – Laser diffraction method (LISST portable XR)  
**POC** – Eurovector elemental analyzer coupled to an Isoprime IRMS  
**DOC** – OI Analytical Aurora 1030 TOC Analyzer  
**SRP & PP** – SpectraMax M2e

### 2. Fine Sediment Deposition


Sediment traps with open and closed bottoms were installed next to subsurface temperature probes during the spring of 2023




**Other deposition effects:**  
 • settling velocity  
 • stream turbulence effects

### 3. Computing Hyporheic Flux

Through diel substrate water temperature fluctuations, we can solve the vertical flux from the 1D advection-diffusion equation using  $\Delta A$  and  $\Delta \Phi$ . Temperature-monitoring probes were installed at the locations of each basket group to estimate local hyporheic flux.



## Results Water Column Fine Sediment Concentrations and $D_{50}$




**Summer high flow events** displayed a wider range of concentrations and **higher peak concentrations on average** than the spring event concentrations.

**Spring high flow events** have a wider range of  $D_{50}$ 's and a **coarser grain size distribution on average** compared to the summer samples.

Average:	<b>Spring</b>	<b>Summer</b>
Concentration ( $\mu\text{L/L}$ )	53.76	95.00
Median Grain Size ( $\mu\text{m}$ )	65.74	54.65

## Results Spring and Summer Fine Sediment Deposition

## Results Seasonal Constituent Hysteresis



A total of 7 summer and 4 snowmelt high flow events were captured for hysteresis

**Particulate constituents (SS, POC and TP) exhibited clockwise hysteresis** in the summer. SS behaved similarly in the spring

**Dissolved constituents (DOC and SRP) differed in their behavior.** Hysteresis for SRP changed for different summer storms. Hysteresis for DOC was consistently counter-clockwise in the summer and consistently clockwise in the spring. This suggests that these constituents might come from different sources

## Discussion and Further Work

- Coarser deposited grain sizes in the spring traps might be due to the transport of larger particle sizes and higher deposited sediment volume could be due to constant exposure to high flows compared to the summer storms.
- Settling velocities for water column suspended sediment samples are yet to be calculated. A more comprehensive analysis of the sediment trap deposition of each grain size class (sand, silt and clay) with hyporheic flux is in development.
- Different summer hysteresis patterns for DOC and particulate constituents suggest they are coming from different sources.
- The different degree of hysteresis for particulates could be explained by the timing of armor layer motion. This timing is currently being explored to better understand our results