

Q1

The following answers are for a Poisson train of 1000 seconds and 35Hz firing rate with a refractory period of 0ms and 5ms respectively. The Fano factor is then calculated for each window width being 10ms, 50ms and 100ms respectively. The Coefficient of Variation is also calculated.

**Refractory period: 0ms**

Window width: 10ms

Fano factor: 1.0036    Coefficient of variation: 0.9967

Window width: 50ms

Fano factor: 1.0006    Coefficient of variation: 0.9967

Window width: 100ms

Fano factor: 0.9898    Coefficient of variation: 0.9967

**Refractory period: 5ms**

Window width: 10ms

Fano factor: 0.7507    Coefficient of variation: 0.8282

Window width: 50ms

Fano factor: 0.6913    Coefficient of variation: 0.8282

Window width: 100ms

Fano factor: 0.6773    Coefficient of variation: 0.8282

Q2

The following answers are calculated from data from rho.dat. The Fano factor and the Coefficient of Variation is calculated for window width 10ms, 50ms, and 100ms respectively.

Window width: 10ms

Fano factor: 1.1176    Coefficient of variation: 2.0085

Window width: 50ms


Fano factor: 2.9297    Coefficient of variation: 2.0085

Window width: 100ms

Fano factor: 4.1029    Coefficient of variation: 2.0085

Q3

The following is an autocorrelogram over the range -100 to +100ms.

Q4

The following is a plot of the spike triggered average over a 100ms window.

