

A

Mapping 120-base QRDR of <i>gyrA</i> at nucleotides 220–339	
Segment 1 (43 bases, ~81.8°C): <i>gyrA</i> nucleotides 253–295	
GCGCCGTCGGTTGCCGAGACCATGGGCAACTA CCACCCGCACGGCGACGCGTCGATCTACGACAGCCTGGTGCGCAT GGCCAGCCCTGGTCGCTGCGCTACCCGCTGGTGGACGGCCAG	

Mapping 282-base QRDR of <i>gyrB</i> at nucleotides 1339–1620	
Segment 2 (42 bases, ~76.4°C): <i>gyrB</i> nucleotides 1340–1381	Segment 3 (22 bases, ~65.0°C): <i>gyrB</i> nucleotides 1494–1515
T CCACGGATCCGCGCAAGTCCGA ACTGTATGTCGTAGAGGT ACTCGGCCGGCGGTTCTGCAAAAAGCGGT CGCGA TTCGATGTTCCAGGCGATACTTCCGCTGCGCGGCAAGATCATCAATGTGGAGAAAGCGGCATCGACCGGTTGCTAA A GAACACCGAAGTTCAGGCGATC ATCACGGCGCTGGGCACCGGGATCCACGACGAGTTCGATATCGGCAAGCTGCGC TACCACAAGATCGTGTGATGGCCGACGCCGATGTTGACGGCCAACATATT	

B

Count	<i>gyrA</i> QRDR Variant	SSMA Screening Segment	Level of Association with Fluoroquinolone Resistance
1	Asp89Asn	1	Assoc w R
2	Gly88Cys	1	Assoc w R
3	Asp94His	1	Assoc w R
4	Asp94Tyr	1	Assoc w R
5	Ser91Pro	1	Assoc w R
6	Asp94Asn	1	Assoc w R
7	Asp94Ala	1	Assoc w R
8	Ala90Val	1	Assoc w R
9	Asp94Gly	1	Assoc w R
10	Gly88Ala	1	Interim Assoc w R

Count	<i>gyrB</i> QRDR Variant	SSMA Screening Segment	Level of Association with Fluoroquinolone Resistance
1	Asn499Asp	3	Assoc w R
2	Asn499Thr	3	Assoc w R
3	Asp461Asn	2	Assoc w R
4	Glu501Asp	3	Assoc w R
5	Glu501Val	3	Interim Assoc w R
6	Asp461His	2	Interim Assoc w R
7	Ser447Phe	2	Interim Assoc w R
8	Ala504Val	3	Interim Assoc w R