

Nicole E. Soltis¹, Susanna Atwell¹, Gongjun Shi^{1,2}, Rachel Fordyce¹, Raoni
Gwinner^{1,3}, Dihan Gao¹, Aysha Shafi¹, Daniel J. Kliebenstein^{*1,4}

¹Department of Plant Sciences, University of California, Davis, One Shields Avenue,
Davis, CA, 95616, USA

²Department of Plant Pathology, North Dakota State University, Fargo, ND, 58102, USA

³Department of Agriculture, Universidade Federal de Lavras, Lavras - MG, 37200-000,
Brazil

⁴DynaMo Center of Excellence, University of Copenhagen, Thorvaldsensvej 40, DK-
1871, Frederiksberg C, Denmark

*Correspondence: kliebenstein@ucdavis.edu Daniel J. Kliebenstein

Title: Interactions of Tomato and *Botrytis* genetic diversity: Parsing the
contributions of host differentiation, domestication and pathogen variation
~~Crop domestication~~ Domesticated tomato ~~dDomestication and pathogen virulence:~~
~~Interactions of tomato host and *Botrytis* genetic diversity~~

Short title: Interactions of tomato and *Botrytis* genetics

Material Distribution Footnote: The author(s) responsible for distribution of materials
integral to the findings presented in this article in accordance with the policy described
in the Instructions for Authors (www.plantcell.org) is (are): Daniel J. Kliebenstein
(kliebenstein@ucdavis.edu).

Abstract

Human selection during crop domestication alters numerous traits, including disease resistance. While the impacts of these processes on crop domestication on specialist pathogens are well-studied, less is known about how the interaction of crop variation and generalist pathogens affects quantitative interactions with generalist pathogens. To study how genetic variation within a domesticated standing genetic variation and domestication within a crop domestication impacts plant resistance to generalist pathogens, we infected a collection of wild and domesticated tomato accessions with a genetically diverse population of the generalist pathogen *Botrytis cinerea*. and to uncover variants in the correspondingly how this interacts with the pathogen's genetics, we infected a collection of wild and domesticated tomato accessions with a genetically diverse population of the generalist pathogen *Botrytis cinerea*. We quantified variation in lesion size of 97 *B. cinerea* genotypes (isolates) on 6 domesticated *Solanum lycopersicum* and 6 wild *S. pimpinellifolium* genotypes. Lesion size was significantly affected by large effects of the host and pathogen's genotype, with a much smaller contribution of domestication controlled by plant domestication, plant genetic variation, and the pathogen's genotype. This pathogen collection also enables Overall, resistance was slightly elevated in the wild tomato accessions. Genome-wide association (GWA) mapping in *B. cinerea*. GWA in the pathogen showed that virulence is identified a highly polygenic collection and involves a diversity of mechanisms of genes. This suggests that breeding against this pathogen would likely need to utilize a diversity of diverse isolates to capture all possible mechanisms. Critically, we identified a discrete subset of *B. cinerea* genes where the allelic variation was linked to altered virulence against the wild versus domesticated tomato accessions, as well as loci that could handle both groups. This indicates that this generalist pathogen already has the necessary a large collection of allelic variation in place to handle the introgression of wild resistance mechanisms into the domesticated crop that must be considered when designing a breeding program.

Introduction

Plant disease is mediated by complex interactions among diverse host and pathogen molecular pathways, and the disease outcome is the sum of host plant susceptibility/resistance and pathogen virulence/sensitivity mechanisms. The specific outcome of any interaction is highly dependent on the genetic variation within these pathways in both the host and pathogen. Over time, mutation and selection have led to distinct genetic architectures in the host and pathogen that are at least partly influenced by the host range of the pathogen. Specialist pathogens are a major focus in plant pathology; virulent on a narrow range of hosts, and often limited to a single species or genus. Most known plant genes for resistance to specialist pathogens confer qualitative resistance through innate immunity via large-effect loci that enable the recognition of the pathogen (Dangl and Jones 2001, Jones and Dangl 2006, Dodds and Rathjen 2010, Pieterse, Van der Does et al. 2012). These recognition signals can be conserved pathogen patterns such as cell-wall polymers or flagellin, or alternatively, specific virulence factors that block perception of the pathogen, and in turn are detected by plant proteins that guard the signaling networks (Jones and Dangl 2006, Bittel and Robatzek 2007, Ferrari, Galletti et al. 2007, Boller and He 2009, Dodds and Rathjen 2010). The evolution of large-effect qualitative loci has partly been driven by the narrow host range for the pathogen that enhances co-evolution between host resistance genes and pathogen virulence mechanisms.

In contrast to specialist pathogens, generalist pathogens are virulent across a wide range of plant host species. Generalist pathogens potentially have less stringent co-evolution to specific hosts and their accompanying resistance mechanisms, because these pathogens can easily shift to new hosts in the environment. This allows generalist pathogens to evade the rapid evolution of new resistance mechanisms within specific hosts until they evolve to counter this new resistance. This niche-shifting ability may partially explain the observation that most natural resistance to generalist pathogens is highly polygenic, and the underlying plant genes for resistance are quantitative (Glazebrook 2005, Nomura, Melotto et al. 2005, Goss and Bergelson 2006, Rowe and Kliebenstein 2008, Barrett, Kniskern et al. 2009, Corwin, Copeland et al. 2016). Plant quantitative resistance genes to generalist pathogens include a broad array of direct

defense genes, like those involved in secondary metabolite production, cell wall formation, and defense proteins (Zhang, Khan et al. 2002, Denby, Kumar et al. 2004, Zipfel, Robatzek et al. 2004, Ferrari, Galletti et al. 2007, Rowe and Kliebenstein 2008, Poland, Balint-Kurti et al. 2009, Corwin, Copeland et al. 2016). -Importantly, these quantitative plant resistance loci do not alter resistance to all genotypes (isolates) of a pathogen but interact with the infecting pathogen's genotype. For example, the ability of the *Arabidopsis* defense metabolite, camalexin, to provide resistance to *Botrytis cinerea* depends upon whether the specific isolate is sensitive or resistant to camalexin (Kliebenstein, Rowe et al. 2005, Pedras and Ahiaonu 2005, Stefanato, Abou-Mansour et al. 2009, Pedras, Hossain et al. 2011) and similarly *B. cinerea* virulence on tomato varies with the isolate's ability to detoxify tomatine (Quidde, Osbourn et al. 1998, Quidde, Büttner et al. 1999). In contrast to the polygenic nature of plant resistance to generalist pathogens, little is known about the genetic architecture of virulence within generalist pathogens, and how this is affected by genetic variation in the plant (Bartoli and Roux 2017). There are no reported naturally variable large-effect virulence loci in generalist pathogens, suggesting that virulence in generalist pathogens is largely quantitative and polygenic. This potential for interaction between polygenic virulence in generalist pathogens and equally polygenic resistance in host plants suggests that we need to work with genetic variation in both the host and pathogen to truly understand quantitative host-pathogen interactions.

A-Domestication of crop plants is a key evolutionary process in plants that has affected resistance to specialist pathogens ~~is the domestication of crop plants~~. Domesticated plant varieties are typically more sensitive to specialist pathogens than their wild relatives (Smale 1996, Rosenthal and Dirzo 1997, Couch, Fudal et al. 2005, Dwivedi, Upadhyaya et al. 2008), and pathogens may evolve higher virulence on domesticated hosts (Stukenbrock and McDonald 2008). Further, domestication typically imposes a genetic bottleneck that reduces genetic diversity in the crop germplasm, including decreased availability of resistance alleles against specialist pathogens (Tanksley and McCouch 1997, Doebley, Gaut et al. 2006, Chaudhary 2013). These general evolutionary patterns, of lower-reduced resistance and allelic diversity found when studying the interaction of specialist pathogens with crop plants, are assumed to

similarly hold for generalist pathogens and their domesticated hosts. However, there is less information about how crop host domestication affects disease caused by generalist pathogens, when the resistance to these pathogens is quantitative and polygenic rather than qualitative and monogenic. As such, there is a need to quantify the effect of domestication on a broad generalist pathogen in comparison to the rest of the crop's standing variation to test ~~conduct a detailed analysis of~~ analyze how domestication may alter the interaction of a plant with a broad generalist pathogen, and correspondingly, how and if how domestication influences the pathogen.

Botrytis cinerea provides a model generalist pathogen for studying quantitative interactions with plant hosts and underlying evolutionary processes. ~~for this generalist in contrast to specialist pathogens.~~ *B. cinerea* is a broad generalist pathogen that can infect most tested plants, from bryophytes to eudicots, and causes wide ranging pre- and post-harvest crop losses (Nicot and Baille 1996, Elad, Williamson et al. 2007, Fillinger and Elad 2015). Individual isolates of *B. cinerea* show the same broad host range ~~(Deighton, Muckenschnabel et al. 2001, Finkers, van Heusden et al. 2007, Ten Have, van Berloo et al. 2007, Corwin, Subedy et al. 2016), (Deighton, Muckenschnabel et al. 2001, Finkers, van Heusden et al. 2007, Ten Have, van Berloo et al. 2007, Corwin, Subedy et al. 2016).~~ This is in contrast to pathogens like *Fusarium oxysporum* where the species can infect diverse hosts, but each isolate is highly host specific (Katan 1999, Ormond, Thomas et al. 2010, Loxdale, Lushai et al. 2011, Barrett and Heil 2012). *B. cinerea* isolates display significant variation in virulence phenotypes, partly due to genetic variation in specific virulence mechanisms, like the production of the phytotoxins, botrydial and botcinic acid (Siewers, Viaud et al. 2005, Dalmais, Schumacher et al. 2011). This genetic variation also influences cell wall degrading enzymes and key regulators of virulence like *VELVET* that quantitatively control virulence on multiple host plants (Rowe and Kliebenstein 2007, Schumacher, Pradier et al. 2012). ~~This genetic variation in diverse~~ This standing diversity in virulence mechanisms can contribute to the formation of quantitative differences in virulence between the isolates (ten Have, Mulder et al. 1998). The phenotypic variation is driven by a high level of sequence diversity spread across the genome (Rowe and Kliebenstein 2007, Fekete, Fekete et al. 2012). The polymorphism rate in *B. cinerea*

~~was measured as, 376.6~~ SNP/kb, ~~which~~ is much more variable than most previously
studied plant pathogens (1-2 SNP/kb in *Blumeria graminis*, 1.5 SNP/kb in *Melampsora*
larici-populina, 5.5 SNP/kb in the compact genome of the obligate biotroph
Plasmodiophora brassicae, 12.3 SNP/kb in the wheat stem rust pathogen *Puccinia*
graminis f. sp. *tritici*) and human pathogens (3-6 SNP/kb in *Mycobacterium*
tuberculosis), and close to the genetic diversity that found in the human pathogen
Mycobacterium tuberculosis (2.9 to 6.2 SNP/kb) (Farhat, Shapiro et al. 2013, Hacquard,
Kracher et al. 2013, Wicker, Oberhaensli et al. 2013, Persoons, Morin et al. 2014,
Upadhyaya, Garnica et al. 2014, Atwell, Corwin et al. 2015, Desjardins, Cohen et al.
2016, Power, Parkhill et al. 2017, Atwell, Corwin et al. 2018). Higher polymorphism
rates are reported for the wheat stem rust pathogen *Puccinia graminis* f. sp. *tritici*, from
a small non-random sample of isolates (12.3 SNP/kb) (Upadhyaya, Garnica et al. 2014).
In addition to SNP diversity, the genomic sequencing showed that *B. cinerea* has a high
level of recombination and genomic admixture, as if it were a randomly intermating
population (Supplemental Figure 1) (Atwell, Corwin et al. 2018). As such, a collection of
B. cinerea isolates contains genetic variation in a wide range of virulence mechanisms,
offering the potential to challenge the host with a blend of diverse virulence
mechanisms. to identify the pathogen variation controlling quantitative virulence
(Corwin, Subedy et al. 2016, Bartoli and Roux 2017, Atwell, Corwin et al. 2018,
Fordyce, Soltis et al. 2018). ~~This can potentially identify the pathogen variation~~
~~controlling quantitative virulence, even in non-model plant systems (Bartoli and Roux~~
~~2017).~~

A model pathosystem for studying quantitative host-pathogen interactions during
domestication is the tomato-*B. cinerea* system, where the pathogen causes crop loss
due to both pre- and post-harvest infection (Dean, Van Kan et al. 2012, Hahn 2014,
Romanazzi and Droby 2016). Resistance to *B. cinerea* is a quantitative trait in tomato
as with most other species, with identified tomato QTLs each explaining up to 15% of
phenotypic variation for lesion size on stems (Diaz, ten Have et al. 2002, Finkers, van
Heusden et al. 2007, Ten Have, van Berloo et al. 2007, Rowe and Kliebenstein 2008,
Corwin, Copeland et al. 2016). Tomato is also a key model system to study how
domestication influences plant physiology and resistance, including alterations in the

circadian clock (Tanksley 2004, Bai and Lindhout 2007, Panthee and Chen 2010, Bergougnoux 2014, Müller, Wijnen et al. 2016), which can modulate resistance to *B. cinerea* (Sauerbrunn and Schlaich 2004, Weyman, Pan et al. 2006, Bhardwaj, Meier et al. 2011, Hevia, Canessa et al. 2015). This suggests that host plant [domestication diversity](#) within tomato can alter traits known [from other systems](#) to influence *B. cinerea* resistance-~~from other systems~~. Tomato domestication is typically considered a single event, followed by extensive crop improvement (Lin, Zhu et al. 2014, Blanca, Montero-Pau et al. 2015). Thus, we are using the tomato-*B. cinerea* pathosystem to directly measure the interaction of [domesticated](#) crop [domestication variation](#) with genetic variation in a generalist pathogen to better understand the evolution of this pathosystem.

In this study, we infected 97 genetically diverse *B. cinerea* isolates on a collection of domesticated tomato, *S. lycopersicum*, and wild tomato, *S. pimpinellifolium*, and quantified the interaction through lesion size in a detached leaf assay. Previous studies have examined *B. cinerea* resistance between domesticated and [distantly related](#) wild tomato species (~~i.e. *S. lycopersicum* and *S. pimpinellifolium*~~) using single isolates of pathogens (Egashira, Kuwashima et al. 2000, Nicot, Moretti et al. 2002, Guimaraes, Chetelat et al. 2004, Ten Have, van Berloo et al. 2007, Finkers, Bai et al. 2008). These previous studies typically used individual wild and domesticated tomato accessions that were the founders of mapping populations and found a wide range of *B. cinerea* resistance. However, it is still unknown how domesticated and [closely related](#) wild tomatoes compare for *B. cinerea* resistance using multiple plant genotypes and a population of the pathogen. We selected accessions to sample major geographic origins of the progenitor species, and focused the domesticated germplasm on diverse mid- to late- 20th century improved germplasm (Lin, Zhu et al. 2014, Blanca, Montero-Pau et al. 2015). In this study, we asked whether *B. cinerea* virulence was controlled by host variation, pathogen variation, or the interaction between them. Lesion size of *B. cinerea* is a quantitative trait that was controlled by plant domestication status, plant genotype and pathogen isolate. ~~We looked for evidence of specialization within our generalist pathogen population. While our *B. cinerea* isolates appear to be generalists across domestication in *Solanum*, a subset of isolates is sensitive to tomato domestication.~~

Finally, we aimed to identify the genetic basis of variation in *B. cinerea* virulence on ~~domesticated and wild tomato~~ *S. lycopersicum* and *S. pimpinellifolium*. We conducted genome-wide association (GWA) in *B. cinerea* to identify pathogen loci where genetic variation leads to altered virulence across the host genotypes, including a specific test for loci that influence responses to crop domestication. Few studies have conducted GWA in plant pathogens for virulence phenotypes, and most of these were limited by few variable loci or few genetically distinct isolates (Dalman, Himmelstrand et al. 2013, Gao, Liu et al. 2016, Talas, Kalih et al. 2016, Wu, Sakthikumar et al. 2017). ~~Our previously-sampled isolate collection includes genetic diversity across 272,672 SNPs (Supplemental Figure 1)(Atwell, Corwin et al. 2015, Zhang, Corwin et al. 2017)(Atwell, Corwin et al. 2015, Zhang, Corwin et al. 2017, Atwell, Corwin et al. 2018).~~ ~~To ensure that genetic inference was independent of the GWA method or SNP diversity reference, we repeated genetic analysis with two different association methods (bigRR and GEMMA) using SNPs called in comparison to two published *B. cinerea* genomes (T4 and B05.10). All methods converged on the same image of genetic architecture; We found that the genetic architecture of~~ virulence of *B. cinerea* is highly quantitative, with hundreds of significant SNPs with small effect sizes associated with lesion area on each tomato genotype. Importantly, there is a subset of loci in the pathogen where allelic variation gives the isolates opposing responses to crop domestication. These pathogen loci could provide tools for understanding how domestication in tomato has influenced generalist pathogen resistance, to inform breeding efforts.

Results

Experimental Design

To measure how tomato ~~domestication genetic variation~~ affects quantitative resistance to a population of a generalist pathogen, we infected a collection of 97 diverse *B. cinerea* isolates (genotypes) on wild and domesticated tomato genotypes ~~(Zhang, Corwin et al. 2017). We compared domesticated and closely related wild tomatoes for *B. cinerea* resistance using multiple plant genotypes and a population of the pathogen.~~ We selected 6 domesticated *Solanum lycopersicum* and 6 wild *S. pimpinellifolium* accessions, the closest wild relative of *S. lycopersicum*, to directly study

how domestication has influenced resistance to *B. cinerea* (Peralta, Spooner et al. 2008, Müller, Wijnen et al. 2016)(Supplemental Figure 24). Our previously collected *B. cinerea* sample includes 97 isolates obtained from various eudicot plant hosts, including tomato stem tissue (2 isolates; T3, KT) and tomato fruit (3 isolates; KGB1, KGB2, Supersteak)(Atwell, Corwin et al. 2015, Zhang, Corwin et al. 2017)(Atwell, Corwin et al. 2015, Zhang, Corwin et al. 2017, Atwell, Corwin et al. 2018). We infected all 97 *B. cinerea* isolates onto each of the 12 plant genotypes in 3-fold replication across 2 independent experiments in a randomized complete block design, giving 6 measurements per plant-pathogen combination, for a total of 3,276 lesions. Digital measurement of the area of the developing lesion provides a composite phenotype controlled by the interaction of host and pathogen genetics. This measurement of the plant-*B. cinerea* interaction has been used successfully in a number of molecular and quantitative genetic studies (Ferrari, Plotnikova et al. 2003, Denby, Kumar et al. 2004, Kliebenstein, Rowe et al. 2005, Ferrari, Galletti et al. 2007, Ten Have, van Berloo et al. 2007, AbuQamar, Chai et al. 2008, Rowe and Kliebenstein 2008, Liu, Hong et al. 2014). It should be noted that we are not focusing on MAMP or PAMP specific host/pathogen interactions with this study; we are instead allowing the identification of any mechanism that may influence the host/pathogen interaction including metabolism, development or any other unknown component. If there is genetic variation affecting the trait, and the trait influences the interaction of host and pathogen, it will be a component of the experiment. This fits with the recently developing view that growth, development and resistance in plants are highly integrated processes that may not be as distinct as once believed (Campos, Yoshida et al. 2016, Ballaré and Pierik 2017, Züst and Agrawal 2017, Izquierdo-Bueno, González-Rodríguez et al. 2018).

Lesion size (phenotypic) variation

We collected images of all lesions at 24, 48, and 72 hours post inoculation. At 24 hours, no visible lesions were present on the tomato leaves. At 48 hours, a thin ring of primary lesion became visible surrounding the location of the spore droplet, but no expansion was visible. At 72 hours significant lesion growth was visible, but no lesions had spread to infect over half of the leaflet. We digitally measured the area of all developing lesions

at 72 hours post infection (HPI) as a measure of virulence (Figure 1). We use the linear measurement of lesion area for several reasons. First, According in previous work to previous studies, at 72 HPI *B. cinerea* lesion area growth appears to enter a relatively linear growth phase (Rowe, Walley et al. 2010); Secondly, and previous research from our group has shown that the linear measurement behaves as a normally distributed trait modeled lesion area as a linear or log-linear trait (Kliebenstein, Rowe et al. 2005, Corwin, Copeland et al. 2016, Fordyce, Soltis et al. 2018) (Kliebenstein, Rowe et al. 2005, Corwin, Copeland et al. 2016, Atwell, Corwin et al. 2018, Fordyce, Soltis et al. 2018). And finally, previous work has shown that Botrytis isolates display large variation in their unit biomass per area lesion area and as such growth in biomass is not the sole factor driving this measure Lesion area is a common measure of the interaction of plant immunity and pathogen virulence, though pathogen biomass does not scale with lesion area (Corwin, Subedy et al. 2016). (Rowe, Walley et al. 2010, Corwin, Copeland et al. 2016) We observed a mean lesion size of 0.67 cm² across the full experiment, with 0.94 CV across the full isolate population on all tomato genotypes. Individual isolates were highly variable in their lesion size across tomato genotypes (Figure 1-c-h), with mean lesion size per isolate of 0.14 cm² to 1.29 cm², and individual isolate coefficient of variation (CV) from 0.51 to 1.68 across all observations on all tomato genotypes (Supplemental Data Set 1). A subset of these isolates is highly virulent on tomato (mean lesion size > 1.05 cm², Figure 1e), and a subset can be considered saprophytic (mean lesion size < 0.3 cm², Figure 1f). Lesion size of *B. cinerea* on tomato was weakly showed a weak positive correlation with lesion size on *A. thaliana* from previous studies; both on domesticated tomato (r=0.247, p= 0.003) and on wild tomato (r=0.301, p= 0.016) (Supplemental Figure 3) (Zhang, Corwin et al. 2017). This lack of correlation suggests the presence of at there are both shared and unique mechanisms of quantitative virulence in the two species.

Contribution of Pathogen Genetics, and Plant Genetics and Crop Domestication Effects on Resistance

To measure the relative contribution of genetic diversity in the plant and the pathogen to variation in the virulence/ susceptibility phenotype, we used a ~~multiple linear regression~~general linear model (~~R Development Core Team 2008~~) (R lme4 package; (Bates, Maechler et al. 2015)). This model directly tested the contribution of pathogen genotype (isolate), plant genotype, and plant domestication status, ~~and pathogen genotype (isolate)~~ to variation in lesion size. The final model showed that genetic variation within both the host plant and the pathogen had significant effects on lesion growth, ~~with pathogen isolate diversity explaining 3.5 fold4 more variance than plant genotype each explaining approximately the same portion of the variance~~, ~~46% of total genetic variance for pathogen isolate vs. 13% for plant genotype~~ (Table 1 and Figure 1c). Interestingly, while tomato domestication status significantly impacted *B. cinerea* virulence, ~~as it was to a much lower level than the other factors shown by the small but significant effects of genetic variation between domesticated and wild tomatoes (3.5% of total genetic variance~~, Table 1). There was no evidence for significant interaction effects between pathogen isolate and plant genotype, ~~but this term contributed the largest proportion of the plant related variance in lesion size (34% of total genetic variance, Table 1). The lack of significance for this term in face of the large fraction of variance may be due to the vast degrees of freedom in this term (Table 1).~~ Thus, the interaction between tomato and *B. cinerea* was significantly controlled by genetic diversity within the host plant and the pathogen, including a slight effect of domestication status.

Table 1. ANOVA results of the interaction between 12 tomato accessions and 95 *B. cinerea* isolates measured as lesion area.

Results of general linear modelling of lesion area for 12 tomato accessions by 95 *B. cinerea* isolates is shown (R lme4 package version 1.1-18-1; (Bates, Maechler et al. 2015)). Two of our 97 isolates did not have replication across 2 experiments, so they were dropped at this stage of analysis. The terms are as follows; Isolate is the 95 *B. cinerea* isolates, Domestication is wild tomato, *S. pimpinellifolium*, versus domesticated tomato, *S. lycopersicum*, Plant is 12 tomato genotypes nested within their respective domestication groupings, Experiment tests the random effect of 2 independent replicate experiments. The nested random effects of whole plant sampled, leaf sampled, and leaflet pair are included. In addition, interactions of these factors are tested (:). The degrees of freedom and p-value are shown. For fixed effects, the type II sum of squares and F-value are shown, and for random effects the likelihood ratio test statistic (LRT) is shown.

<u>Fixed Effect</u>	<u>SS</u>	<u>F value</u>	<u>DF</u>	<u>p</u>
<u>Isolate</u>	<u>37.8</u>	<u>1.7</u>	<u>94</u>	<u>0.007</u>
<u>Domestication</u>	<u>3.4</u>	<u>14.1</u>	<u>1</u>	<u>0.0006</u>
<u>Domest/Plant</u>	<u>39.3</u>	<u>16.2</u>	<u>10</u>	<u>5e-11</u>
<u>Iso:Domest</u>	<u>15.8</u>	<u>0.7</u>	<u>94</u>	<u>0.99</u>
<u>Iso:Domest/Plant</u>	<u>179.1</u>	<u>0.8</u>	<u>940</u>	<u>1</u>
<u>Random Effect</u>	<u>LRT</u>	<u>DF</u>	<u>p</u>	
<u>1 Experiment</u>	<u>136</u>	<u>1</u>	<u><2e-16</u>	
<u>1 Whole Plant</u>	<u>0.21</u>	<u>1</u>	<u>0.65</u>	
<u>1 WP/Leaf</u>	<u>22.4</u>	<u>1</u>	<u>2e-06</u>	
<u>1 WP/Leaf/Leaflet Pair</u>	<u>0</u>	<u>1</u>	<u>1</u>	
<u>1 Exp:Iso</u>	<u>321</u>	<u>1</u>	<u><2e-16</u>	

Pathogen Specialization to Source Host

One evolutionary model of plant-generalist pathogen interactions suggests that ~~generalist~~ pathogen isolates within ~~a-a generalist~~ species may specialize for interaction with specific hosts. Alternatively, generalist isolates may show no host specialization or preference. Our collection of *B. cinerea* includes five isolates that may be adapted to tomato, as they were collected from *S. lycopersicum*. To test if there is evidence for specialization to the source host, we compared the virulence of the *B. cinerea* isolates obtained from tomato to the broader pathogen population. For *B. cinerea* genotypes isolated from tomato tissue vs. other hosts, there was no significant difference in lesion size ~~on domesticated tomato (t-test; t=1.10, n = 97, p=0.33), wild tomato (t-test; t=1.09, n = 97, p=0.33) or~~ across all tomato genotypes (t-test; n = 97, p=0.14) (Figure 1g). In fact, one isolate collected from tomato tissue (KGB1) was within the 10 least-virulent isolates and another (Triple3) was within the 10 most-virulent isolates (Figure 1g). This demonstrated significant genetic variation in virulence across the *B. cinerea* isolates, and that this collection of *B. cinerea* isolates from tomato do not display a strong host-specificity for tomato (Martinez, Blancard et al. 2003, Ma and Michailides 2005, Rowe and Kliebenstein 2007, Samuel, Veloukas et al. 2012).

Pathogen Specialization to Host Genotype

Though we did not find evidence for *B. cinerea* ~~adaptation to~~ preference for tomato based on isolate host source, the *B. cinerea* isolates may contain genetic variation at individual loci that allow them to better attack subsets of the tomato genotypes (Rowe and Kliebenstein 2007, Kretschmer and Hahn 2008, Corwin, Subedy et al. 2016). A visual analysis of the data suggested an interaction between the genomes of *B. cinerea* and tomato (Figure 1 c-h). However, when using the full model, we found no significant interaction between isolate and individual host genotype, even though there was a large fraction of variance within these terms (Table 1). This may indicate a lack of interaction between genetic variation in the host and pathogen. Interaction effects in large datasets can be difficult to identify using mixed models, so we used a second standard statistical approach, a Wilcoxon signed-rank test, ~~to~~ We used model-adjusted lesion sizes as input to test if the rank of *B. cinerea* isolate-induced lesion size significantly changes between pairs of tomato genotypes. This showed that when using the full isolate population, the rank performance of the isolates does significantly vary between host genotypes. When comparing mean lesion size between paired plant genotypes, ~~58.59%~~ 58.59% (~~38-39~~ out of 66) of tomato accession pairs had significantly different ranking of the isolates (Wilcoxon signed-rank test with Benjamini-Hochberg FDR-correction, Table 2, Supplemental Figure 42). A significant p-value indicates that the two host genotypes show evidence for different virulence interactions with the population of *B. cinerea* isolates, providing evidence for host x pathogen genotypic interactions. This pattern was consistent across domesticated host pairs, wild host pairs, or between-species host pairs (Wilcoxon signed-rank test with B-H FDR-correction, Table 2). This provides evidencesuggests that the population of *B. cinerea* does display differential responses to the tomato genetic variation.

To focus on whether specific *B. cinerea* isolates may be sensitive to tomato domestication, we applied a Wilcoxon and ANOVA approach. Overall, most isolates (78/97, 80%) are more virulent on domesticated than wild tomato (Figure 3; Supplemental Data Set 1). ~~Using a The~~ Wilcoxon signed-rank test, to compare the rank of model-corrected mean lesion size of all the *B. cinerea* isolates on wild versus domesticated tomato, ~~was showed we found a significant differencesignificant~~ (Wilcoxon signed-rank test, $W = 59465801$, $p\text{-value} = 0.0020007$) (Figure 3). While this shows a

general population behavior, we used single-isolate ANOVAs ~~To to identify test if any~~
~~specific~~ the pathogen genotypes ~~most had a significant sensitive to strongly associated~~
~~with domestication~~ association with domestication. ~~, we conducted single-isolate~~
~~ANOVAs on~~ These general linear models ~~including included~~ the fixed effects of plant,
domestication, and ~~the random effect of~~ experiment. ~~, After adjusting for multiple testing,~~
~~this identified and found~~ two isolates (Fd2, Rose) with a significant effect of
domestication on lesion size ($p < 0.05$, FDR corrected) (Figure 1h), both of which are
more virulent on domesticated tomato (Supplemental Data Set 3). ~~These included one~~
~~of the highly virulent isolates (Fd2), and one of the largely saprophytic isolates (Rose),~~
~~which suggests that isolate virulence level on tomato does not predict *B. cinerea* genetic~~
~~response to tomato domestication. Both of these isolates were more virulent on~~
~~domesticated than on wild tomato. These results suggest that this *B. cinerea* population~~
~~contains two highly domestication-sensitive associated isolates which are more virulent~~
~~on domesticated tomato, and a broader pattern of *B. cinerea* sensitivity to tomato~~
~~genetic variation.~~

To assess whether isolates could appear domestication-associated due to
random chance, we bootstrapped assignment of plant accessions to domestication
groups. ~~Across the 100 bootstraps, we~~ 96 of the 100 bootstraps identified no isolates
~~with domestication sensitivity, and the other four bootstraps identified only identified 2~~
~~isolates showing significant domestication association (FDR < 0.01) to domestication in~~
~~4% of the random datasets.~~ Therefore, our individual isolate observations are in the
96th percentile. ~~This While this is suggestive, suggests that a more precise estimate of~~
~~isolate x domestication interactions would require larger experiments using either more~~
~~replication or additional plant genotypes.~~

Table 2. Rank order shifts of 97 *B. cinerea* isolates by lesion area across all of the tomato accessions.

Wilcoxon signed-rank test on comparing ~~model-corrected~~ mean *B. cinerea* lesion area
on tomato accessions. This tests for a change in the rank order of the 97 isolates
between each pair of tomato accessions. A significant p-value suggests that the relative
performance of individual isolates is altered from one host to the other. The lower left
corner of the chart includes B-H FDR-corrected p-values, the upper right corner
includes the test statistic (W). Bold text indicates significance at $p < 0.01$ after
correction, italicized text indicates suggestive p-values $0.01 < p < 0.1$. NS shows non-
significant interactions.

		Wild					Domesticated						
		LA1547	LA1589	LA1684	LA2093	LA2176	LA480	LA2706	LA3008	LA3475	LA410	LA4345	LA4355
Wild	LA1547		2978	3988	2927	1865	3008	1710	3460	1597	1135	3928	2944
	LA1589	<0.001		5401	4699	3359	4662	3014	4918	2938	2340	5536	4454
	LA1684	NS	0.029		3709	2552	3690	2296	4004	2205	1690	4537	3571
	LA2093	<0.001	NS	0.049		3013	4496	2732	4889	2588	1947	5534	4264
	LA2176	<0.001	0.004	<0.001	<0.001		5837	4029	6002	3963	3276	6706	5583
	LA480	<0.001	NS	0.044	NS	0.001		6143	4192	6286	6855	3575	4702
Domesticated	LA2706	<0.001	<0.001	<0.001	<0.001	NS	<0.001		6311	4523	3876	6917	5940
	LA3008	0.009	NS	NS	NS	<0.001	NS	<0.001		2619	2082	5100	4049
	LA3475	<0.001	<0.001	<0.001	<0.001	NS	<0.001	NS	<0.001		3815	7088	5984
	LA410	<0.001	<0.001	<0.001	<0.001	0.002	<0.001	NS	<0.001	NS		7567	6602
	LA4345	0.16	0.011	NS	0.011	<0.001	0.021	<0.001	NS	<0.001	<0.001		3439
	LA4355	<0.001	NS	0.02	NS	0.008	NS	<0.001	NS	<0.001	<0.001	<0.001	

Domestication and Lesion Size Variation

Existing literature predominantly reports that crop domestication decreases plant resistance to pathogens (Smale 1996, Rosenthal and Dirzo 1997, Couch, Fudal et al. 2005, Dwivedi, Upadhyaya et al. 2008, Stukenbrock and McDonald 2008). While we did observe the expected decreased resistance (by 18%) in domesticated tomato (Figure 2 and 3, Table 1), domestication was a minor player in controlling lesion size variation, with most of the plant genetic signature coming from variation within both the wild and domesticated tomato species, contributing 12-fold more variation in resistance than domestication alone (Table 1). ~~In our analysis, we identified a significantly greater (18%) resistance of wild tomato in comparison to domesticated tomato across the~~

population of *B. cinerea* isolates (Figure 2 and 3, Table 1). However, this domestication effect was not the dominant source of variation, as genetic variation within the domesticated and wild genotypes contributed 3.8-fold more variation in resistance than domestication alone (Table 1). While we did observe the expected decreased resistance in domesticated tomato, domestication was a minor player in controlling lesion size variation, with most of the plant genetic signature coming from variation within both the wild and domesticated tomato species. Removing the two domestication-associated isolates (Fd2, Rose) from our population did not eliminate the effect of tomato domestication on lesion size, as Domestication still accounted for 1.2% of genetic variation, with a significant effect on lesion size as it was still significant and *B. cinerea* was still more virulent on domesticated tomato by 17% (Supplemental Table 1). ——— To test how this mild domestication effect might be sensitive to shifts in the collection of tomato genotypes, we used the bootstrapping of domesticationsame bootstraps from above for the full model from above. Our observed domestication effect was in the top 80th percentile across all bootstraps, suggesting that itwhile the domestication effect is small, it is relatively stable in response to shifts in the genotypes. However, a larger sample of *S. lycopersicum* and *S. pimpinellifolium* genotypes would be needed to develop a more precise estimate of any domestication effect on lesion size.

In addition to altering trait means, domestication commonly decreases genetic variation in comparison to wild germplasm due to bottlenecks, including for tomato (Tanksley and McCouch 1997, Doebley, Gaut et al. 2006, Bai and Lindhout 2007). We would expect this decreased genetic variation to limit phenotypic variation, including disease phenotypes. Interestingly in this tomato population, we did not observe reduced variation in lesion size in the wild-domesticated tomato. Rather, the domesticated tomato genotypes had a wider range of average lesion size than wild genotypes; the 90th percentile range (95th percentile to 5th percentile) spanned 2.03 cm² lesion size variation on domesticated tomato (standard deviation = 0.68 cm²) versus 1.76 cm² variation on wild tomato (standard deviation = 0.58 cm²). Additionally, the The wild and domesticated tomato genotypes showed statistically similar variation in resistance (F-test, $F_{96,96}=1.39$, $p=0.11$) (Figure 3, Supplemental Figure 24). Overall, there is a slight

domestication impact on average resistance to *B. cinerea*, ~~but and~~ no evidence of a phenotypic bottleneck due to domestication. This suggests that in the tomato-*B. cinerea* pathosystem, domestication is not a major part of the variation.

Quantitative Genetics of Pathogen Virulence on Tomato

Genetic variation within *B. cinerea* had a large effect on virulence on tomato and showed some evidence for interactioned with tomato domestication (Table 1). This suggests that there is genetic variation within the pathogen, in which some alleles enhance, and other alleles decrease virulence depending upon the plant's genotype. To identify variable pathogen genes controlling differential virulence across plant genotypes, we conducted GWA mapping analysis within the pathogen ~~by two approaches, using 272,672 SNPs compared to the *B. cinerea* T4 reference genome (Supplemental Figure 1)~~ (Atwell, Corwin et al. 2018). Due to the large effect of plant genotype on resistance to *B. cinerea*, we performed GWA using ~~the~~ model-corrected least-squared mean virulence measured on each tomato genotype as separate traits. We ~~first~~ used a ridge-regression approach (bigRR) ~~in combination with 272,672 SNPs from *B. cinerea* compared to the T4 reference genome~~ to estimate the phenotypic effects across the genome ~~(Shen, Alam et al. 2013, Corwin, Copeland et al. 2016, Corwin, Subedy et al. 2016, Francisco, Joseph et al. 2016)~~ (Shen, Alam et al. 2013, Corwin, Copeland et al. 2016, Corwin, Subedy et al. 2016, Francisco, Joseph et al. 2016, Atwell, Corwin et al. 2018). ~~To verify these patterns, we also implemented a Genome-wide Efficient Mixed Model Association (GEMMA) analysis with a centered relatedness matrix to control for the effects of population structure (Zhou and Stephens 2012). In GEMMA, we included 237,878 SNPs from *B. cinerea* compared to the B05.10 reference genome.~~ To determine significance of SNP effects under ~~both~~ GWA methods, we permuted phenotypes 1000 times to calculate 95, 99, and 99.9% effect size thresholds within each plant host. At Under 1000x permutations, the 99.9% threshold is imprecise, but we included this approximate threshold to more conservatively identify conservatively SNP associations. ~~Under both methods,~~ GWA analysis showed that the genetic basis of *B. cinerea* virulence on tomato is highly polygenic. Consistent with a polygenic structure of this trait in the pathogen, ~~neither method of GWA identified~~ GWA

499 ~~did not identify~~ large-effect SNPs (Figure 4). The ~~number of significant *B. cinerea*~~
500 ~~virulence SNPs identified by this~~ ridge-regression approach (bigRR) ~~identified~~
501 ~~from~~ ~~varied by plant accession, from~~ 1,284 to 25,421 SNPs ~~on within *B. cinerea* that~~
502 ~~were significantly associated with altered virulence on~~ the 12 different host genotypes
503 (significance was determined by the SNP effect size estimate exceeding the 99% 1000-
504 permutation threshold). ~~The model accounting for population structure (GEMMA)~~
505 ~~confirmed our finding of a highly polygenic nature of lesion size in the pathogen~~
506 ~~(Supplemental Figure 3), with 2,530 to 8,221 SNPs significantly associated with~~
507 ~~virulence at the 99% threshold, and 288 to 1,361 SNPs at the 99.9% threshold~~
508 ~~(significance was determined using an empirically determined 1000-permutation~~
509 ~~threshold).~~

510 At the SNP level, fewer loci contribute to virulence across all hosts ~~genotypes~~.
511 We found five *B. cinerea* SNPs significantly linked to altered lesion size on all 12 tomato
512 accessions (Figure 4b). 215 SNPs were called in at least ten hosts, and 3-,300k SNPs
513 were called in at least half of the hosts while 27% (46,000) of the significant SNPs were
514 linked to virulence on only a single host tomato genotype. These levels of overlap
515 exceed the expected overlap due to random chance (Figure 5a). While only a small
516 subset of these *B. cinerea* SNPs were linked to virulence on all the tomato genotypes,
517 we obtained better overlap across host genotypes by focusing on gene windows.

518 To focus on the small-effect genes linked to *B. cinerea* virulence, we classified a
519 gene as significantly associated if there was 1 SNP linked to a trait using a 2kbp window
520 surrounding the start and stop codon for a given gene. This analysis identified 14 genes
521 linked to differential virulence in all 12 tomato accessions by bigRR (Figure 5b,
522 Supplemental Data [Set 2-ae](#)), as some SNPs within a gene had accession-specific
523 phenotypes (significant in <12 tomato accessions). A further 1045 genes were linked to
524 differential virulence on 7 to 11 of the tomato accessions by bigRR (Figure 5b,
525 Supplemental Data [Set 2a-e](#)). ~~At the 99.9% SNP significance threshold, GEMMA~~
526 ~~identified 23 genes across 7 to 9 of the tomato accessions (Supplemental Figure 4).~~

527 Of the 14 genes with SNPs significantly associated with *B. cinerea* virulence on
528 all tomato genotypes by bigRR, most have not been formally linked to pathogen
529 virulence. However, SNPs within a pectinesterase gene (BcT4_6001, Bcin14g00870)

were associated to virulence across 11 tomato accessions. Pectinesterases are key enzymes for attacking the host cell wall, suggesting that variation in this pectinesterase locus and the other loci may influence pathogen virulence across all the tomato genotypes (Valette-Collet, Cimerman et al. 2003). Therefore, [as an example of a virulence gene identified by our GWA methods](#), we looked for evidence of multiple haplotypes in this locus linked to virulence by visualizing the SNP effects across the pectinesterase gene. We plotted the effect sizes for all SNPs in this gene and investigated the linkage disequilibrium amongst these SNPs (Figure 6). This showed that the effect of SNPs across this gene vary in effect direction depending on tomato host genotype (Figure 6a). We identified two haplotype blocks contributing to the association of this gene to the virulence phenotype (Figure 6b). One block is associated with SNPs in the 5' untranslated region in SNPs 5-11, and the second block is SNPs that span the entirety of the gene in SNPs 13-26. Interestingly, there are only two SNPs in the open reading frame of the associated gene (Figure 6). This suggests that the major variation surrounding this locus is controlling the regulatory motifs for this pectinesterase. Thus, there is significant genetic variation in *B. cinerea* virulence that is dependent upon the host's genetic background. This suggests that the pathogen relies on polygenic small effect loci, potentially allowing selection to customize virulence on the different tomato hosts.

~~To identify genes consistently associated with *B. cinerea* virulence on tomato across GWA methods, we examined the gene overlap between significant associations identified by GEMMA on the B05.10 genome and bigRR on the T4 genome. We conservatively identified genes within 2kb of significant SNPs at the 99% permutation threshold for bigRR, and at the 99.9% permutation threshold for GEMMA. Among these, 263 genes were linked to at least two plant genotypes by both methods (Supplemental Data 2 a). These genes include transporters and enzymes that can be important for Botrytis toxin production and/or detoxification of plant defense compounds and are key to virulence. Other known and predicted pathogen virulence functions were largely not identified (Supplemental Data 2 a, c).~~

~~At the SNP level, fewer loci contribute to virulence across all hosts and both GWA methods. We found five *B. cinerea* SNPs significantly linked to altered lesion size~~

on all 12 tomato accessions using the bigRR analysis (Figure 4b). 215 SNPs were called in at least ten hosts, and 3.3k SNPs were called in at least half of the hosts while 27% (46,000) of the significant SNPs were linked to virulence on only a single host tomato genotype. These levels of overlap exceed the expected overlap due to random chance (Figure 5a). GEMMA analysis also found significant SNP overlap between hosts at the 99% permutation threshold, with 89 SNPs in at least ten hosts, 859 SNPs in at least half of the hosts, and 63% (19,270) of significant SNPs unique to a single host. SNP calling between hosts was lower for GEMMA at the 99.9% permutation threshold, with 78% of significant SNPs (4269) in a single host, and 38 SNPs significant across at least half of the hosts (Supplemental Figure 4 a). While only a small subset of these *B. cinerea* SNPs were linked to virulence on all the tomato genotypes, we obtained better overlap across host genotypes by focusing on gene windows.

Quantitative Genetics of Pathogen Response to Tomato Domestication

The identification of two isolates that ~~distinctly respond~~differed on wild and domesticated tomato to tomato domestication suggests that there is indicated that there may be some natural genetic variation in *B. cinerea* ~~that is affected by tomato domestication~~linked to this phenotypic variation. To directly map *B. cinerea* genes that control differential virulence on wild versus domesticated ated tomatoes, we used the least-squared mean virulence of each isolate across all wild and all domesticated tomato genotypes as two traits. We also calculated a domestication sensitivity trait; the relative difference in lesion size for each isolate between domesticated and wild hosts. Using these three traits, we conducted bigRR GWA within *B. cinerea* to map genes in the pathogen that respond to domestication shifts in the plant. Using the mean lesion area of the *B. cinerea* isolates on the wild or domesticated tomato hosts identified a complex, highly polygenic pattern of significant SNPs, similar to the individual tomato accessions (Figure 4, Figure 7). ~~This~~The significant SNP sets had a high degree of overlap between the wild phenotype and domesticated phenotype. In contrast, ~~the~~ Domestication Sensitivity ~~trait~~identified a much-more limited set of SNPs ~~that had~~with less overlap ~~with~~to the mean lesion area on either Domesticated or Wild tomato (Figure 7). ~~GWA of these domestication traits by GEMMA identified similar patterns of polygenic~~

structure, high overlap between SNPs and genes on wild or domesticated tomato hosts, and rare overlap with Domestication Sensitivity (Supplemental Figure 5). To ~~begin~~ ~~queryingquery~~ the underlying gene functions for these ~~various~~ *B. cinerea* loci, we called genes as significant if there was one SNP within 2kb of the gene (Figure 7c). ~~We also examined the genes associated with these domestication virulence traits found by both bigRR and GEMMA. This overlap identified 200 unique genes including several transporters and enzymes, with few predicted virulence genes (Supplemental Data 2 b). One gene from this overlap list (Bcin01g05800) contains TPR repeats, which are common in bacterial virulence proteins (Cerveny, Straskova et al. 2013) and are among the proteins secreted by the plant pathogen Ustilago maydis (Lo Presti, López-Díaz et al. 2016).~~ Using all 1251 genes linked to domestication traits by bigRR for a functional enrichment analysis found only 22 significantly overrepresented biological functions (Fisher exact test, $p < 0.05$, Supplemental Data [Set 2b-f](#)) when compared to the whole-genome T4 gene annotation. ~~We also examined functional enrichment for the genes associated with domestication traits by both GEMMA and bigRR. We found 41 significantly overrepresented biological functions (Supplemental Data 2 d). In both datasets, the~~ ~~The~~ enrichments were largely surrounding enzyme and transport functions, which are known to be key components of how the pathogen produces toxic metabolites and conversely detoxifies plant defense compounds. Thus, there is an apparent subset of *B. cinerea* genes that may be specific to the genetic changes that occurred in tomato during domestication. Further work is needed to assess if and how variation in these genes may link to altered virulence on domestic~~ated~~ and wild tomatoes.

Discussion

The genetics of plant resistance to generalist pathogens are mostly quantitative, depend upon pathogen isolate, and rely on genetic variation in both signal perception and direct defense genes (Kover and Schaal 2002, Parlevliet 2002, Glazebrook 2005, Nomura, Melotto et al. 2005, Goss and Bergelson 2006, Tiffin and Moeller 2006, Rowe and Kliebenstein 2008, Barrett, Kniskern et al. 2009, Corwin, Copeland et al. 2016, Zhang, Corwin et al. 2017). Previous studies ~~on-of~~ tomato resistance to *B. cinerea* have

found a quantitative genetic architecture that varies between domesticated and wild tomato species, with higher resistance in the wild species (Egashira, Kuwashima et al. 2000, Nicot, Moretti et al. 2002, Guimaraes, Chetelat et al. 2004, Finkers, van Heusden et al. 2007, Ten Have, van Berloo et al. 2007, Finkers, Bai et al. 2008). However, it was not known how the choice of *B. cinerea* isolate may change this plant-pathogen interaction. To address these questions, we used genetic variation in wild and domesticated tomato accessions in conjunction with a population of *B. cinerea* isolates. ~~This also allowed us to test how domestication variation within tomato influenced the interaction at the level of the pathogen population and individual genes in the pathogen.~~ *B. cinerea* virulence on tomato, as measured by lesion size, was significantly affected by pathogen isolate, host genotype, and domestication status (Table 1). Pathogen isolate and tomato genotype were the strongest determinants of the interaction with only ~~Tomato domestication led to~~ a slight but significant decrease in resistance to the pathogen associated with domestication. ~~but critically~~ Equally, there was no evidence of a domestication bottleneck, with similar variance in resistance between the wild and domesticated tomato accessions (Table 1, Figure 2). There was also little evidence in this *B. cinerea* population for specialization to tomato, supporting the hypothesis that *B. cinerea* is a generalist at the isolate and species level (Figure 1 c-h) (Giraud, Fortini et al. 1999, Martinez, Blancard et al. 2003, Ma and Michailides 2005). GWA mapping within the pathogen showed that the genetics underlying *B. cinerea* virulence on tomato are highly quantitative and vary across tomato genotypes and domestication status (Figure 5, Figure 7). This analysis identified a small subset of pathogen genes whose variation contributes to differential virulence on most of the hosts tested, and a set of pathogen genes whose variation is responsive to tomato domestication (Supplemental Data Set 2 b, d, fb). ~~We also identified a conservative subset of genes whose association to differential *Botrytis cinerea* virulence is consistent across GWA methods and reference genomes (Supplemental Data 2 a, b, c, d).~~

Domestication and altered pathogen virulence genetics

~~These results provide evidence of a mild tomato domestication effect on resistance to the generalist pathogen, *B. cinerea*. We measured an 18% increase in~~

susceptibility across domesticated varieties, but this represents explains less than 1% of
the a small portion of the total variance of *B. cinerea* lesion size on tomato (Table 1a).
As such, domestication status alone is a poor predictor of a specific tomato host's
resistance to infection by *B. cinerea*. This suggests that while tomato domestication
does affect this plant-pathogen interaction, it is not the primary factor defining the
measured trait. Functional plant defenses within both the domesticated and wild tomato
accessions may act to buffer variation in *B. cinerea* virulence, as has been shown for *A.*
thaliana (Zhang, Corwin et al. 2017). The effect of tomato domestication varied across
the *B. cinerea* isolates, with specific loci linked to differential virulence across wild and
domesticated tomatoes (Figure 1 c-h, Figure 7, Supplemental Data Set 3, Supplemental
Data Set 4)). If a study relies on one or a few isolates, it could obtain a falsely high or
falsely low estimation of how host domestication influences pathogen resistance. This
shows the As such, future studies need to utilize a population of *B. cinerea* to
understand the factors contributing to *B. cinerea* virulence and how this is altered by
crop domestication.

In biotrophic pathogens, host domestication has decreased the diversity of
resistance alleles because they are lost in the domestication bottleneck as found for
specialist pathogens (Tanksley and McCouch 1997, Doebley, Gaut et al. 2006, Hyten,
Song et al. 2006, Chaudhary 2013). Surprisingly, we did not find evidence for a
domestication bottleneck in the phenotypic resistance to *B. cinerea* (Figure 2, Figure 3).
This is in contrast to genomic studies that explicitly show a genotypic bottleneck within
tomato domestication (Miller and Tanksley 1990, Koenig, Jiménez-Gómez et al. 2013).
Previous work in *A. thaliana* with these isolates has shown that if plant defenses such
as jasmonic acid and salicylic acid signaling are non-functional, there is increased
variation in *B. cinerea* virulence (Zhang, Corwin et al. 2017). Conversely, if major plant
defense Thus, if these pathways such as jasmonic acid and salicylic acid lost
functionality in had large effect differences between wild and domesticated tomato
tomato domestication, we would expect to see a wider range of *B. cinerea* virulence
phenotypes in domesticated tomato (Zhang, Corwin et al. 2017). However The similarity
in the variance suggests that any differences we are seeing are not caused by large
effect changes that abolish or greatly diminish specific defense signaling networks, we

~~also do not find evidence of phenotypic plasticity in domesticated tomato resistance to *B. cinerea* (Figure 2, Figure and 3). This suggests that at least for this generalist pathogen, the genetic bottleneck of tomato domestication has not imparted a phenotypic bottleneck. One possible explanation is that resistance to this pathogen is so polygenic in the plant that our experiment is not sufficiently large to pick up any genetic bottleneck effect using phenotypic variance.~~ These patterns, of mild decrease in resistance to *B. cinerea* due to plant domestication, and within-species plant variation exceeding the contribution of domestication itself, may be unique to interactions between *B. cinerea* and tomato, or more general. ~~It remains to be seen if these patterns hold for *B. cinerea* on its other host plants.~~ It is unclear whether ~~domestication this pattern is unique to tomato has a universal effect on plant resistance to *B. cinerea*,~~ or if each domestication event is unique.

Polygenic quantitative virulence and breeding complications

Our results indicate a highly polygenic basis of quantitative virulence of the generalist *B. cinerea* on tomato. ~~similar to the Previous studies have found a highly polygenic basis of *B. cinerea* quantitative virulence on the host side of the interaction~~ (Zhang, Corwin et al. 2017). The variation in lesion size is linked to numerous *B. cinerea* SNPs, each with small effect sizes (Figure 4a). Importantly, the tomato host accession greatly influenced which *B. cinerea* loci were significantly associated to lesion size (Figure 5). Thus, it possible that there is specialization at the gene level, in which different alleles within the pathogen link to differential virulence on specific host genotypes (Giraud, Fortini et al. 1999, Rowe and Kliebenstein 2007, Blanco-Ulate, Morales-Cruz et al. 2014). This polygenic architecture of virulence is ~~distinctly~~ different from virulence architecture in specialist pathogens that often have one or a few large effect genes that control virulence (Keen 1992, De Feyter, Yang et al. 1993, Abramovitch and Martin 2004, Boyd, Ridout et al. 2013, Vleeshouwers and Oliver 2014). Further studies are needed to compare how the host plant species may affect this image of genetic variation in virulence.

These results indicate particular challenges for breeding durable resistance to *B. cinerea*, and possibly other generalist pathogens. The highly polygenic variation in

virulence, combined with genomic sequencing showing that this pathogen is an inter-breeding population, suggests that the pathogen is actively blending a large collection of polymorphic virulence loci (~~Rowe and Kliebenstein 2007, Fekete, Fekete et al. 2012, Atwell, Corwin et al. 2015~~) (Rowe and Kliebenstein 2007, Fekete, Fekete et al. 2012, Atwell, Corwin et al. 2015, Atwell, Corwin et al. 2018). Thus, it is ~~not~~insufficient to breed crop resistance against a single isolate of *B. cinerea*, as this resistance mechanism would likely be rapidly overcome by new genotypes within the field population of *B. cinerea*. In contrast, it is likely necessary to breed resistance using a population of the pathogen, and to focus on plant loci that target entire virulence pathways or mechanisms. The results in this study indicate that the specific genetics of the plant host, the host's general domestication status, and the specific genetics of the pathogen isolate will all combine to affect how the estimated breeding value inferred from any experiment will translate to a field application (Table 1). As such, utilizing a single or even a few pathogen isolates to guide resistance breeding in plants is unlikely to translate to durable resistance against *B. cinerea* as a species. Further, the lack of evidence for a domestication bottleneck on tomato resistance to *B. cinerea* suggests that, at least for tomato, allelic variation in this generalist pathogen is sufficient to overcome introgression of wild resistance genes or alleles into the domesticated crop.

This study examined the contributions of host and pathogen natural genetic variation to the quantitative interaction in the tomato-*B. cinerea* pathosystem. ~~In addition, the study explicitly tested the effects of tomato domestication on this pathosystem.~~ *B. cinerea* has a highly quantitative genetic basis of virulence on tomato, which is dominated by pathogen effects but also sensitive to genetic variation linked to tomato domestication. Future studies are necessary to test if this pattern of domestication responses in tomato is similar to patterns in other crops. Because this population of *B. cinerea* can infect a wide range of hosts, it will be possible to directly conduct this study. By extending future work to additional domestication events, it may be possible to test whether independent crop domestication events have a consistent underlying genetic signal of *B. cinerea* adaptation to plant domestication.

Methods

Tomato genetic resources

We obtained seeds for 12 selected tomato genotypes in consultation with the UC Davis Tomato Genetics Resource Center. These include a diverse sample of 6 genotypes of domesticated tomato's closest wild relative (*S. pimpinellifolium*) sampling across its major geographic regions (Peru, Ecuador) and 6 heritage and modern varieties of *S. lycopersicum*, focusing on mid- to late-20th century improved varieties (Lin, Zhu et al. 2014, Blanca, Montero-Pau et al. 2015). While genetic data is not available for all of our *S. pimpinellifolium* accessions, 9 of the 12 accessions have been genotyped and span the mappable diversity in domesticated tomato and its close relatives (Sim, Durstewitz et al. 2012) (Supplemental Figure 24). We bulked all genotypes in long-day (16h photoperiod) greenhouse conditions at UC Davis in fall 2014. We grew plants under metal-halide lamps using day/night temperatures at 25°C/18°C in 4" pots filled with standard potting soil (Sunshine mix #1, Sun Gro Horticulture). Plants were watered once daily and pruned and staked to maintain upright growth. Fruits were collected at maturity and stored at 4°C in dry paper bags until seed cleaning. To clean the seeds, we incubated seeds and locule contents at 24°C in 1% protease solution (Rapidase C80 Max) for 2h, then rinsed them in deionized water and air-dried. We then stored seeds in a cool, dry, dark location until use.

To grow plants for detached leaf assays, we bleach-sterilized all seeds and germinated them on paper in the growth chamber using flats covered with humidity domes. At 7 days we transferred seedlings to soil (SunGro Horticulture, Agawam, MA) and grew all plants in growth chambers in 20°C, short-day (10h photoperiod) conditions with 180-190 μ M light intensity and 60% RH. We bottom-watered with deionized water every two days for two weeks, and at week 3 watered every two days with added nutrient solution (0.5% N-P-K fertilizer in a 2-1- 2 ratio; Grow More 4-18-38). The plants were used for detached leaf assays 6 weeks after transferring seedlings to soil. Flowering in this system did not occur until minimally 9 weeks of age for any accession, and as such we were sampling midway between the juvenile/adult transition and any flowering time decision. This window has been successful to minimize any major ontogenetic effects on the pathogen/host interaction in other systems (Corwin, Copeland et al. 2016).

***B. cinerea* genetic resources**

We utilized a previously described collection of *B. cinerea* isolates that were isolated as single spores from natural infections of fruit and vegetable tissues collected in California and internationally ~~-(Atwell, Corwin et al. 2015, Zhang, Corwin et al. 2017)~~ (Atwell, Corwin et al. 2015, Zhang, Corwin et al. 2017, Atwell, Corwin et al. 2018). This included five isolates obtained from natural infections of tomato. We maintained *B. cinerea* isolates as conidial suspensions in 30% glycerol for long-term storage at -80°C. For regrowth, we diluted spore solutions to 10% concentration in filter-sterilized 50% grape juice, and then inoculated onto 39g/L potato dextrose agar (PDA) media. We grew isolates at 25°C in 12h light and propagated every 2 weeks. Sequencing failed for 6 out of our 97 phenotyped isolates. For bigRR GWA mapping with the 91 isolates genotyped in this study, we utilized a total of 272,672 SNPs against the *B. cinerea* T4 genome with minor allele frequency (MAF) 0.20 or greater, and less than 10% missing calls across the isolates (SNP calls in at least 82/ 91 isolates) ~~(Atwell, Corwin et al. 2018). For GEMMA mapping, we used 91 isolates with a total of 237,878 SNPs against the *B. cinerea* B05.10 genome with MAF 0.20 or greater and less than 10% missing calls. The overall SNP number was similar when using either reference genome.~~

Detached leaf assay

To study the effect of genetic variation in host and pathogen on lesion formation, we infected detached leaves of 12 diverse tomato varieties with the above 97 *B. cinerea* isolates. We used a randomized complete block design for a total of 6 replicates across 2 experiments. In each experiment, this included a total of 10 plants per genotype randomized in 12 flats in 3 growth chambers. Each growth chamber block corresponded with a replicate of the detached leaf assay, such that growth chamber and replicate shared the same environmental block. At 6 weeks of age, we selected 5 leaves per plant (expanded leaves from second true leaf or younger), and 2 leaflet pairs per leaf. We randomized the order of leaves from each plant, and the leaflets were placed on 1% phytoagar in planting flats, with humidity domes. Our inoculation protocol followed previously described methods (Denby, Kumar et al. 2004, Kliebenstein, Rowe et al.

2005). Spores were collected from mature *B. cinerea* cultures grown on canned peach plates and diluted to 10 spores/ μ L in filter-sterilized 50% organic grape juice. Spores in grape juice were maintained in 4°C refrigeration or on ice from the time of collection, to inhibit germination prior to inoculation. The diluted spore suspensions were homogenized by agitation continuously during the entire process of applying the spores to all samples. This maintains the spores in the suspension and ensures even application across samples, then 4 μ L droplets were placed onto the detached leaflets at room temperature. The entire inoculation took approximately 2 hours of time per experiment. Mock-inoculated control leaves were treated with 4 μ L of 50% organic grape juice without spores. Digital photos were taken of all leaflets at 24, 48, and 72 hours post inoculation and automated image analysis was used to measure lesion size.

Automated Image Analysis

Lesion area was digitally measured using the EBIImage and CRIImage packages (Pau, Fuchs et al. 2010, Failmezger, Yuan et al. 2012) in the R statistical environment (R Development Core Team 2008), as previously described (Corwin, Copeland et al. 2016, Corwin, Subedy et al. 2016). Leaflets were identified as objects with green hue, and lesions were identified as low-saturation objects within leaves. Images masks were generated for both the leaf and lesion, then manually refined by a technician to ensure accurate object calling. The area of these leaves and lesions were then automatically measured as pixels per lesion and converted to area using a 1 cm reference within each image.

Data analysis

We analyzed lesion areas using a general linear models for the full experiment to determine the contributions of plant and pathogen genotype, including the fixed effects of isolate genotype, plant domestication (*S. lycopersicum* or *S. pimpinellifolium*), plant genotype (which is nested within domestication status), experiment, and block (nested within experiment) on lesion area, as well as their interactions ((R lme4 package; (Bates, Maechler et al. 2015)). Two of our 97 isolates that did not have replication across

2 experiments were dropped at this stage of analysis. We used the following linear models throughout our analyses.

Main mixed-effect model of lesion size variation

$$Y = I + D/P + I:D + I:D/P + W_R/L/A + E_R + E_R:I$$

Within-plant accession mixed-effect model of lesion size

$$Y = I + W_R/L/A + E_R + E_R:I$$

Within-isolate mixed-effect model of lesion size

$$Y = D/P + E_R$$

Where I represents fungal genotype (isolate), P represents plant genotype (accession), D represents domestication status, E represents experiment, W represents whole plant, L represents leaf, A represents leaflet position. Factors with the subscript R are included in the analysis as random effects.

~~The significance of individual terms in the model did not change if experiment and block were treated as random effects. Adding terms for individual plant, leaf, and leaflet position did not significantly improve the full model, so they were omitted from further analysis. This~~ The within-plant accession model was used to calculate the significance of each factor and to obtain the least-squared means of lesion size for each *B. cinerea* isolate x tomato accession as well as for each *B. cinerea* isolate x domesticated/wild tomato. We also calculated a domestication sensitivity phenotype, Sensitivity = (Domesticated lesion size – Wild lesion size) / Domesticated lesion size.

We bootstrapped assignment of plant accessions to domestication groups in order to assess the robustness of our observed domestication effects. We randomly drew three genotypes from the domesticated and wild groupings and assigned them to a new pseudo-wild grouping. The other six genotypes were assigned as a pseudo-domesticated grouping and the model was rerun. This bootstrapping was repeated 100 times with each representing a random draw. We used these to repeat the full model and to repeat the individual isolate models, as a test of the robustness of the tomato domestication effect.

Using tomato sequence data from the SolCAP diversity panel that contained 9 of our 12 accessions, we determined pairwise genetic distances between our accessions (Sim, Durstewitz et al. 2012). We calculated pairwise Euclidean distances between 426 wild and domesticated tomato accessions from Infinium SNP genotyping at 7,720 loci

using the R adegenet package (Jombart , Sim, Durstewitz et al. 2012). Clustering is by R hclust (in the stats package) default UPGMA method (R Development Core Team 2008).

We used several methods to examine host specialization to tomato within *B. cinerea*. First, we split our *B. cinerea* population into isolates collected from tomato tissue vs. other hosts. We compared these groups by t-test for virulence on domesticated tomato genotypes, wild tomato genotypes, or all tomato genotypes. Next, we used a Wilcoxon signed-rank test to compare the rank order distribution of [model-adjusted](#) lesion sizes across paired tomato genotypes. ~~To~~ [Also, to](#) examine host specialization to tomato domestication within *B. cinerea*, we used a Wilcoxon signed-rank test to compare the rank order of [model-adjusted](#) lesion sizes across all domesticated vs. all wild tomato genotypes. Finally, we conducted single-isolate ANOVAs with FDR correction [on general linear models](#) to identify isolates with a significant response to plant genotype or domestication status.

The model means and Domestication Sensitivity were used as the phenotypic input for GWA using bigRR, a heteroskedastic ridge regression method that incorporates SNP-specific shrinkage (Shen, Alam et al. 2013). This approach has previously had a high validation rate (Ober, Huang et al. 2015, Corwin, Copeland et al. 2016, Francisco, Joseph et al. 2016, Kooke, Kruijer et al. 2016). The *B. cinerea* bigRR GWA used 272,672 SNPs at MAF 0.20 or greater and <10% missing SNP calls as described above (Atwell, Corwin et al. 2018). Because bigRR provides an estimated effect size, but not a p-value, significance was estimated using 1000 permutations to determine effect significance at 95%, 99%, and [\(approximately\)](#) 99.9% thresholds (Doerge and Churchill 1996, Shen, Alam et al. 2013, Corwin, Copeland et al. 2016). SNPs were annotated by custom R scripts with gene transfer format file construction from the T4 gene models for genomic DNA by linking the SNP to genes within a 2kbp window (<http://www.broadinstitute.org>, (Staats and van Kan 2012)). Functional annotations are based on the T4 gene models for genomic DNA (<http://www.broadinstitute.org>, *B. cinerea*; (Staats and van Kan 2012)). Additional genes of interest, based on a broad literature search of known virulence loci, were taken from

NCBI (<https://www.ncbi.nlm.nih.gov/>) and included by mapping sequence to the T4 reference using MUMmer v3.0 (Kurtz, Phillippy et al. 2004).

To predict expected overlap of significant SNPs across plant genotypes, we used the average number of significant SNPs per each of the 12 plant genotypes (14,000 SNPs) and calculated expected overlap between those 12 lists using binomial coefficients. ~~The *B. cinerea* GEMMA used 237,878 SNPs at MAF 0.20 or greater, and less than 10% missing SNP calls as described above. To determine significance of SNPs by GEMMA, we used 1000 permutations to determine p-value significance at the 99%, and 99.9% thresholds (Doerge and Churchill 1996, Shen, Alam et al. 2013, Corwin, Copeland et al. 2016). SNPs were annotated using a custom R script linking the SNP to genes within a 2kbp window from the gene transfer format file construction from the B05.10 gene models for genomic DNA (Staats and van Kan 2012, Zerbino, Achuthan et al. 2017). A table of gene name translations across genome annotations was pulled from the gene overlap between the bigRR T4 annotation and GEMMA B05.10 annotation using a custom R script and gene name translations pulled from the INRA *Botrytis cinerea* Portal (Choquer, Fournier et al. 2007, Viaud, Adam-Blondon et al. 2012).~~ Functional annotations of the overlap_gene lists are based on the T4 gene models for genomic DNA (<http://www.broadinstitute.org>, *B. cinerea*; (Staats and van Kan 2012).

Supplemental Data Files

Supplemental Data Set 1. Mean \pm SE of *B. cinerea* lesion size of all isolates across all tomato accessions.

Supplemental Data Set 2. Gene and Function Annotation from ~~B05.10 and~~ T4 GWA Results

Supplemental Data Set 3. Results of single-isolate ANOVA on mixed effect model

Supplemental Table 1. Results of ANOVA following removal of domestication-associated isolates

Supplemental Figure 1. Allele frequency spectrum of *B. cinerea* SNPs.

Supplemental Figure 24. Genetic distance between selected tomato accessions.

[Supplemental Figure 3. Correlation between *B. cinerea* lesion size on tomato and on *A. thaliana*](#)

[Supplemental Figure 42. Rank order plot of *B. cinerea* lesion size on two tomato genotypes.](#)

[Supplemental Figure 3. GWA by GEMMA of *B. cinerea* lesion size on individual tomato genotypes.](#)

[Supplemental Figure 4. Frequency of overlap in *B. cinerea* GEMMA GWA significance across tomato accessions.](#)

[Supplemental Figure 5. GEMMA GWA analysis of domestication sensitivity in *B. cinerea*.](#)

Figure Legends

Figure 1. *Botrytis cinerea* x tomato diversity in detached leaf assay and digital image analysis. a) Individual tomato leaflets of 6 *S. lycopersicum* genotypes and 6 *S. pimpinellifolium* genotypes are in randomized rows, spore droplets of individual *B. cinerea* isolates are in randomized columns. Digital images are collected 72 hours post inoculation. Single droplets of 40 *B. cinerea* spores are infected on randomized leaflets using randomized isolates, and digital images are taken 72 hours post inoculation. b) Digital masking of leaf and lesion is followed by automated measurement of area for each lesion. c-h) Variation in lesion size resulting of the interaction of *B. cinerea* and diverse tomato genotypes. c) Average lesion size of single *B. cinerea* isolates (line traces) across tomato host genotypes grouped by domestication status. d) Highlight of the common reference *B. cinerea* isolate B05.10. e) Highlight of the ten highest-virulence isolates, as estimated by mean virulence across all tomato genotypes. f) Highlight of the ten most saprophytic, or low virulence, isolates, as estimated by mean virulence across all genotypes. g) Highlight of the five isolates collected from tomato tissue. h) Highlight of the two isolates with significant domestication sensitivity.

Figure 2. Distribution of tomato genotype susceptibility to infection with 97 genetically diverse *B. cinerea* isolates. Violin plots show the distribution of lesion size caused by *B. cinerea* isolates on each tomato host genotype. Individual points are mean lesion size for each of the 97 different isolate-host pairs. The boxes show the 75th percentile distribution, and the horizontal

line shows the mean resistance of the specific host genotype. The tomato genotypes are grouped based on their status as wild or domesticated germplasm.

Figure 3. Distribution of *B. cinerea* virulence by tomato domestication status.

The violin plots show the mean virulence of each *B. cinerea* isolate on the tomato genotypes, grouped as wild or domesticated germplasm. The domestication effect on lesion size is significant (Table 1, ANOVA, $p < 2e-16 = 0.0006$). The interaction plot between the two violin plots connects the average lesion size of a single *B. cinerea* isolate between the wild and domesticated germplasm.

Figure 4. GWA of *B. cinerea* lesion size on individual tomato genotypes.

Botrytis cinerea chromosomes are differentiated by shading, alternating light and dark grey.

a) Manhattan plot of estimated SNP effect sizes from bigRR for *B. cinerea* lesion size using a single tomato accession, LA2093. Permutation-derived thresholds are shown in horizontal dashed lines.

b) The number of tomato accessions for which a *B. cinerea* SNP was significantly linked to lesion development by bigRR using the 99% permutation threshold. Frequency is number of phenotypes in which the SNP exceeds the threshold. Vertical dotted lines identify regions with overlap between the top 100 large-effect SNPs for LA2093 and significance across the majority (≥ 6) of tomato genotypes tested.

Figure 5. Frequency of overlap in *B. cinerea* GWA significance across tomato accessions.

a) The frequency with which the *B. cinerea* SNPs significantly associate with lesion size on the 12 tomato accessions using bigRR and the 99% permutation threshold. The black line indicates the expected frequency of random overlap, given the number of significant SNPs per plant genotype and size of total SNP set. The inset zooms in on the distribution for overlapping SNPs above 6 plant genotypes for easier visualization. There were no SNPs expected to overlap by random chance in the inset.

b) The frequency with which *B. cinerea* genes significantly associated with lesion size on the 12 tomato accessions. Genes were called as significant if there was one significant SNP called at the 99% permutation threshold within the gene body, or within 2kb of the gene body.

Figure 6. Host specificity of significant SNPs linked to the gene BcT4_6001 (Bcin14g00870).

a) SNPs with effects estimates above the 99% permutation threshold are colored by trait (plant [phenotype-accession](#) in which the effect was estimated). [Wild accessions are oranges \(yellow to red shades\) and domesticated accessions are blues \(green to purple shades\)](#). BcT4_6001 (Bcin14g00870) is a pectinesterase gene linked to at least one significant SNP on all 12 of the tested tomato accessions by bigRR. The annotated exons are depicted as turquoise rectangles, with the start codon marked with an arrow indicating the direction of transcription. Red rectangles indicate corresponding linkage disequilibrium blocks from Figure 6b.

b) Linkage disequilibrium plot, including all pairwise comparisons of SNPs in the 2kb region surrounding Bcin14g00870. The color scheme for each SNP pair is D'/LOD: white if LOD <2 and D' <1, bright red for LOD ≥2 and D'=1, intermediate shades for LOD ≥2 and D' <1.

Figure 7. GWA analysis of domestication sensitivity in *B. cinerea*.

Domestication sensitivity of each isolate was estimated using the average virulence on the wild and domesticated tomato germplasm and using calculated Sensitivity. This was then utilized for GWA mapping by bigRR.

a) The top 1000 SNPs that significantly affect lesion size across domesticated tomato, wild tomato or domestication sensitivity are shown. Significance is called as crossing the 99% permutation threshold.

b) Venn diagram of overlapping SNPs identified as crossing the 99% permutation threshold for each trait.

c) Venn diagram of overlapping genes identified as crossing the 99% permutation threshold for each trait. Genes were called as significant if there was one significant SNP within the gene body or within 2kb of the gene body.

[Acknowledgements](#)

[Financial support for this work was provided by the National Research Foundation DNR grant 99, US NSF grants IOS 1339125, MCB 1330337 and IOS 1021861, and the USDA National Institute of Food and Agriculture, Hatch project number CA-D-PLS-7033-H.](#)

1041 **References**

- 1042 Abramovitch, R. B. and G. B. Martin (2004). "Strategies used by bacterial pathogens to suppress plant
1043 defenses." Current opinion in plant biology **7**(4): 356-364.
- 1044 AbuQamar, S., M.-F. Chai, H. Luo, F. Song and T. Mengiste (2008). "Tomato protein kinase 1b mediates
1045 signaling of plant responses to necrotrophic fungi and insect herbivory." The Plant Cell **20**(7): 1964-1983.
- 1046 Atwell, S., J. Corwin, N. Soltis and D. Kliebenstein (2018). "Resequencing and association mapping of the
1047 generalist pathogen *Botrytis cinerea*." bioRxiv.
- 1048 Atwell, S., J. Corwin, N. Soltis, A. Subedy, K. Denby and D. J. Kliebenstein (2015). "Whole genome
1049 resequencing of *Botrytis cinerea* isolates identifies high levels of standing diversity." Frontiers in
1050 microbiology **6**: 996.
- 1051 Bai, Y. and P. Lindhout (2007). "Domestication and breeding of tomatoes: what have we gained and
1052 what can we gain in the future?" Annals of botany **100**(5): 1085-1094.
- 1053 Ballaré, C. L. and R. Pierik (2017). "The shade-avoidance syndrome: multiple signals and ecological
1054 consequences." Plant, cell & environment **40**(11): 2530-2543.
- 1055 Barrett, L. G. and M. Heil (2012). "Unifying concepts and mechanisms in the specificity of plant–enemy
1056 interactions." Trends in plant science **17**(5): 282-292.
- 1057 Barrett, L. G., J. M. Kniskern, N. Bodenhausen, W. Zhang and J. Bergelson (2009). "Continua of specificity
1058 and virulence in plant host–pathogen interactions: causes and consequences." New Phytologist **183**(3):
1059 513-529.
- 1060 Bartoli, C. and F. Roux (2017). "Genome-Wide Association Studies In Plant Pathosystems: Toward an
1061 Ecological Genomics Approach." Frontiers in plant science **8**.
- 1062 Bates, D., M. Maechler, B. Bolker and S. Walker (2015). "Fitting Linear Mixed-Effects Models Using
1063 lme4." Journal of Statistical Software **67**(1): 1-48.
- 1064 Bergougnoux, V. (2014). "The history of tomato: from domestication to biopharming." Biotechnology
1065 advances **32**(1): 170-189.
- 1066 Bhardwaj, V., S. Meier, L. N. Petersen, R. A. Ingle and L. C. Roden (2011). "Defence responses of
1067 *Arabidopsis thaliana* to infection by *Pseudomonas syringae* are regulated by the circadian clock." PLoS
1068 one **6**(10): e26968.
- 1069 Bittel, P. and S. Robatzek (2007). "Microbe-associated molecular patterns (MAMPs) probe plant
1070 immunity." Current opinion in plant biology **10**(4): 335-341.
- 1071 Blanca, J., J. Montero-Pau, C. Sauvage, G. Bauchet, E. Illa, M. J. Díez, D. Francis, M. Causse, E. van der
1072 Knaap and J. Cañizares (2015). "Genomic variation in tomato, from wild ancestors to contemporary
1073 breeding accessions." BMC genomics **16**(1): 257.
- 1074 Blanco-Ulate, B., A. Morales-Cruz, K. C. Amrine, J. M. Labavitch, A. L. Powell and D. Cantu (2014).
1075 "Genome-wide transcriptional profiling of *Botrytis cinerea* genes targeting plant cell walls during
1076 infections of different hosts." Frontiers in plant science **5**.
- 1077 Boller, T. and S. Y. He (2009). "Innate immunity in plants: an arms race between pattern recognition
1078 receptors in plants and effectors in microbial pathogens." Science **324**(5928): 742-744.
- 1079 Boyd, L. A., C. Ridout, D. M. O'Sullivan, J. E. Leach and H. Leung (2013). "Plant–pathogen interactions:
1080 disease resistance in modern agriculture." Trends in genetics **29**(4): 233-240.
- 1081 Campos, M. L., Y. Yoshida, I. T. Major, D. de Oliveira Ferreira, S. M. Weraduwege, J. E. Froehlich, B. F.
1082 Johnson, D. M. Kramer, G. Jander and T. D. Sharkey (2016). "Rewiring of jasmonate and phytochrome B
1083 signalling uncouples plant growth-defense tradeoffs." Nature communications **7**: 12570.
- 1084 Cervený, L., A. Strasková, V. Danková, A. Hartlová, M. Cecková, F. Staud and J. Stulík (2013).
1085 "Tetratricopeptide repeat motifs in the world of bacterial pathogens: role in virulence mechanisms."
1086 Infection and immunity **81**(3): 629-635.

1087 Chaudhary, B. (2013). "Plant domestication and resistance to herbivory." International journal of plant
1088 genomics **2013**.

1089 Choquer, M., E. Fournier, C. Kunz, C. Levis, J.-M. Pradier, A. Simon and M. Viaud (2007). "Botrytis cinerea
1090 virulence factors: new insights into a necrotrophic and polyphageous pathogen." FEMS microbiology
1091 letters **277**(1): 1-10.

1092 Corwin, J. A., D. Copeland, J. Feusier, A. Subedy, R. Eshbaugh, C. Palmer, J. Maloof and D. J. Kliebenstein
1093 (2016). "The quantitative basis of the Arabidopsis innate immune system to endemic pathogens
1094 depends on pathogen genetics." PLoS Genet **12**(2): e1005789.

1095 Corwin, J. A., A. Subedy, R. Eshbaugh and D. J. Kliebenstein (2016). "Expansive phenotypic landscape of
1096 Botrytis cinerea shows differential contribution of genetic diversity and plasticity." Molecular Plant-
1097 Microbe Interactions **29**(4): 287-298.

1098 Couch, B. C., I. Fudal, M.-H. Lebrun, D. Tharreau, B. Valent, P. Van Kim, J.-L. Nottéghem and L. M. Kohn
1099 (2005). "Origins of host-specific populations of the blast pathogen Magnaporthe oryzae in crop
1100 domestication with subsequent expansion of pandemic clones on rice and weeds of rice." Genetics
1101 **170**(2): 613-630.

1102 Dalmais, B., J. Schumacher, J. Moraga, P. Le Pecheur, B. Tudzynski, I. G. Collado and M. Viaud (2011).
1103 "The Botrytis cinerea phytotoxin botcinic acid requires two polyketide synthases for production and has
1104 a redundant role in virulence with botrydial." Molecular plant pathology **12**(6): 564-579.

1105 Dalman, K., K. Himmelstrand, Å. Olson, M. Lind, M. Brandström-Durling and J. Stenlid (2013). "A
1106 genome-wide association study identifies genomic regions for virulence in the non-model organism
1107 Heterobasidion annosum ss." PLoS One **8**(1): e53525.

1108 Dangl, J. L. and J. D. Jones (2001). "Plant pathogens and integrated defence responses to infection."
1109 nature **411**(6839): 826-833.

1110 De Feyter, R., Y. Yang and D. W. Gabriel (1993). "Gene-for-genes interactions between cotton R genes
1111 and Xanthomonas campestris pv. malvacearum avr genes." Molecular plant-microbe interactions: MPMI
1112 **6**(2): 225-237.

1113 Dean, R., J. A. Van Kan, Z. A. Pretorius, K. E. Hammond-Kosack, A. Di Pietro, P. D. Spanu, J. J. Rudd, M.
1114 Dickman, R. Kahmann and J. Ellis (2012). "The Top 10 fungal pathogens in molecular plant pathology."
1115 Molecular plant pathology **13**(4): 414-430.

1116 Deighton, N., I. Muckenschnabel, A. J. Colmenares, I. G. Collado and B. Williamson (2001). "Botrydial is
1117 produced in plant tissues infected by Botrytis cinerea." Phytochemistry **57**(5): 689-692.

1118 Denby, K. J., P. Kumar and D. J. Kliebenstein (2004). "Identification of Botrytis cinerea susceptibility loci
1119 in Arabidopsis thaliana." The Plant Journal **38**(3): 473-486.

1120 Desjardins, C. A., K. A. Cohen, V. Munsamy, T. Abeel, K. Maharaj, B. J. Walker, T. P. Shea, D. V. Almeida,
1121 A. L. Manson and A. Salazar (2016). "Genomic and functional analyses of Mycobacterium tuberculosis
1122 strains implicate ald in D-cycloserine resistance." Nature genetics **48**(5): 544-551.

1123 Diaz, J., A. ten Have and J. A. van Kan (2002). "The role of ethylene and wound signaling in resistance of
1124 tomato to Botrytis cinerea." Plant physiology **129**(3): 1341-1351.

1125 Dodds, P. N. and J. P. Rathjen (2010). "Plant immunity: towards an integrated view of plant-pathogen
1126 interactions." Nature Reviews Genetics **11**(8): 539-548.

1127 Doebley, J. F., B. S. Gaut and B. D. Smith (2006). "The molecular genetics of crop domestication." Cell
1128 **127**(7): 1309-1321.

1129 Doerge, R. W. and G. A. Churchill (1996). "Permutation tests for multiple loci affecting a quantitative
1130 character." Genetics **142**(1): 285-294.

1131 Dwivedi, S. L., H. D. Upadhyaya, H. T. Stalker, M. W. Blair, D. J. Bertioli, S. Nielsen and R. Ortiz (2008).
1132 "Enhancing crop gene pools with beneficial traits using wild relatives." Plant Breeding Reviews **30**: 179.

1133 Egashira, H., A. Kuwashima, H. Ishiguro, K. Fukushima, T. Kaya and S. Imanishi (2000). "Screening of wild
 1134 accessions resistant to gray mold (*Botrytis cinerea* Pers.) in *Lycopersicon*." Acta physiologiae plantarum
 1135 **22**(3): 324-326.
 1136 Elad, Y., B. Williamson, P. Tudzynski and N. Delen (2007). *Botrytis* spp. and diseases they cause in
 1137 agricultural systems—an introduction. Botrytis: Biology, pathology and control, Springer: 1-8.
 1138 Failmezger, H., Y. Yuan, O. Rueda, F. Markowitz and M. H. Failmezger (2012). "CRImage: CRImage a
 1139 package to classify cells and calculate tumour cellularity." R package version 1.24.0.
 1140 Farhat, M. R., B. J. Shapiro, K. J. Kieser, R. Sultana, K. R. Jacobson, T. C. Victor, R. M. Warren, E. M.
 1141 Streicher, A. Calver and A. Sloutsky (2013). "Genomic analysis identifies targets of convergent positive
 1142 selection in drug-resistant *Mycobacterium tuberculosis*." Nature genetics **45**(10): 1183-1189.
 1143 Fekete, É., E. Fekete, L. Irinyi, L. Karaffa, M. Árnayasi, M. Asadollahi and E. Sándor (2012). "Genetic
 1144 diversity of a *Botrytis cinerea* cryptic species complex in Hungary." Microbiological Research **167**(5): 283-
 1145 291.
 1146 Ferrari, S., R. Galletti, C. Denoux, G. De Lorenzo, F. M. Ausubel and J. Dewdney (2007). "Resistance to
 1147 *Botrytis cinerea* induced in *Arabidopsis* by elicitors is independent of salicylic acid, ethylene, or
 1148 jasmonate signaling but requires PHYTOALEXIN DEFICIENT3." Plant physiology **144**(1): 367-379.
 1149 Ferrari, S., J. M. Plotnikova, G. De Lorenzo and F. M. Ausubel (2003). "Arabidopsis local resistance to
 1150 *Botrytis cinerea* involves salicylic acid and camalexin and requires EDS4 and PAD2, but not SID2, EDS5 or
 1151 PAD4." The Plant Journal **35**(2): 193-205.
 1152 Fillinger, S. and Y. Elad (2015). Botrytis-the Fungus, the Pathogen and Its Management in Agricultural
 1153 Systems, Springer.
 1154 Finkers, R., Y. Bai, P. van den Berg, R. van Berloo, F. Meijer-Dekens, A. Ten Have, J. van Kan, P. Lindhout
 1155 and A. W. van Heusden (2008). "Quantitative resistance to *Botrytis cinerea* from *Solanum neorickii*." Euphytica
 1156 **159**(1-2): 83-92.
 1157 Finkers, R., A. W. van Heusden, F. Meijer-Dekens, J. A. van Kan, P. Maris and P. Lindhout (2007). "The
 1158 construction of a *Solanum habrochaites* LYC4 introgression line population and the identification of QTLs
 1159 for resistance to *Botrytis cinerea*." Theoretical and Applied Genetics **114**(6): 1071-1080.
 1160 Fordyce, R., N. Soltis, C. Caseys, G. Gwinner, J. Corwin, S. Atwell, D. Copeland, J. Feusier, A. Subedy, R.
 1161 Eshbaugh and D. Kliebenstein (2018). "Combining Digital Imaging and GWA Mapping to Dissect Visual
 1162 Traits in Plant/Pathogen Interactions." Plant Physiology.
 1163 Francisco, M., B. Joseph, H. Caligagan, B. Li, J. A. Corwin, C. Lin, R. E. Kerwin, M. Burow and D. J.
 1164 Kliebenstein (2016). "Genome wide association mapping in *Arabidopsis thaliana* identifies novel genes
 1165 involved in linking allyl glucosinolate to altered biomass and defense." Frontiers in plant science **7**.
 1166 Gao, Y., Z. Liu, J. D. Faris, J. Richards, R. S. Brueggeman, X. Li, R. P. Oliver, B. A. McDonald and T. L.
 1167 Friesen (2016). "Validation of genome-wide association studies as a tool to identify virulence factors in
 1168 *Parastagonospora nodorum*." Phytopathology **106**(10): 1177-1185.
 1169 Giraud, T., D. Fortini, C. Levis, C. Lamarque, P. Leroux, K. LoBuglio and Y. Brygoo (1999). "Two sibling
 1170 species of the *Botrytis cinerea* complex, *transposa* and *vacuata*, are found in sympatry on numerous
 1171 host plants." Phytopathology **89**(10): 967-973.
 1172 Glazebrook, J. (2005). "Contrasting mechanisms of defense against biotrophic and necrotrophic
 1173 pathogens." Annu. Rev. Phytopathol. **43**: 205-227.
 1174 Goss, E. M. and J. Bergelson (2006). "Variation in resistance and virulence in the interaction between
 1175 *Arabidopsis thaliana* and a bacterial pathogen." Evolution **60**(8): 1562-1573.
 1176 Guimaraes, R. L., R. T. Chetelat and H. U. Stotz (2004). "Resistance to *Botrytis cinerea* in *Solanum*
 1177 *lycopersicoides* is dominant in hybrids with tomato, and involves induced hyphal death." European
 1178 journal of plant pathology **110**(1): 13-23.

1179 Hacquard, S., B. Kracher, T. Maekawa, S. Vernaldi, P. Schulze-Lefert and E. V. L. van Themaat (2013).
 1180 "Mosaic genome structure of the barley powdery mildew pathogen and conservation of transcriptional
 1181 programs in divergent hosts." Proceedings of the National Academy of Sciences **110**(24): E2219-E2228.
 1182 Hahn, M. (2014). "The rising threat of fungicide resistance in plant pathogenic fungi: Botrytis as a case
 1183 study." Journal of chemical biology **7**(4): 133-141.
 1184 Hevia, M. A., P. Canessa, H. Müller-Esparza and L. F. Larrondo (2015). "A circadian oscillator in the
 1185 fungus Botrytis cinerea regulates virulence when infecting Arabidopsis thaliana." Proceedings of the
 1186 National Academy of Sciences **112**(28): 8744-8749.
 1187 Hyten, D. L., Q. Song, Y. Zhu, I.-Y. Choi, R. L. Nelson, J. M. Costa, J. E. Specht, R. C. Shoemaker and P. B.
 1188 Cregan (2006). "Impacts of genetic bottlenecks on soybean genome diversity." Proceedings of the
 1189 National Academy of Sciences **103**(45): 16666-16671.
 1190 Izquierdo-Bueno, I., V. E. González-Rodríguez, A. Simon, B. Dalmais, J. M. Pradier, P. Le Pêcheur, A.
 1191 Mercier, A. S. Walker, C. Garrido and I. G. Collado (2018). "Biosynthesis of abscisic acid in fungi:
 1192 Identification of a sesquiterpene cyclase as the key enzyme in Botrytis cinerea." Environmental
 1193 microbiology.
 1194 Jombart, T. (2008). "adeqnet: a R package for the multivariate analysis of genetic markers."
 1195 Bioinformatics **24**(11): 1403-1405.
 1196 Jones, J. D. and J. L. Dangi (2006). "The plant immune system." Nature **444**(7117): 323-329.
 1197 Katan, T. (1999). "Current status of vegetative compatibility groups in Fusarium oxysporum."
 1198 Phytoparasitica **27**(1): 51-64.
 1199 Keen, N. (1992). "The molecular biology of disease resistance." Plant molecular biology **19**(1): 109-122.
 1200 Kliebenstein, D. J., H. C. Rowe and K. J. Denby (2005). "Secondary metabolites influence
 1201 Arabidopsis/Botrytis interactions: variation in host production and pathogen sensitivity." The Plant
 1202 Journal **44**(1): 25-36.
 1203 Koenig, D., J. M. Jiménez-Gómez, S. Kimura, D. Fulop, D. H. Chitwood, L. R. Headland, R. Kumar, M. F.
 1204 Covington, U. K. Devisetty and A. V. Tat (2013). "Comparative transcriptomics reveals patterns of
 1205 selection in domesticated and wild tomato." Proceedings of the National Academy of Sciences **110**(28):
 1206 E2655-E2662.
 1207 Kooke, R., W. Kruijer, R. Bours, F. F. Becker, A. Kuhn, J. Buntjer, T. Doeswijk, J. Guerra, H. J. Bouwmeester
 1208 and D. Vreugdenhil (2016). "Genome-wide association mapping and genomic prediction elucidate the
 1209 genetic architecture of morphological traits in Arabidopsis thaliana." Plant Physiology: pp. 00997.02015.
 1210 Kover, P. X. and B. A. Schaal (2002). "Genetic variation for disease resistance and tolerance among
 1211 Arabidopsis thaliana accessions." Proceedings of the National Academy of Sciences **99**(17): 11270-
 1212 11274.
 1213 Kretschmer, M. and M. Hahn (2008). "Fungicide resistance and genetic diversity of Botrytis cinerea
 1214 isolates from a vineyard in Germany." Journal of Plant Diseases and Protection: 214-219.
 1215 Kurtz, S., A. Phillippy, A. L. Delcher, M. Smoot, M. Shumway, C. Antonescu and S. L. Salzberg (2004).
 1216 "Versatile and open software for comparing large genomes." Genome biology **5**(2): R12.
 1217 Lin, T., G. Zhu, J. Zhang, X. Xu, Q. Yu, Z. Zheng, Z. Zhang, Y. Lun, S. Li and X. Wang (2014). "Genomic
 1218 analyses provide insights into the history of tomato breeding." Nature genetics **46**(11): 1220.
 1219 Liu, B., Y.-B. Hong, Y.-F. Zhang, X.-H. Li, L. Huang, H.-J. Zhang, D.-Y. Li and F.-M. Song (2014). "Tomato
 1220 WRKY transcriptional factor SIDRW1 is required for disease resistance against Botrytis cinerea and
 1221 tolerance to oxidative stress." Plant Science **227**: 145-156.
 1222 Lo Presti, L., C. López Díaz, D. Turrà, A. Di Pietro, M. Hampel, K. Heimel and R. Kahmann (2016). "A
 1223 conserved co-chaperone is required for virulence in fungal plant pathogens." New Phytologist **209**(3):
 1224 1135-1148.
 1225 Loxdale, H. D., G. Lushai and J. A. Harvey (2011). "The evolutionary improbability of 'generalism' in
 1226 nature, with special reference to insects." Biological Journal of the Linnean Society **103**(1): 1-18.

1227 Ma, Z. and T. J. Michailides (2005). "Genetic structure of *Botrytis cinerea* populations from different host
1228 plants in California." Plant disease **89**(10): 1083-1089.

1229 Martinez, F., D. Blancard, P. Lecomte, C. Levis, B. Dubos and M. Fermaud (2003). "Phenotypic differences
1230 between vacuina and transposa subpopulations of *Botrytis cinerea*." European Journal of Plant
1231 Pathology **109**(5): 479-488.

1232 Miller, J. and S. Tanksley (1990). "RFLP analysis of phylogenetic relationships and genetic variation in the
1233 genus *Lycopersicon*." TAG Theoretical and Applied Genetics **80**(4): 437-448.

1234 Müller, N. A., C. L. Wijnen, A. Srinivasan, M. Ryngajllo, I. Ofner, T. Lin, A. Ranjan, D. West, J. N. Maloof
1235 and N. R. Sinha (2016). "Domestication selected for deceleration of the circadian clock in cultivated
1236 tomato." Nature genetics **48**(1): 89-93.

1237 Nicot, P., A. Moretti, C. Romiti, M. Bardin, C. Caranta and H. Ferriere (2002). "Differences in
1238 susceptibility of pruning wounds and leaves to infection by *Botrytis cinerea* among wild tomato
1239 accessions." TGC Report **52**: 24-26.

1240 Nicot, P. C. and A. Baille (1996). Integrated control of *Botrytis cinerea* on greenhouse tomatoes. Aerial
1241 Plant Surface Microbiology, Springer: 169-189.

1242 Nomura, K., M. Melotto and S.-Y. He (2005). "Suppression of host defense in compatible plant–
1243 *Pseudomonas syringae* interactions." Current opinion in plant biology **8**(4): 361-368.

1244 Ober, U., W. Huang, M. Magwire, M. Schlather, H. Simianer and T. F. Mackay (2015). "Accounting for
1245 genetic architecture improves sequence based genomic prediction for a *Drosophila* fitness trait." PLoS
1246 One **10**(5): e0126880.

1247 Ormond, E. L., A. P. Thomas, P. J. Pugh, J. K. Pell and H. E. Roy (2010). "A fungal pathogen in time and
1248 space: the population dynamics of *Beauveria bassiana* in a conifer forest." FEMS microbiology ecology
1249 **74**(1): 146-154.

1250 Panthee, D. R. and F. Chen (2010). "Genomics of fungal disease resistance in tomato." Current genomics
1251 **11**(1): 30-39.

1252 Parlevliet, J. E. (2002). "Durability of resistance against fungal, bacterial and viral pathogens; present
1253 situation." Euphytica **124**(2): 147-156.

1254 Pau, G., F. Fuchs, O. Sklyar, M. Boutros and W. Huber (2010). "EBImage—an R package for image
1255 processing with applications to cellular phenotypes." Bioinformatics **26**(7): 979-981.

1256 Pedras, M. S. C. and P. W. Ahiaonu (2005). "Metabolism and detoxification of phytoalexins and analogs
1257 by phytopathogenic fungi." Phytochemistry **66**(4): 391-411.

1258 Pedras, M. S. C., S. Hossain and R. B. Snitynsky (2011). "Detoxification of cruciferous phytoalexins in
1259 *Botrytis cinerea*: Spontaneous dimerization of a camalexin metabolite." Phytochemistry **72**(2): 199-206.

1260 Peralta, I., D. Spooner and S. Knapp (2008). "The taxonomy of tomatoes: a revision of wild tomatoes
1261 (*Solanum* section *Lycopersicon*) and their outgroup relatives in sections *Juglandifolium* and
1262 *Lycopersicoides*." Syst Bot Monogr **84**: 1-186.

1263 Persoons, A., E. Morin, C. Delaruelle, T. Payen, F. Halkett, P. Frey, S. De Mita and S. Duplessis (2014).
1264 "Patterns of genomic variation in the poplar rust fungus *Melampsora larici-populina* identify
1265 pathogenesis-related factors." Frontiers in plant science **5**.

1266 Pieterse, C. M., D. Van der Does, C. Zamioudis, A. Leon-Reyes and S. C. Van Wees (2012). "Hormonal
1267 modulation of plant immunity." Annual review of cell and developmental biology **28**: 489-521.

1268 Poland, J. A., P. J. Balint-Kurti, R. J. Wisser, R. C. Pratt and R. J. Nelson (2009). "Shades of gray: the world
1269 of quantitative disease resistance." Trends in plant science **14**(1): 21-29.

1270 Power, R. A., J. Parkhill and T. de Oliveira (2017). "Microbial genome-wide association studies: lessons
1271 from human GWAS." Nature Reviews Genetics **18**(1): 41-50.

1272 Quidde, T., P. Büttner and P. Tudzynski (1999). "Evidence for three different specific saponin-detoxifying
1273 activities in *Botrytis cinerea* and cloning and functional analysis of a gene coding for a putative
1274 avenacinase." European Journal of Plant Pathology **105**(3): 273-283.

1275 Quidde, T., A. Osbourn and P. Tudzynski (1998). "Detoxification of α -tomatine by *Botrytis cinerea*."
 1276 Physiological and Molecular Plant Pathology **52**(3): 151-165.
 1277 R Development Core Team (2008). "R: A language and environment for statistical computing." R
 1278 Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0.
 1279 Romanazzi, G. and S. Droby (2016). Control Strategies for Postharvest Grey Mould on Fruit Crops.
 1280 Botrytis—the Fungus, the Pathogen and its Management in Agricultural Systems, Springer: 217-228.
 1281 Rosenthal, J. P. and R. Dirzo (1997). "Effects of life history, domestication and agronomic selection on
 1282 plant defence against insects: evidence from maizes and wild relatives." Evolutionary Ecology **11**(3): 337-
 1283 355.
 1284 Rowe, H. C. and D. J. Kliebenstein (2007). "Elevated genetic variation within virulence-associated *Botrytis*
 1285 *cinerea* polygalacturonase loci." Molecular Plant-Microbe Interactions **20**(9): 1126-1137.
 1286 Rowe, H. C. and D. J. Kliebenstein (2008). "Complex genetics control natural variation in *Arabidopsis*
 1287 *thaliana* resistance to *Botrytis cinerea*." Genetics **180**(4): 2237-2250.
 1288 Rowe, H. C., J. W. Walley, J. Corwin, E. K.-F. Chan, K. Dehesh and D. J. Kliebenstein (2010). "Deficiencies
 1289 in jasmonate-mediated plant defense reveal quantitative variation in *Botrytis cinerea* pathogenesis."
 1290 PLoS Pathog **6**(4): e1000861.
 1291 Samuel, S., T. Veloukas, A. Papavasileiou and G. S. Karaoglanidis (2012). "Differences in frequency of
 1292 transposable elements presence in *Botrytis cinerea* populations from several hosts in Greece." Plant
 1293 disease **96**(9): 1286-1290.
 1294 Sauerbrunn, N. and N. L. Schlaich (2004). "PCC1: a merging point for pathogen defence and circadian
 1295 signalling in *Arabidopsis*." Planta **218**(4): 552-561.
 1296 Schumacher, J., J.-M. Pradier, A. Simon, S. Traeger, J. Moraga, I. G. Collado, M. Viaud and B. Tudzynski
 1297 (2012). "Natural variation in the VELVET gene *bcvel1* affects virulence and light-dependent
 1298 differentiation in *Botrytis cinerea*." PLoS One **7**(10): e47840.
 1299 Shen, X., M. Alam, F. Fikse and L. Rönnegård (2013). "A novel generalized ridge regression method for
 1300 quantitative genetics." Genetics **193**(4): 1255-1268.
 1301 Siewers, V., M. Viaud, D. Jimenez-Teja, I. G. Collado, C. S. Gronover, J.-M. Pradier, B. Tudzynski and P.
 1302 Tudzynski (2005). "Functional analysis of the cytochrome P450 monooxygenase gene *bcbot1* of *Botrytis*
 1303 *cinerea* indicates that botrydial is a strain-specific virulence factor." Molecular plant-microbe
 1304 interactions **18**(6): 602-612.
 1305 Sim, S.-C., G. Durstewitz, J. Plieske, R. Wieseke, M. W. Ganai, A. Van Deynze, J. P. Hamilton, C. R. Buell,
 1306 M. Causse and S. Wijeratne (2012). "Development of a large SNP genotyping array and generation of
 1307 high-density genetic maps in tomato." PloS one **7**(7): e40563.
 1308 Smale, M. (1996). "Understanding global trends in the use of wheat diversity and international flows of
 1309 wheat genetic resources."
 1310 Staats, M. and J. A. van Kan (2012). "Genome update of *Botrytis cinerea* strains B05. 10 and T4."
 1311 Eukaryotic cell **11**(11): 1413-1414.
 1312 Stefanato, F. L., E. Abou-Mansour, A. Buchala, M. Kretschmer, A. Mosbach, M. Hahn, C. G. Bochet, J. P.
 1313 Métraux and H. j. Schoonbeek (2009). "The ABC transporter *BcatrB* from *Botrytis cinerea* exports
 1314 camalexin and is a virulence factor on *Arabidopsis thaliana*." The Plant Journal **58**(3): 499-510.
 1315 Stukenbrock, E. H. and B. A. McDonald (2008). "The origins of plant pathogens in agro-ecosystems."
 1316 Annu. Rev. Phytopathol. **46**: 75-100.
 1317 Talas, F., R. Kalih, T. Miedaner and B. A. McDonald (2016). "Genome-wide association study identifies
 1318 novel candidate genes for aggressiveness, deoxynivalenol production, and azole sensitivity in natural
 1319 field populations of *Fusarium graminearum*." Molecular Plant-Microbe Interactions **29**(5): 417-430.
 1320 Tanksley, S. D. (2004). "The genetic, developmental, and molecular bases of fruit size and shape
 1321 variation in tomato." The plant cell **16**(suppl 1): S181-S189.

Tanksley, S. D. and S. R. McCouch (1997). "Seed banks and molecular maps: unlocking genetic potential from the wild." Science **277**(5329): 1063-1066.

ten Have, A., W. Mulder, J. Visser and J. A. van Kan (1998). "The endopolygalacturonase gene Bcpg1 is required for full virulence of *Botrytis cinerea*." Molecular Plant-Microbe Interactions **11**(10): 1009-1016.

Ten Have, A., R. van Berloo, P. Lindhout and J. A. van Kan (2007). "Partial stem and leaf resistance against the fungal pathogen *Botrytis cinerea* in wild relatives of tomato." European journal of plant pathology **117**(2): 153-166.

Tiffin, P. and D. A. Moeller (2006). "Molecular evolution of plant immune system genes." Trends in genetics **22**(12): 662-670.

Upadhyaya, N. M., D. P. Garnica, H. Karaoglu, J. Sperschneider, A. Nemri, B. Xu, R. Mago, C. A. Cuomo, J. P. Rathjen and R. F. Park (2014). "Comparative genomics of Australian isolates of the wheat stem rust pathogen *Puccinia graminis* f. sp. *tritici* reveals extensive polymorphism in candidate effector genes." Frontiers in plant science **5**.

Valette-Collet, O., A. Cimerman, P. Reignault, C. Levis and M. Boccara (2003). "Disruption of *Botrytis cinerea* pectin methylesterase gene Bcpme1 reduces virulence on several host plants." Molecular Plant-Microbe Interactions **16**(4): 360-367.

Viaud, M., A.-F. Adam-Blondon, J. Amselem, P. Bally, A. Cimerman, B. Dalmais-Lenaers, N. Lapalu, M.-H. Lebrun, B. Poinssot and J. M. Pradier (2012). "Le génome de *Botrytis* décrypté." Revue des oenologues et des techniques vitivinicoles et oenologiques(142): 9-11.

Vleeshouwers, V. G. and R. P. Oliver (2014). "Effectors as tools in disease resistance breeding against biotrophic, hemibiotrophic, and necrotrophic plant pathogens." Molecular plant-microbe interactions **27**(3): 196-206.

Weyman, P. D., Z. Pan, Q. Feng, D. G. Gilchrist and R. M. Bostock (2006). "A circadian rhythm-regulated tomato gene is induced by arachidonic acid and *Phytophthora infestans* infection." Plant physiology **140**(1): 235-248.

Wicker, T., S. Oberhaensli, F. Parlange, J. P. Buchmann, M. Shatalina, S. Roffler, R. Ben-David, J. Doležal, H. Šimková and P. Schulze-Lefert (2013). "The wheat powdery mildew genome shows the unique evolution of an obligate biotroph." Nature Genetics **45**(9): 1092-1096.

Wu, J. Q., S. Sakthikumar, C. Dong, P. Zhang, C. A. Cuomo and R. F. Park (2017). "Comparative genomics integrated with association analysis identifies candidate effector genes corresponding to Lr20 in phenotype-paired *Puccinia triticina* isolates from Australia." Frontiers in plant science **8**.

Zerbino, D. R., P. Achuthan, W. Akanni, M. R. Amode, D. Barrell, J. Bhai, K. Billis, C. Cummins, A. Gall and C. G. Girón (2017). "Ensembl 2018." Nucleic acids research **46**(D1): D754-D761.

Zhang, L., A. Khan, D. Nino-Liu and M. Foolad (2002). "A molecular linkage map of tomato displaying chromosomal locations of resistance gene analogs based on a *Lycopersicon esculentum* × *Lycopersicon hirsutum* cross." Genome **45**(1): 133-146.

Zhang, W., J. A. Corwin, D. Copeland, J. Feusier, R. Eshbaugh, F. Chen, S. Atwell and D. J. Kliebenstein (2017). "Plastic transcriptomes stabilize immunity to pathogen diversity: the jasmonic acid and salicylic acid networks within the *Arabidopsis*/*Botrytis* pathosystem." The Plant Cell: tpc. 00348.02017.

Zhou, X. and M. Stephens (2012). "Genome-wide efficient mixed-model analysis for association studies." Nature genetics **44**(7): 821.

Zipfel, C., S. Robatzek, L. Navarro and E. J. Oakeley (2004). "Bacterial disease resistance in *Arabidopsis* through flagellin perception." Nature **428**(6984): 764.

Züst, T. and A. A. Agrawal (2017). "Trade-offs between plant growth and defense against insect herbivory: an emerging mechanistic synthesis." Annual review of plant biology **68**: 513-534.