Learning strategic behavior in social and economic networks

Nicolò Pagan ¹ Florian Dörfler ¹

¹ Automatic Control Laboratory, ETH Zürich, Switzerland

Complex Networks 2018

 7^{th} International Conference on Complex Networks and Their Applications

December, 11-13 2018, Cambridge, UK





Observations

Actors decide with whom they want to interact.



- Actors decide with whom they want to interact.
- Network positions provide benefits to the actors.



- ► Actors decide with whom they want to interact.
- Network positions provide benefits to the actors.





- Actors decide with whom they want to interact.
- Network positions provide benefits to the actors.



- Actors decide with whom they want to interact.
- Network positions provide benefits to the actors.





Social Influence



The more people we are connected to, the more we can influence them.

Social Influence



The more people we are connected to, the more we can influence them.

Social Support



The more our friends' friends are our friends, the safer we feel.

[Heider (1946); Coleman (1990)] (Structural Balance theory)

Social Influence



The more people we are connected to, the more we can influence them.

Social Support



The more our friends' friends are our friends, the safer we feel.

[Heider (1946); Coleman (1990)] (Structural Balance theory)

Brokerage



The more we are on the path between people, the more we can control.

[Burt (1992)]

(Structural Holes theory)

Social Influence



The more people we are connected to, the more we can influence them.

Social Support



The more our friends' friends are our friends, the safer we feel.

[Heider (1946); Coleman (1990)] (Structural Balance theory)

Brokerage



The more we are on the path between people, the more we can control. [Burt (1992)] (Structural Holes theory)

Betweenness Centrality

ETH zürich

Degree Centrality

Clustering Coefficient

Ingredients

▶ **Directed weighted** network \mathcal{G} with $\mathcal{N} = \{1, \dots, N\}$ agents.



Ingredients

- ▶ **Directed weighted** network \mathcal{G} with $\mathcal{N} = \{1, ..., N\}$ agents.
- ▶ $a_{ij} \in [0,1]$ quantifies the importance of the friendship among i and j from i's point of view.



Ingredients

- ▶ **Directed weighted** network \mathcal{G} with $\mathcal{N} = \{1, ..., N\}$ agents.
- ▶ $a_{ij} \in [0,1]$ quantifies the importance of the friendship among i and j from i's point of view.
- A typical action of agent i is:

$$a_i = [a_{i1}, \dots, a_{i,i-1}, a_{i,i+1}, \dots, a_{iN}] \in \mathcal{A} = [0,1]^{N-1},$$



Ingredients

- ▶ **Directed weighted** network \mathcal{G} with $\mathcal{N} = \{1, ..., N\}$ agents.
- ▶ $a_{ij} \in [0,1]$ quantifies the importance of the friendship among i and j from i's point of view.
- ▶ A typical action of agent i is:

$$a_i = [a_{i1}, \dots, a_{i,i-1}, a_{i,i+1}, \dots, a_{iN}] \in \mathcal{A} = [0,1]^{N-1},$$

▶ **Rational** agents: every agent *i* is endowed with a payoff function *V_i* and is looking for

$$a_i^\star \in \arg\max_{a_i \in \mathcal{A}} V_i(a_i, \mathbf{a_{-i}})$$




$$V_i(a_i, \mathbf{a_{-i}}) = t_i(a_i, \mathbf{a_{-i}}) + u_i(a_i, \mathbf{a_{-i}}) - c_i(a_i),$$



$$V_i(a_i, \mathbf{a_{-i}}) = t_i(a_i, \mathbf{a_{-i}}) + u_i(a_i, \mathbf{a_{-i}}) - c_i(a_i),$$

■ Social influence on friends,

$$t_i(a_i, \mathbf{a}_{-\mathbf{i}}) = \sum_{j \neq i} a_{ji}$$



$$V_i(a_i, \mathbf{a_{-i}}) = t_i(a_i, \mathbf{a_{-i}}) + u_i(a_i, \mathbf{a_{-i}}) - c_i(a_i),$$

■ Social influence on friends, on friends of friends, with $\delta_i \in [0,1]$:

$$t_i(a_i, \mathbf{a_{-i}}) = \sum_{j \neq i} a_{ji} + \delta_i \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji}$$
paths of length 2




$$V_i(a_i, \mathbf{a_{-i}}) = t_i(a_i, \mathbf{a_{-i}}) + u_i(a_i, \mathbf{a_{-i}}) - c_i(a_i),$$

Social influence on friends, on friends of friends, ... with $\delta_i \in [0,1]$:

$$\begin{split} t_i(a_i, \mathbf{a_{-i}}) &= \sum_{j \neq i} a_{ji} + \delta_i \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji} + \\ & \underbrace{\text{paths of length 2}}_{\text{paths of length 3}} + \delta_i^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}, \end{split}$$

[Jackson and Wolinsky (1996)]



$$V_i(a_i, \mathbf{a_{-i}}) = t_i(a_i, \mathbf{a_{-i}}) + u_i(a_i, \mathbf{a_{-i}}) - c_i(a_i),$$

■ Social influence on friends, on friends of friends, ... with $\delta_i \in [0,1]$:

$$\begin{split} t_i(a_i, \mathbf{a_{-i}}) &= \sum_{j \neq i} a_{ji} + \delta_i \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji} + \\ & \text{paths of length 2} \\ &+ \delta_i^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}, \\ & \text{paths of length 3} \end{split}$$

[Jackson and Wolinsky (1996)]

■ Clustering coefficient: number of closed triads: $u_i(a_i, \mathbf{a_{-i}}) = \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i,j} a_{ik} a_{kj} \right)$,

[Burger and Buskens (2009)]



$$V_i(a_i, \mathbf{a_{-i}}) = t_i(a_i, \mathbf{a_{-i}}) + u_i(a_i, \mathbf{a_{-i}}) - c_i(a_i),$$

■ Social influence on friends, on friends of friends, ... with $\delta_i \in [0,1]$:

$$\begin{split} t_i(a_i, \mathbf{a_{-i}}) = & \sum_{j \neq i} a_{ji} + \delta_i \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji} + \\ & \text{paths of length 2} \\ + & \delta_i^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}, \\ & \text{paths of length 3} \end{split}$$

[Jackson and Wolinsky (1996)]

■ Clustering coefficient: number of closed triads: $u_i(a_i, \mathbf{a_{-i}}) = \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i,j} a_{ik} a_{kj} \right)$,

[Burger and Buskens (2009)]

Cost of maintaining ties: $c_i(a_i) = \sum_{j \neq i} a_{ij}$.



$$\begin{aligned} V_i(a_i, \mathbf{a_{-i}} | P_i) &= \alpha_i t_i(a_i, \mathbf{a_{-i}}) + \beta_i u_i(a_i, \mathbf{a_{-i}}) - \gamma_i c_i(a_i), & \alpha_i \geq 0, \, \beta_i \in \mathbb{R}, \, \gamma_i > 0 \\ P_i &= \{\alpha_i, \beta_i, \gamma_i, \delta_i\} \end{aligned}$$

■ Social influence on friends, on friends of friends, ... with $\delta_i \in [0,1]$:

$$\begin{split} t_i(a_i, \mathbf{a_{-i}}) &= \sum_{j \neq i} a_{ji} + \delta_i \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji} + \\ & \text{paths of length 2} \\ &+ \delta_i^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}, \\ & \text{paths of length 3} \end{split}$$

[Jackson and Wolinsky (1996)]

■ Clustering coefficient: number of closed triads: $u_i(a_i, \mathbf{a_{-i}}) = \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i,j} a_{ik} a_{kj} \right)$,

[Burger and Buskens (2009)]

Cost of maintaining ties: $c_i(a_i) = \sum_{j \neq i} a_{ij}$.






Definition (Nash equilibrium).

$$V_i(\mathbf{a_i}, \mathbf{a_{-i}}^{\star}|P_i) \leq V_i(\mathbf{a_i}^{\star}, \mathbf{a_{-i}}^{\star}|P_i), \forall \mathbf{a_i} \in \mathcal{A}.$$


Definition (Nash equilibrium).

$$V_i(\mathbf{a_i}, \mathbf{a_{-i}}^{\star}|P_i) \leq V_i(\mathbf{a_i}^{\star}, \mathbf{a_{-i}}^{\star}|P_i), \forall \mathbf{a_i} \in \mathcal{A}.$$




Definition (Nash equilibrium).

$$V_i(\mathbf{a_i}, \mathbf{a_{-i}}^{\star}|P_i) \leq V_i(\mathbf{a_i}^{\star}, \mathbf{a_{-i}}^{\star}|P_i), \forall \mathbf{a_i} \in \mathcal{A}.$$



Definition (Nash equilibrium).


$$V_i(\mathbf{a_i}, \mathbf{a_{-i}}^{\star}|P_i) \leq V_i(\mathbf{a_i}^{\star}, \mathbf{a_{-i}}^{\star}|P_i), \forall \mathbf{a_i} \in \mathcal{A}.$$



Definition (Nash equilibrium).

The network G^* is a NE if for all agents i,

$$V_i(a_i, \mathbf{a_{-i}}^{\star}|P_i) \leq V_i(a_i^{\star}, \mathbf{a_{-i}}^{\star}|P_i), \forall a_i \in \mathcal{A}.$$



Question: For which individual parameters P_i is \mathcal{G}^* in equilibrium?

Homogeneous agents

Assumption

Individual preferences $P_i = P$, for all agents i.


Homogeneous agents

Assumption

Individual preferences $P_i = P$, for all agents i.

Result

Analytic characterization of individual behavior, based on some specific network motifs.




Homogeneous agents


Assumption

Individual preferences $P_i = P$, for all agents i.

Result

Analytic characterization of individual behavior, based on some specific network motifs.





Bala and Goyal (2000), Buechel and Buskens (2013), Buechel and Buskens (2013), Buskens and van de Rijt (2008).

Let the unkown heterogeneous individual preferences set $P_i = \{\alpha_i, \beta_i, \gamma_i, \delta_i\}$. Define:

$$\Phi_i(a_i,P_i) = V_i\left(a_i,a_{-i}^{\star}|P_i\right) - V_i\left(a_i^{\star},a_{-i}^{\star}|P_i\right).$$

Let the unkown heterogeneous individual preferences set $P_i = \{\alpha_i, \beta_i, \gamma_i, \delta_i\}$. Define:

$$\Phi_i(a_i, P_i) = V_i\left(a_i, a_{-i}^{\star}|P_i\right) - V_i\left(a_i^{\star}, a_{-i}^{\star}|P_i\right).$$

Note: $\Phi_i(a_i, P_i) > 0$ if P_i violates the Nash equilibrium condition.

Let the unkown heterogeneous individual preferences set $P_i = \{\alpha_i, \beta_i, \gamma_i, \delta_i\}$. Define:

$$\Phi_i(a_i,P_i) = V_i\left(a_i,a_{-i}^{\star}|P_i\right) - V_i\left(a_i^{\star},a_{-i}^{\star}|P_i\right).$$

Note: $\Phi_i(a_i, P_i) > 0$ if P_i violates the Nash equilibrium condition.

Integrate over the all action space, taking into account only the violations.

$$\Psi_i(P_i) = \int_{\mathcal{A}} \max \left\{ 0, \Phi_i(a_i, P_i) \right\} da_i.$$

Let the unkown heterogeneous individual preferences set $P_i = \{\alpha_i, \beta_i, \gamma_i, \delta_i\}$. Define:

$$\Phi_i(a_i,P_i) = V_i\left(a_i,a_{-i}^{\star}|P_i\right) - V_i\left(a_i^{\star},a_{-i}^{\star}|P_i\right).$$

Note: $\Phi_i(a_i, P_i) > 0$ if P_i violates the Nash equilibrium condition.

Integrate over the all action space, taking into account only the violations.

$$\Psi_i(P_i) = \int_{\mathcal{A}} \max \left\{ 0, \Phi_i(a_i, P_i) \right\} da_i.$$

Minimize the violations

$$\hat{P}_i \in \arg\min_{P_i \in \mathcal{P}} \{\Psi_i(P_i)\}$$



Let the unkown heterogeneous individual preferences set $P_i = \{\alpha_i, \beta_i, \gamma_i, \delta_i\}$. Define:

$$\Phi_i(a_i,P_i) = V_i\left(a_i,a_{-i}^{\star}|P_i\right) - V_i\left(a_i^{\star},a_{-i}^{\star}|P_i\right).$$

Note: $\Phi_i(a_i, P_i) > 0$ if P_i violates the Nash equilibrium condition.

Integrate over the all action space, taking into account only the violations.

$$\Psi_i(P_i) = \int_{\mathcal{A}} \max \left\{ 0, \Phi_i(a_i, P_i) \right\} da_i.$$

Minimize the violations

$$\hat{P}_i \in \arg\min_{P_i \in \mathcal{P}} \left\{ \Psi_i(P_i) \right\} \iff \hat{P}_i \in \arg\max_{P_i \in \mathcal{P}} \left\{ L_i(P_i) \right\},$$

where $L_i(P_i) = -\Psi_i(P_i)$ is the likelihood function.








Figure: Australian bank data set, Pattison et al. (2000).









Low hierarchical positions occupy the periphery of the network.



Presence of star-like motifs embedded in the network (e.g. Branch Manager).





- Competitive behavior and reciprocity of high-ranking positions.
- Low-ranking positions inclined towards social support.




Figure: Maximum likelihood estimate of strategic behavior \hat{P}_i .



Summary








$$V_i(a_i, \mathbf{a_{-i}}|P_i) = \alpha_i t_i(a_i, \mathbf{a_{-i}}) + \beta_i u_i(a_i, \mathbf{a_{-i}}) - \gamma_i c_i(a_i),$$

$$\alpha_i \geq 0, \, \beta_i \in \mathbb{R}, \, \gamma_i > 0.$$



 $pagann@ethz.ch, \quad people.ee.ethz.ch/{\sim}pagann/$

- Bala, V. and Goyal, S. (2000). A Noncooperative Model of Network Formation. *Econometrica*, 68(5):1181–1229.
- Buechel, B. and Buskens, V. (2013). The Dynamics of Closeness and Betweenness. *The Journal of Mathematical Sociology*, 37(July):159–191.
- Burger, M. J. and Buskens, V. (2009). Social context and network formation: An experimental study. *Social Networks*, 31(1):63–75.
- Burt, R. S. (1992). Structural hole. Harvard Business School Press, Cambridge, MA.
- Buskens, V. and van de Rijt, A. (2008). Dynamics of Networks if Everyone Strives for Structural Holes. *American Journal of Sociology*, 114(2):371–407.
- Coleman, J. (1990). Foundations of social theory. Cambridge, MA: Belknap.
- Heider, F. (1946). Attitudes and cognitive organization. *The Journal of psychology*, 21(1):107–112.
- Jackson, M. O. and Wolinsky, A. (1996). A Strategic Model of Social and Economic Networks. *Journal of Economic Theory*, 71(1):44–74.
- Pattison, P., Wasserman, S., Robins, G., and Kanfer, A. M. (2000). Statistical evaluation of algebraic constraints for social networks. *Journal of mathematical psychology*, 44(4):536–568.

