

Social network formation: from individuals incentives to systemic stability

Nicolò Pagan¹ Florian Dörfler¹

¹ Automatic Control Laboratory, ETH Zürich, Switzerland

GAMENET at NetSci 2018

International School and Conference on Network Science

Paris, 12.06.2018

Social Networks



Social Networks provide benefits

Social Networks provide benefits

Social Influence



The more people we are connected to, the more we can influence them.


Social Networks provide benefits

Social Influence



The more people we are connected to, the more we can influence them.

Safety




The more our friends' friends are our friends, the safer we feel.

[Heider (1946); Coleman (1990)]
(Structural Balance theory)


Social Networks provide benefits

Social Influence




The more people we are connected to, the more we can influence them.

Safety



The more our friends' friends are our friends, the safer we feel.
[Heider (1946); Coleman (1990)]
(Structural Balance theory)

Efficacy



The more we are on the path between people, the more we can control.
[Burt (1992)]
(Structural Holes theory)

Social Networks provide benefits


Social Influence



The more people we are connected to, the more we can influence them.

Degree Centrality

Safety




The more our friends' friends are our friends, the safer we feel.

[Heider (1946); Coleman (1990)]
(Structural Balance theory)

Clustering Coefficient

Efficacy




The more we are on the path between people, the more we can control.

[Burt (1992)]
(Structural Holes theory)

Betweenness Centrality

Network Formation




Analysis of the co-evolution of individual behavior and network topology.

- Probabilistic approach
- **Game Theoretical approach** (Strategic Network Formation)

Social Network Formation Model

Ingredients


- **Directed weighted** network \mathcal{G} with $N \geq 3$ agents.



Social Network Formation Model

Ingredients


- ▶ **Directed weighted** network \mathcal{G} with $N \geq 3$ agents.
- ▶ $a_{ij} \in [0, 1]$ quantifies the importance of the friendship among i and j **from i 's point of view**.



Social Network Formation Model

Ingredients

- ▶ **Directed weighted** network \mathcal{G} with $N \geq 3$ agents.
- ▶ $a_{ij} \in [0, 1]$ quantifies the importance of the friendship among i and j **from i 's point of view**.
- ▶ **Homogeneous** agents.




Social Network Formation Model

Ingredients

- ▶ **Directed weighted** network \mathcal{G} with $N \geq 3$ agents.
- ▶ $a_{ij} \in [0, 1]$ quantifies the importance of the friendship among i and j **from i 's point of view**.
- ▶ **Homogeneous** agents.
- ▶ A typical action of agent i is:

$$a_i = [a_{i1}, \dots, a_{i,i-1}, a_{i,i+1}, \dots, a_{iN}]$$

living in the action space $\mathcal{A} = [0, 1]^{N-1}$.



Social Network Formation Model

Ingredients


- ▶ **Directed weighted** network \mathcal{G} with $N \geq 3$ agents.
- ▶ $a_{ij} \in [0, 1]$ quantifies the importance of the friendship among i and j **from i 's point of view**.
- ▶ **Homogeneous** agents.
- ▶ A typical action of agent i is:

$$a_i = [a_{i1}, \dots, a_{i,i-1}, a_{i,i+1}, \dots, a_{iN}]$$

living in the action space $\mathcal{A} = [0, 1]^{N-1}$.


- ▶ **Rational** agents: i is endowed with a payoff function V_i and is looking for

$$a_i^* \in \arg \max_{a_i \in \mathcal{A}} V_i(a_i, \mathbf{a}_{-i})$$



From individual incentives to the payoff function

$$V_i(a_i, \mathbf{a}_{-i}) =$$




From individual incentives to the payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = t_i(a_i, \mathbf{a}_{-i})$$

- **Social influence** on friends,

$$t_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ji}$$




From individual incentives to the payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = t_i(a_i, \mathbf{a}_{-i})$$

- **Social influence** on friends, on friends of friends, with $\delta \in [0, 1]$:

$$t_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ji} + \underbrace{\delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji}}_{\text{paths of length 2}}$$




From individual incentives to the payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = t_i(a_i, \mathbf{a}_{-i})$$

- **Social influence** on friends, on friends of friends, ... with $\delta \in [0, 1]$:

$$t_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ji} + \underbrace{\delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji}}_{\text{paths of length 2}} + \underbrace{\delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}}_{\text{paths of length 3}},$$




From individual incentives to the payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = t_i(a_i, \mathbf{a}_{-i})$$

- **Social influence** on friends, on friends of friends, ... with $\delta \in [0, 1]$:

$$t_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ji} + \underbrace{\delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji}}_{\text{paths of length 2}} + \underbrace{\delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}}_{\text{paths of length 3}},$$

[Jackson and Wolinsky (1996)]



From individual incentives to the payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = t_i(a_i, \mathbf{a}_{-i}) + u_i(a_i, \mathbf{a}_{-i})$$


- Social influence** on friends, on friends of friends, ... with $\delta \in [0, 1]$:

$$t_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ji} + \underbrace{\delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji}}_{\text{paths of length 2}} + \underbrace{\delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}}_{\text{paths of length 3}},$$

[Jackson and Wolinsky (1996)]

- Clustering coefficient**: number of closed triads:

$$u_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i, j} a_{ik} a_{kj} \right),$$



From individual incentives to the payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = t_i(a_i, \mathbf{a}_{-i}) \pm u_i(a_i, \mathbf{a}_{-i})$$

- **Social influence** on friends, on friends of friends, ... with $\delta \in [0, 1]$:


$$t_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ji} + \underbrace{\delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji}}_{\text{paths of length 2}} + \underbrace{\delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}}_{\text{paths of length 3}},$$

[Jackson and Wolinsky (1996)]

- **Clustering coefficient**: number of closed triads:

$$u_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i, j} a_{ik} a_{kj} \right),$$

[Burger and Buskens (2009)]



From individual incentives to the payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = t_i(a_i, \mathbf{a}_{-i}) \pm u_i(a_i, \mathbf{a}_{-i}) - c_i(a_i),$$

- **Social influence** on friends, on friends of friends, ... with $\delta \in [0, 1]$:

$$t_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ji} + \underbrace{\delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji}}_{\text{paths of length 2}} + \underbrace{\delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}}_{\text{paths of length 3}},$$

[Jackson and Wolinsky (1996)]


- **Clustering coefficient**: number of closed triads:

$$u_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i, j} a_{ik} a_{kj} \right),$$

[Burger and Buskens (2009)]

- **Cost of maintaining ties**:

$$c_i(a_i) = \sum_{j \neq i} a_{ij}.$$



From individual incentives to the payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha t_i(a_i, \mathbf{a}_{-i}) + \beta u_i(a_i, \mathbf{a}_{-i}) - \gamma c_i(a_i), \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0$$

- **Social influence** on friends, on friends of friends, ... with $\delta \in [0, 1]$:

$$t_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ji} + \underbrace{\delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji}}_{\text{paths of length 2}} + \underbrace{\delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji}}_{\text{paths of length 3}},$$

[Jackson and Wolinsky (1996)]


- **Clustering coefficient**: number of closed triads:

$$u_i(a_i, \mathbf{a}_{-i}) = \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i, j} a_{ik} a_{kj} \right),$$

[Burger and Buskens (2009)]


- **Cost of maintaining ties**:

$$c_i(a_i) = \sum_{j \neq i} a_{ij}.$$



Stability Definitions


Nash equilibrium (NE)



$$V_i(\mathbf{a}_i, \mathbf{a}_{-i}^*)$$

Stability Definitions

Nash equilibrium (NE)




C 1. for all agents i ,


$$V_i(a_i, \mathbf{a}_{-i}^*) \leq V_i(a_i^*, \mathbf{a}_{-i}^*), \forall a_i \in \mathcal{A}.$$

Stability Definitions

Nash equilibrium (NE)



Pairwise-Nash equilibrium (PNE)




C 1. for all agents i ,

$$V_i(a_i, \mathbf{a}_{-i}^*) \leq V_i(a_i^*, \mathbf{a}_{-i}^*), \forall a_i \in \mathcal{A}.$$

$$V_i(a_{ij}, a_{ji}, \mathbf{a}_{-(i,j)}^*), V_j(a_{ji}, a_{ij}, \mathbf{a}_{-(i,j)}^*)$$

Stability Definitions


Nash equilibrium (NE)



C 1. for all agents i ,

$$V_i(\mathbf{a}_i, \mathbf{a}_{-i}^*) \leq V_i(a_i^*, \mathbf{a}_{-i}^*), \forall \mathbf{a}_i \in \mathcal{A}.$$

Pairwise-Nash equilibrium (PNE)



C 2. for all agents i and j , and $\forall \mathbf{a}_{ij} \in [0, 1]$,

$$V_i(\mathbf{a}_{ij}, \mathbf{a}_{i-(i,j)}^*, \mathbf{a}_{-i}^*) \leq V_i(a_{ij}^*, \mathbf{a}_{i-(i,j)}^*, \mathbf{a}_{-i}^*),$$

C 3. for all pairs (i, j) , and $\forall (\mathbf{a}_{ij}, \mathbf{a}_{ji}) \in [0, 1]^2$,


$$V_i(\mathbf{a}_{ij}, \mathbf{a}_{ji}, \mathbf{a}_{-(i,j)}^*) > V_i(a_{ij}^*, \mathbf{a}_{ji}^*, \mathbf{a}_{-(i,j)}^*)$$

\Downarrow

$$V_j(\mathbf{a}_{ji}, \mathbf{a}_{ij}, \mathbf{a}_{-(i,j)}^*) < V_j(a_{ji}^*, \mathbf{a}_{ij}^*, \mathbf{a}_{-(i,j)}^*).$$

Stability Definitions

Nash equilibrium (NE)




C 1. for all agents i ,

$$V_i(\mathbf{a}_i, \mathbf{a}_{-i}^*) \leq V_i(\mathbf{a}_i^*, \mathbf{a}_{-i}^*), \forall \mathbf{a}_i \in \mathcal{A}.$$

C 1. and **C 2.** are selfish conditions.

C 3. is a cooperative (Pareto optimality) condition.

Pairwise-Nash equilibrium (PNE)



C 2. for all agents i and j , and $\forall \mathbf{a}_{ij} \in [0, 1]$,

$$V_i(\mathbf{a}_{ij}, \mathbf{a}_{i-(i,j)}^*, \mathbf{a}_{-i}^*) \leq V_i(\mathbf{a}_{ij}^*, \mathbf{a}_{i-(i,j)}^*, \mathbf{a}_{-i}^*),$$


C 3. for all pairs (i, j) , and $\forall (\mathbf{a}_{ij}, \mathbf{a}_{ji}) \in [0, 1]^2$,

$$V_i(\mathbf{a}_{ij}, \mathbf{a}_{ji}, \mathbf{a}_{-(i,j)}^*) > V_i(\mathbf{a}_{ij}^*, \mathbf{a}_{ji}^*, \mathbf{a}_{-(i,j)}^*)$$


\Downarrow

$$V_j(\mathbf{a}_{ji}, \mathbf{a}_{ij}, \mathbf{a}_{-(i,j)}^*) < V_j(\mathbf{a}_{ji}^*, \mathbf{a}_{ij}^*, \mathbf{a}_{-(i,j)}^*).$$

Stability Conditions



Stability Conditions




Definition (Nash equilibrium).

The network \mathcal{G}^* is a NE if for all agents i

$$V_i(a_i, \mathbf{a}_{-i}^*) \leq V_i(a_i^*, \mathbf{a}_{-i}^*), \forall a_i \in \mathcal{A}.$$

Stability Conditions



Definition (Nash equilibrium).

The network \mathcal{G}^* is a NE if for all agents i

$$V_i(a_i, \mathbf{a}_{-i}^*) \leq V_i(a_i^*, \mathbf{a}_{-i}^*), \forall a_i \in \mathcal{A}.$$

Stability conditions depend on:


- network \mathcal{G}^* : $\{a_i^*, i \in \mathcal{N}\}$,
- parameters: $\{\alpha, \beta, \gamma, \delta, N\}$.

Remark (Payoff function).

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha t_i(a_i, \mathbf{a}_{-i}, \delta) + \beta u_i(a_i, \mathbf{a}_{-i}) - \gamma c_i(a_i),$$

$\alpha \geq 0, \delta \in [0, 1], \quad \beta \in \mathbb{R}, \quad \gamma > 0.$

Stability Conditions



Definition (Nash equilibrium).

The network \mathcal{G}^* is a NE if for all agents i

$$V_i(a_i, \mathbf{a}_{-i}^*) \leq V_i(a_i^*, \mathbf{a}_{-i}^*), \forall a_i \in \mathcal{A}.$$

Stability conditions depend on:

- network \mathcal{G}^* : $\{a_i^*, i \in \mathcal{N}\}$,
- parameters: $\{\alpha, \beta, \gamma, \delta, N\}$.

Remark (Payoff function).

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha t_i(a_i, \mathbf{a}_{-i}, \delta) + \beta u_i(a_i, \mathbf{a}_{-i}) - \gamma c_i(a_i),$$

$\alpha \geq 0, \delta \in [0, 1], \quad \beta \in \mathbb{R}, \quad \gamma > 0.$

Question: For which parameters is a certain topology \mathcal{G}^* stable?

Network Topologies




Figure: Empty Network




Figure: Complete Network




Figure: Complete Balanced Bipartite Network






Figure: Star Network

Empty Network stability



Theorem .

- \mathcal{G}^{EN} is always a NE.
- \mathcal{G}^{EN} is a PNE if and only if



$$\frac{\alpha}{\gamma} \leq \frac{1}{(1 + \delta + \delta^2)}.$$

- No agent has a selfish incentive to create a link \rightarrow **always a NE**.
- Cooperation reduces the region of pairwise-Nash equilibrium.

Remark (Payoff function).

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha t_i(a_i, \mathbf{a}_{-i}, \delta) + \beta u_i(a_i, \mathbf{a}_{-i}) - \gamma c_i(a_i), \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0.$$

Complete Network stability



Theorem . Define

$$\bar{\gamma}_{NE} := \begin{cases} \alpha\delta(1 + \delta(2N - 3)) + \beta(N - 2), & \text{if } \beta > 0 \\ \alpha\delta(1 + \delta(2N - 3)) + 2\beta(N - 2), & \text{if } \beta \leq 0, \end{cases}$$

$$\bar{\gamma}_{PNE} := \alpha\delta(1 + \delta(2N - 3)) + 2\beta(N - 2),$$

then,



- \mathcal{G}^{CN} is a NE $\iff \gamma \leq \bar{\gamma}_{NE}$,
- \mathcal{G}^{CN} is a PNE $\iff \gamma \leq \bar{\gamma}_{PNE}$.

Remark (Payoff function).

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha t_i(a_i, \mathbf{a}_{-i}, \delta) + \beta u_i(a_i, \mathbf{a}_{-i}) - \gamma c_i(a_i), \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0.$$

- Complete network stability is correlated with **large** values of β (**high clustering**).
- Phase transition: Selfish agents prefer to drop all the links simultaneously.

Balanced Complete Bipartite Network stability





At equilibrium,

- ties across different node partitions should not be dismissed;
- ties within the same node partition should not be initiated.
- Balanced Complete Bipartite network stability is correlated with **negative** values of β (**betweenness centrality**).
- Cooperation reduces the region of pairwise-Nash equilibrium.

Remark (Payoff function).

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha t_i(a_i, \mathbf{a}_{-i}, \delta) + \beta u_i(a_i, \mathbf{a}_{-i}) - \gamma c_i(a_i), \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0.$$

Star stability



At equilibrium,

- ties from the central node should not be dismissed;
- ties to the central node should not be dismissed;
- ties between peripheral nodes should not be initiated.

- Star network stability needs **negative** values of β (**betweenness centrality**) and **large** values of α (**degree centrality**). However, an upper bound on α exists (for NE).
- Cooperation reduces the region of pairwise-Nash equilibrium.

Remark (Payoff function).

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha t_i(a_i, \mathbf{a}_{-i}, \delta) + \beta u_i(a_i, \mathbf{a}_{-i}) - \gamma c_i(a_i), \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0.$$

Phase diagram




Figure: Nash Equilibrium





Figure: Pairwise-Nash Equilibrium

- Different stable topologies can co-exist in the state space.
- Cooperation between agents often reduces the stability region.


Individual incentives and Pairwise-Nash stable topologies



Payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha \left(\sum_{j \neq i} a_{ji} + \delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji} + \delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji} \right) \\ + \beta \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i, j} a_{ik} a_{kj} \right) - \gamma \sum_{j \neq i} a_{ij}, \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0$$


Individual incentives and Pairwise-Nash stable topologies




- **Empty network stability and high cost** [Bala and Goyal (2000); Buechel and Buskens (2013)],

Payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha \left(\sum_{j \neq i} a_{ji} + \delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji} + \delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji} \right) + \beta \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i, j} a_{ik} a_{kj} \right) - \gamma \sum_{j \neq i} a_{ij}, \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0$$




Individual incentives and Pairwise-Nash stable topologies




- **Empty network stability and high cost** [Bala and Goyal (2000); Buechel and Buskens (2013)],
- **Complete network stability and high clustering** [Buechel and Buskens (2013)],

Payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha \left(\sum_{j \neq i} a_{ji} + \delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji} + \delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji} \right) + \beta \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i, j} a_{ik} a_{kj} \right) - \gamma \sum_{j \neq i} a_{ij}, \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0$$




Individual incentives and Pairwise-Nash stable topologies




- **Empty network** stability and **high cost** [Bala and Goyal (2000); Buechel and Buskens (2013)],
- **Complete network** stability and **high clustering** [Buechel and Buskens (2013)],
- **Balanced Complete Bipartite network** stability and **betweenness centrality** [Buskens and van de Rijt (2008)],

Payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha \left(\sum_{j \neq i} a_{ji} + \delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji} + \delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji} \right) + \beta \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i, j} a_{ik} a_{kj} \right) - \gamma \sum_{j \neq i} a_{ij}, \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0$$




Individual incentives and Pairwise-Nash stable topologies



- **Empty network** stability and **high cost** [Bala and Goyal (2000); Buechel and Buskens (2013)],
- **Complete network** stability and **high clustering** [Buechel and Buskens (2013)],
- **Balanced Complete Bipartite network** stability and **betweenness centrality** [Buskens and van de Rijt (2008)],
- **Star network** stability and **degree centrality** [Bala and Goyal (2000)].

Payoff function

$$V_i(a_i, \mathbf{a}_{-i}) = \alpha \left(\sum_{j \neq i} a_{ji} + \delta \sum_{k \neq j} \sum_{j \neq i} a_{kj} a_{ji} + \delta^2 \sum_{l \neq k} \sum_{k \neq j} \sum_{j \neq i} a_{lk} a_{kj} a_{ji} \right) + \beta \sum_{j \neq i} a_{ij} \left(\sum_{k \neq i, j} a_{ik} a_{kj} \right) - \gamma \sum_{j \neq i} a_{ij}, \quad \alpha \geq 0, \beta \in \mathbb{R}, \gamma > 0$$



- Bala, V. and Goyal, S. (2000). A Noncooperative Model of Network Formation. *Econometrica*, 68(5):1181–1229.
- Buechel, B. and Buskens, V. (2013). The Dynamics of Closeness and Betweenness. *The Journal of Mathematical Sociology*, 37(July):159–191.
- Burger, M. J. and Buskens, V. (2009). Social context and network formation: An experimental study. *Social Networks*, 31(1):63–75.
- Burt, R. S. (1992). Structural hole. *Harvard Business School Press, Cambridge, MA*.
- Buskens, V. and van de Rijt, A. (2008). Dynamics of Networks if Everyone Strives for Structural Holes. *American Journal of Sociology*, 114(2):371–407.
- Coleman, J. (1990). Foundations of social theory. *Cambridge, MA: Belknap*.
- Heider, F. (1946). Attitudes and cognitive organization. *The Journal of psychology*, 21(1):107–112.
- Jackson, M. O. and Wolinsky, A. (1996). A Strategic Model of Social and Economic Networks. *Journal of Economic Theory*, 71(1):44–74.