

Unknown is not normal: separating language-model extraction from rule-based decision logic for clinical risk scores

Nicolás Vera Zúñiga — Independent researcher, Chile — nicovera@quetru.cl

Abstract

Background. Large language models (LLMs) are increasingly used to compute clinical risk scores from free-text notes. Notes are often incomplete, and the usual convention of treating undocumented findings as normal can silently misclassify patients. An end-to-end LLM agent can instead ask the clinician for missing data, but it may ask unnecessary questions or answer before it has enough information.

Objective. To test whether separating calibrated, three-state extraction (an LLM) from decision logic (deterministic code computing score bounds over unknown inputs) reduces the questions asked without losing decision accuracy or safety, compared with asking for everything, with an end-to-end LLM agent, and with a binary missing-equals-normal schema.

Methods.

- *Cohort and notes.* 1,200 synthetic emergency cases, 200 each for HEART, CURB-65, qSOFA, PERC, Wells (pulmonary embolism) and Cockcroft-Gault. Notes were generated by an LLM from structured ground truth and validated for faithfulness.
- *Extraction.* An LLM labelled each input present, absent or unknown, with an exact evidence span and a confidence. Code computed the set of decision categories consistent with the unknowns and asked a simulated clinician only about inputs that could change the category.
- *Comparators.* Asking for every missing input; the same pipeline with missing inputs treated as normal; an end-to-end agent (Claude Opus 5.5) with calculator and ask-the-clinician tools; and a variant with value-of-information ordering and confirmation of low-confidence values.
- *Conditions.* Clean and messy notes, and an ideal and a noisy clinician (who does not know, misremembers, or answers vaguely).
- *External check.* 684 real case reports from MedCalc-Bench Verified.
- *Outcomes.* Decision-category accuracy (abstention counted as incorrect) and questions per case. Secondary outcomes were under- and over-triage with a higher-risk fallback, premature commitment, and irrelevant questions.

Results.

- *Ideal conditions.* With a small commercial model (Claude Haiku 4.5) as extractor, the bounds policy matched ask-all accuracy (99.4% vs 99.4%) while asking half as many questions (0.92 vs 1.78 per case; $p<0.001$) and no irrelevant ones (vs 48%).
- *Missing treated as normal.* This dropped accuracy to 91.2% and under-triaged 8.5% of patients (95% CI 7.1–10.2); the bounds policy under-triaged 0.1%. The binary schema's under-triage persisted in every condition (8.2–10.2%), even when its headline accuracy matched the bounds policy under a noisy clinician.
- *End-to-end agent.* The Opus agent was as accurate under ideal conditions (99.6%) but asked 9.5% irrelevant questions. With a noisy clinician it was less accurate than the bounds policy (83.5% vs 87.0%; $p<0.001$) and committed prematurely in 2.7% of cases (bounds: 0%).
- *Local model.* A 9B local model as extractor reached oracle-level accuracy (99.8%).
- *Real notes.* Only 52% of the 584 real case reports in the MedCalc-Bench training split contained enough information to determine the category (HEART 13%). Missing-equals-normal gave the correct category in 69%.

Conclusions. Treating undocumented findings as normal systematically under-triages patients, and benchmark conventions hide this. Routing decisions through code that reasons explicitly about unknowns removes premature commitment and irrelevant questions, halves the questions asked, and works with small local models. A frontier agent is competitive on accuracy but not on these safety properties.

1. Introduction

Clinical decision rules and scores (HEART for chest pain, CURB-65 for pneumonia, qSOFA, PERC and the Wells criteria for pulmonary embolism, and Cockcroft-Gault for renal dosing) are defined over a small set of inputs with explicit thresholds. Several groups now use LLMs to compute them from clinical text [MedCalc-Bench; MedRaC; MedMCP-Calc]. The standard benchmark, MedCalc-Bench, scores the numeric answer and, like most implementations, treats an input that the note does not mention as absent or normal [MedCalc-Bench].

In emergency practice that convention is not neutral. “No mention of hemoptysis” is not “no hemoptysis”, and a score computed as if it were can place a patient in a lower-risk category than their true one. The safe behaviour is to recognise when the documented facts do not determine the decision, and then to ask only for the facts that could change it. Recent work shows that LLMs handle this poorly in both directions: they commit to a score when the category is still undetermined, and they abstain when it is already determined [ClinDet-Bench]. Interactive benchmarks show that models asked to gather information before answering often ask too little or too much [MediQ; AgentClinic; CRAFT-MD].

We test a simple division of labour. A language model reads the note into typed, three-valued facts: present with value, explicitly absent, or unknown. Each fact carries an exact evidence span and a confidence. Deterministic code then computes the score’s possible values over the unknown inputs (its *bounds*), decides whether the decision category is already determined, and, if not, identifies the unknown inputs whose values could change the category. Only those are asked. We compare this design with asking for every missing input, with the same pipeline using a binary schema (missing = normal), and with an end-to-end frontier LLM agent that has a calculator and an ask-the-clinician tool.

We pre-specified four hypotheses:

- **H1.** The bounds policy asks far fewer questions than ask-all without losing accuracy.
- **H2.** An end-to-end agent both over-asks and under-asks.
- **H3.** Three-valued typing reduces silent missing-as-absent errors compared with a binary schema.
- **H4.** Confirming low-confidence, decision-critical extracted values catches most extraction errors at a small question cost.

2. Methods

2.1 Calculators and decision categories

Six calculators were implemented as pure functions from their primary sources [HEART; CURB-65; qSOFA/Sepsis-3; PERC; Wells; Cockcroft-Gault]. Decision categories were:

- HEART: low 0–3, moderate 4–6, high ≥ 7 .
- CURB-65: 0–1, 2, and ≥ 3 .
- qSOFA: positive if ≥ 2 .
- PERC: negative if all criteria are met.
- Wells: two-tier, PE likely if > 4 .
- Cockcroft-Gault: creatinine clearance < 30 , 30 to < 60 , and ≥ 60 mL/min.

Where the primary source was ambiguous, or differed from common implementations, we implemented a documented reading and listed it for review (Supplementary Table S1). Examples are the HEART troponin bands, the BUN equivalent of CURB-65 urea, and the PERC oxygen-saturation threshold. The author, an emergency physician, reviewed these readings.

Units are normalised by code, never by the model. For example, blood urea nitrogen or urea in mg/dL is converted to urea in mmol/L, and creatinine in $\mu\text{mol/L}$ to mg/dL.

2.2 Score bounds and decision relevance

Each input is boolean, ordinal, or numeric within a physiological range. Numeric inputs used through thresholds are divided into regions by the calculator’s cut-points, with the inclusivity of each cut-point ($\geq$ vs $>$) declared explicitly. The continuous score, Cockcroft-Gault, is monotone in each input.

Given a partial assignment (known values, intervals for inputs documented as “normal”, and unknowns), the achievable scores are enumerated exactly over the regions, or computed at interval corners for the continuous score. This yields the set of possible categories. The category is *determined* when only one category is possible. An unknown input is *decision-relevant* if changing its value changes the category for some completion of the other unknowns. This definition keeps inputs that matter only jointly.

Property-based tests with Hypothesis checked, for every calculator, that every completion of a partial assignment falls within the computed bounds, and that inputs labelled irrelevant never change the category.

2.3 Synthetic cohort

For each calculator we sampled hidden patient truth from plausible emergency-department priors (Supplementary Table S2). The author reviewed the priors. Each input was then marked documented positive, documented negative or normal, or not documented, with 30% missingness per input; systolic and diastolic blood pressure were documented together.

Rejection sampling produced exactly 50% of cases whose category was undetermined from the documented facts. This rejection preserves each stratum’s conditional distribution, so results can be reweighted to the natural share of undetermined cases (Supplementary Table S3).

Cases were tagged with five kinds of trap:

- an embedded negation;
- a value reported in another unit;
- a value from a previous encounter;
- a superseded same-visit reading, chosen where possible to change the category if misread;
- a comorbidity implied only by a medication.

Table 1 describes the cohort.

2.4 Clinical notes

Generation. Notes were written by an LLM (Claude Sonnet 5) from a per-case fact sheet. The fact sheet told the model to state each documented value exactly, to state negatives and normals explicitly, and to omit undocumented inputs entirely, including indirect cues such as rhythm descriptors that imply a heart rate, complete medication lists, or gendered wording. The note was never to name the score.

Validation. Every note passed two layers of validation:

- deterministic rule checks: required numbers present in exact surface form, and no score names;
- an independent LLM judge (Claude Opus 5.5), which classified every input as stated, negated or normal, not mentioned, or implied, compared against ground truth.

Notes that failed were regenerated up to twice (22 of 1,200 needed one retry, and all passed). The prompt was fixed at version v3 after two pilot rounds, and all 1,200 main notes used it.

Messy notes. A “messy” variant rendered 600 of the same cases (100 per calculator) from the same fact sheets in an end-of-shift style, with fragments, heavy abbreviation, typos outside numbers, copied-forward text and no headings. Of these, 41 still failed validation after three attempts and were excluded, leaving 559 cases paired across conditions.

2.5 Extraction

Schema. The extractor returned, for every input:

- a status (present, absent or unknown);
- a value in the input's domain;
- a unit token from a fixed list;
- the shortest exact supporting quote;
- a confidence.

Code-side checks. Code located the quote in the note and rejected any claim whose quote was not an exact substring; the input then became unknown. Code also normalised units, and rejected “absent” for inputs that cannot be negated (age, weight, sex) and values outside plausible ranges.

Extractors.

- Claude Haiku 4.5, accessed through headless Claude Code (`claude -p`) with a replaced system prompt, all tools disabled and JSON-schema-constrained output. Confidence was self-reported.
- Qwen3.5-9B (4-bit quantised), run locally through Ollama with a JSON schema, thinking disabled, temperature 0, and a fixed seed. Confidence was taken from token log-probabilities as $P(\text{status tokens}) \times P(\text{value tokens})$.

Calibration. Temperature scaling and isotonic regression were fitted on a seeded 30% development split per calculator and evaluated on the remainder. S4 used isotonic calibration.

2.6 Policies

- **S1 ask-all:** asks for every input the extraction left unknown.
- **S3 bounds:** asks only decision-relevant unknowns, one at a time, recomputing bounds after each answer.
- **S4:** S3 plus value-of-information ordering (the probability, under population priors, that the answer alone determines the category) and confirmation of extracted values whose calibrated confidence was below 0.9 and that were decision-critical.
- **S3-bin:** S3 with missing inputs treated as absent or normal. This is the MedCalc-Bench convention.
- **S2 end-to-end agent:** Claude Opus 5.5 received the note and parameter list and returned one JSON action per turn: ask the clinician about an input, run the calculator on supplied values, or answer with a category or “cannot determine”.

A code policy abstains when no remaining question can determine the category. For secondary analyses, an abstention is replaced by the highest-risk category still possible. For the agent, which does not track bounds, the calculator's highest-risk category is used.

2.7 Simulated clinician

A deterministic simulator answered from the hidden truth. Troponin was “not yet available” in 10% of cases. The noisy clinician, drawn independently per case and input, was:

- unable to answer 10% of questions;
- wrong 5% of the time (a yes/no flipped, a graded input one level off, or a number off by 10–25%);
- vague for 25% of numeric answers (a $\pm 10\%$ range containing the truth).

The noise does not depend on question order, so all policies face the same clinician.

2.8 Outcomes and statistics

Primary outcomes. Decision-category accuracy, with abstention counted as incorrect, and questions per case.

Secondary outcomes:

- accuracy when committing;
- under-triage (decision lower-risk than truth) and over-triage (decision higher-risk than truth), using the fallback;

- premature commitment (answering while the category was undetermined by the documented facts plus answers received);
- irrelevant questions (asked when the input could not change the category);
- silent missing-as-absent claims;
- extraction accuracy by documented state. A comorbidity conveyed only through a medication (e.g. lisinopril, never naming hypertension) is not determined by the note, so both “present” and “unknown” are accepted there, and “absent” is an error. For the same reason, the reference knowledge used to judge the agent’s questions and premature commitment treats such comorbidities as unknown;
- calibration (Brier score, expected calibration error).

Tests. Proportions have Wilson 95% confidence intervals and mean questions have 2,000-resample bootstrap intervals. Systems were compared on the same cases with exact McNemar tests (accuracy) and Wilcoxon signed-rank tests (questions). P-values were Holm-adjusted within each condition across four pre-specified comparisons (S3 vs S1, S3 vs S2, S3 vs S3-bin, S4 vs S3). Errors were attributed to abstention, extraction, routing (premature commitment), clinician error, or bounds.

2.9 External check on real notes

We used the MedCalc-Bench Verified release [MedCalc-Bench; label audits: Ye et al.; Krohn-Grimberghe] for the five overlapping calculators (qSOFA is not included). This comprised 100 test notes and a seeded sample of 584 training notes, with duplicates of test notes excluded. These are published, de-identified case reports licensed CC-BY-SA 4.0. They were used under a documented exception to the project’s synthetic-only rule and were not redistributed.

For each note we report:

- whether our calculator, applied to MedCalc’s annotated inputs, reproduces their label;
- the share of notes whose category our pipeline found determined;
- whether the true category remained within our bounds;
- category accuracy under MedCalc’s missing-equals-normal convention.

2.10 Implementation and reproducibility

Cost and model access. All runs used \$0 of API spend. Claude models were accessed through a Claude Code subscription, and Qwen ran on a laptop (Apple M1 Pro, 16 GB).

Reproducibility. Every run is driven by one configuration file and a fixed seed. Every model call was cached on disk under a hash of provider, model, prompt and parameters, so the author’s tables and figures regenerate from that cache without further model calls (calc-bounds paper configs/main.yaml). The cache is not released; re-running the pipeline re-queries the same models with the same prompts.

Code and tests. The code (Python 3.12, MIT licence; <https://github.com/nicoveraz/calc-bounds>) has 149 automated tests, including the property tests.

3. Results

3.1 Ideal conditions (Table 2, Figure 2)

H1. With oracle extraction, the bounds policy (S3) matched ask-all (S1) accuracy (99.8% vs 99.8%) with 0.92 vs 1.74 questions per case (Wilcoxon $p < 0.001$). Of S1’s questions, 47% could not change the decision; S3 asked none. The pattern held with both LLM extractors:

- Haiku: 99.4% for both, 0.92 vs 1.78 questions.
- Qwen: 99.8% for both, 1.23 vs 2.21 questions.

Across missingness levels of 10%, 30% and 50%, reweighted to the natural share of undetermined cases, S3 needed 34–49% fewer questions than S1 at equal accuracy (Supplementary Table S3).

H3. Treating missing inputs as normal (S3-bin) reduced accuracy to 91.2–91.8% (McNemar $p < 0.001$ vs S3; 99–105 vs 2 discordant cases). S3-bin committed while the category was undetermined in 39–49% of cases and under-triaged 8.2–8.8% of patients. S3 under-triaged 0.0–0.1%.

The agent under ideal conditions. The Opus agent reached 99.6% accuracy with 0.99 questions per case. It was not significantly different from S3 in accuracy, and it asked more questions ($p < 0.001$). Its behaviour included:

- 9.5% of its questions could not change the decision;
- it answered prematurely in 0.2% of cases.

H4. Confirming low-confidence values (S4) raised Haiku-based accuracy from 99.4% to 99.7% and asked fewer questions overall than S3 (0.90 vs 0.92; $p = 0.005$), because value-of-information ordering offset the confirmation questions. The accuracy gain was not significant (3 vs 0 discordant cases, adjusted $p = 0.75$).

Extractors. Both were accurate and conservative:

- *Accuracy of claims.* 99.6% of Haiku’s and 99.5% of Qwen’s present/absent claims were correct.
- *How Qwen fails.* Qwen missed 17% of documented negatives, but it labelled them unknown rather than absent, and S3 recovered them by asking.
- *How Haiku fails.* Haiku converted an undocumented input into “absent” in 16 claims (0.022 per case). This is the error H3 targets, and it accounts for most of Haiku’s accuracy loss under S3.
- *Calibration.* Isotonic calibration reduced expected calibration error from 0.014 to 0.002 (Haiku) and from 0.048 to 0.002 (Qwen).

3.2 Harder conditions (Table 3, Figures 2–3)

Noisy clinician. A noisy clinician reduced every system’s accuracy by 12–16 points, mainly through abstentions: a “don’t know” to the one decisive question leaves the category open. When committing, S1, S3 and S4 were 96.8–97.8% correct.

The agent under noise (full cohort, Haiku extraction):

- The agent was less accurate than S3 (83.5% vs 87.0%; 23 vs 65 discordant cases; adjusted $p < 0.001$).
- It committed prematurely in 2.7% of cases; the code policies did so in none.

The binary schema under noise. S3-bin’s headline accuracy (86.6–88.4%) matched S3’s, because it rarely asked and so rarely met a “don’t know”. When it committed, however, it was correct only 89–91% of the time, and it continued to under-triage 8.4–10.2% of patients (Figure 3).

Under- and over-triage with the fallback.

	S3 bounds	S3-bin	Agent
Under-triage, all conditions	0.0–0.5%	8.2–10.2%	up to 1.1%
Over-triage, noisy clinician	9.6–12.9% (S3 and S4)	lower	12.0–13.8%

With the fallback, S3 and S4 over-triaged when information was genuinely unavailable, which is the safe direction.

Messy notes (559 paired cases, ideal clinician).

- Haiku’s S3 accuracy fell from 99.1% on clean notes to 98.6% on messy notes, and its silent missing-as-absent rate doubled (0.045 per case). S4 recovered part of this (99.1%).
- Qwen was unaffected (99.5%).
- The agent reading messy notes directly was more accurate than S3 with Haiku extraction (99.8% vs 98.6%; 7 vs 0 discordant cases; adjusted $p = 0.047$) and similar to S3 with Qwen extraction.
- With a noisy clinician on messy notes, S3 and the agent did not differ (85.5% vs 85.0%).

3.3 Real notes (Table 4, Figure 4)

Our code reproduced MedCalc-Bench’s labels from its annotated inputs for 100% of HEART, CURB-65, PERC and Wells notes. Cockcroft-Gault reproduced only 51% of labels, because MedCalc-Bench chooses the weight input by body-mass index (actual, ideal or adjusted), whereas we used actual weight. One test-split HEART disagreement came from counting a transient ischaemic attack as atherosclerotic disease.

On 584 real training-split case reports:

- Only 52% (95% CI 48–56) had a category determined by the documented facts with Haiku extraction, and 47% with Qwen.
- By calculator: PERC 90%, Cockcroft-Gault 88%, CURB-65 49%, Wells 19%, HEART 13%.
- Under MedCalc’s missing-equals-normal convention, the category was correct for 69% of notes (HEART 22%).
- The true category remained possible within our bounds for 93–95% of notes. The misses were extraction errors, mainly in HEART inputs (agreement with annotations 73–74%).

The 100-note test split gave similar results: 59% determined, and 97% with the truth still possible.

4. Discussion

4.1 Principal findings

The most consistent finding is a safety one. Treating undocumented findings as normal under-triaged about one patient in twelve in every condition we tested, including clean notes and a perfect clinician. On real case reports it produced the correct category for only about two-thirds of notes. Headline accuracy can hide this: under a noisy clinician the binary schema’s accuracy equalled that of the bounds policy, because a system that rarely asks rarely meets an unanswerable question.

Separating extraction from decision logic removed premature commitment and irrelevant questions entirely, and halved the number of questions compared with asking for everything. When information was genuinely unobtainable, the bounds policy defaulted to the higher-risk category.

A frontier end-to-end agent was a strong baseline. Under ideal conditions it was as accurate as the bounds policy, and on messy notes it was more accurate than the pipeline with a small commercial extractor. We therefore do not claim that the modular design is more accurate than a capable agent. Its advantages lie elsewhere:

- it made no premature commitments and asked no irrelevant questions;
- it had the lowest under-triage;
- it was deterministic and auditable;
- its behaviour depended on reading, not reasoning, so a 9B model running locally matched oracle accuracy. The agent, by contrast, needs a frontier model at every turn (about 2.5 turns per case).

4.2 Relation to prior work

ClinDet-Bench [ClinDet-Bench] formalises the same determinacy question over score ranges and shows that LLMs both commit prematurely and over-abstain. Our work adds the question-asking step and a comparison with an agent that can ask. MediQ, AgentClinic and CRAFT-MD [MediQ; AgentClinic; CRAFT-MD] evaluate information-seeking diagnostic dialogue rather than calculators.

MedCalc-Bench and its audits [MedCalc-Bench; Ye et al.; Krohn-Grimberghe] evaluate one-shot score computation. Our real-note analysis suggests that its missing-equals-normal convention substantially overstates how often a score can safely be computed from the note alone. The design connects to active feature acquisition, selective prediction, and partial evaluation of rules over unknown inputs. Healthcare standards for executable guidelines (CQL, FHIR Clinical Practice Guidelines) would be a natural host for the determinacy check.

4.3 Limitations

- **Synthetic notes.** The main cohort’s notes were generated and validated by LLMs from the same family as the Claude extractor. An exploratory analysis of 46 cases with notes from both Sonnet and Qwen suggested a small same-family advantage (+3.2 points, 95% CI 0.4–6.4). Synthetic notes are also more complete and cleaner than real ones, as the real-note analysis shows, so absolute accuracies are optimistic. Differences in question counts and safety are the more transferable results.
- **Simulated clinician.** It follows a simple parametric model and has not been calibrated to real clinicians.
- **Independent priors.** Patient characteristics were sampled independently (apart from blood pressure), which can produce implausible combinations, such as established coronary disease with no risk factors. It does not change scores, but it lowers the face validity of some cases. We flagged 12 such cases (all HEART: established coronary disease with no risk factors, or age under 40 with three or more risk factors). Excluding them left the results unchanged (Supplementary Table S10).
- **Model access.** Claude models were accessed through Claude Code’s headless mode rather than a bare API. This adds about 1,200 tokens of harness context, and their confidence is self-reported.
- **Real-note evaluation.** It does not include asking questions, because no real clinician was available to answer. Its labels have known errors [Ye et al.; Krohn-Grimberghe].
- **Calculator readings.** The readings (Supplementary Table S1) and the population priors were reviewed by a single physician, the author, not by an independent panel. Other readings would change some scores, but they apply equally to all strategies.
- **Near-ceiling accuracy.** Accuracy under ideal conditions leaves little room for differences, and several accuracy comparisons are underpowered.

4.4 Implications

For builders of clinical decision support and documentation tools:

- keep “unknown” distinct from “absent” end to end;
- decide with code whether the documented facts determine the decision;
- ask only what can change it;
- when information cannot be obtained, fall back to the higher-risk category.

For evaluators, benchmarks that fill missing inputs with normal values should also report how often the category is actually determined by the note, and how often the convention under-triages.

5. Conclusions

Unknown is not normal. Treating undocumented findings as normal quietly under-triages patients and inflates benchmark accuracy. A simple design in which a language model reads and code decides makes missing information explicit, asks only questions that matter, never commits prematurely, and works with small local models, while matching the accuracy of ask-everything policies and remaining competitive with a frontier agent.

Declarations

- **Data availability.** Only the code is released (see Code availability). The synthetic cohort is regenerated deterministically by the code from the configuration and seed. The synthetic notes, extractions, traces and the LLM response cache are not released; running the code regenerates them with the same prompts, models and seeds, but LLM outputs are not guaranteed to be byte-identical. MedCalc-Bench Verified is available from its authors (CC-BY-SA 4.0).
- **Code availability.** <https://github.com/nicoveraz/calc-bounds> (MIT licence).
- **Funding.** None.
- **Competing interests.** None.

- **Use of AI.** LLMs generated the synthetic notes and served as extractors, judge and agent, as described. Code and analysis were developed with AI assistance (Claude Code).
- **Ethics.** Synthetic data, plus published de-identified case reports; no patient data. Ethics review was not required.

References

[To be formatted for the target venue; all verified 2026-09-26.]

- **HEART.** Six AJ, Backus BE, Kelder JC. Chest pain in the emergency room: value of the HEART score. *Neth Heart J.* 2008;16(6):191–196.
- **CURB-65.** Lim WS, et al. Defining community acquired pneumonia severity on presentation to hospital. *Thorax.* 2003;58(5):377–382.
- **qSOFA/Sepsis-3.**
 - Seymour CW, et al. Assessment of clinical criteria for sepsis. *JAMA.* 2016;315(8):762–774.
 - Singer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA.* 2016;315(8):801–810.
- **PERC.**
 - Kline JA, et al. Clinical criteria to prevent unnecessary diagnostic testing in emergency department patients with suspected pulmonary embolism. *J Thromb Haemost.* 2004;2(8):1247–1255.
 - Kline JA, et al. Prospective multicenter evaluation of the pulmonary embolism rule-out criteria. *J Thromb Haemost.* 2008;6(5):772–780.
- **Wells.**
 - Wells PS, et al. Derivation of a simple clinical model to categorize patients probability of pulmonary embolism. *Thromb Haemost.* 2000;83(3):416–420.
 - van Belle A, et al. Effectiveness of managing suspected pulmonary embolism using an algorithm combining clinical probability, D-dimer testing, and computed tomography. *JAMA.* 2006;295(2):172–179.
- **Cockcroft-Gault.** Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron.* 1976;16(1):31–41.
- **MedCalc-Bench.** Khandekar N, et al. MedCalc-Bench: evaluating large language models for medical calculations. NeurIPS Datasets and Benchmarks, 2024. Verified release: huggingface.co/datasets/nsk7153/MedCalc-Bench-Verified.
- **Ye et al.** Scalable stewardship of an LLM-assisted clinical benchmark with physician oversight. arXiv:2512.19691, 2025–2026.
- **Krohn-Grimberghe.** arXiv:2603.02222, 2026 (audit of calculator implementations in MedCalc-Bench).
- **ClinDet-Bench.** Watanabe Y, et al. arXiv:2602.22771, 2026.
- **MediQ.** Li SS, et al. MediQ: question-asking LLMs and a benchmark for reliable interactive clinical reasoning. NeurIPS 2024.
- **AgentClinic.** Schmidgall S, et al. AgentClinic: a multimodal agent benchmark to evaluate AI in simulated clinical environments. arXiv:2405.07960.
- **CRAFT-MD.** Johri S, et al. An evaluation framework for clinical use of large language models in patient interaction tasks. *Nat Med.* 2025.
- **MedRaC.** From scores to steps. EMNLP 2025.
- **MedMCP-Calc.** ACL 2026; arXiv:2601.23049.

Figures

Tables

Table 1. Synthetic cohort (n = 1,200). Undetermined: category not determined by the documented facts (50% by design).




Figure 1: Pipeline. A language model reads the note into typed tri-state facts (present / absent / unknown, with an exact evidence span and a confidence). Code normalizes units, computes the score’s bounds over unknown inputs and, if the decision category is not determined, asks the clinician only for inputs that could change it; each answer updates the bounds.




Figure 2: Decision-category accuracy (abstention counted as incorrect) versus mean questions per case, Haiku 4.5 extraction, 559 cases with validated notes in both styles. Left: clean notes, ideal clinician. Right: messy notes, noisy clinician.




Figure 3: Under-triage (decision lower-risk than the truth) and over-triage (higher-risk) with a higher-risk fallback for undetermined cases, by system and condition (Haiku 4.5 extraction, 559 paired cases). Error bars: Wilson 95% CIs.




Figure 4: Real case reports (MedCalc-Bench Verified training split; Haiku 4.5 extraction): share whose decision category is determined by the documented facts, versus share whose category is correct when missing inputs are treated as normal.

Calculator	Cases	Parameters	True categories	Undetermined from note	Params not documented, %	Cases with traps
HEART	200	11	low 82, moderate 103, high 15	100	25	56
CURB-65	200	6	low 95, moderate 68, high 37	100	24	61
qSOFA	200	3	negative 154, positive 46	100	33	53
PERC	200	8	negative 86, positive 114	100	29	61
Wells PE	200	7	pe_unlikely 162, pe_likely 38	100	29	56
Cockcroft-Gault	200	4	crcl_lt_30 23, crcl_30_to_lt_60 91, crcl_ge_60 86	100	24	39

Table 2. Main results: clean notes, ideal clinician (n = 1,200 per row). S1 ask-all; S2 end-to-end agent (Opus 5.5; reads the notes, so identical across extraction rows); S3 bounds; S4 bounds + value-of-information ordering + confidence echo; S3-bin bounds with missing = normal. Accuracy counts abstention as incorrect; under-triage uses the higher-risk fallback. 95% CIs: Wilson (proportions), bootstrap (questions).

Extraction	System	Accuracy, % (95% CI)	Questions/case (95% CI)	Irrelevant, %	Premature, %	Under-triage, % (95% CI)
Oracle	S1	99.8 (99.4–100.0)	1.74 (1.67–1.82)	47.3	0.0	0.0 (0.0–0.3)
Oracle	S2	99.6 (99.0–99.8)	0.99 (0.92–1.06)	9.5	0.2	0.1 (0.0–0.5)
Oracle	S3	99.8 (99.4–100.0)	0.92 (0.85–0.99)	0.0	0.0	0.0 (0.0–0.3)
Oracle	S4	99.8 (99.4–100.0)	0.87 (0.81–0.93)	0.0	0.0	0.0 (0.0–0.3)
Oracle	S3-bin	91.8 (90.1–93.2)	0.21 (0.19–0.24)	0.0	39.2	8.2 (6.8–9.9)
Haiku 4.5	S1	99.4 (98.8–99.7)	1.78 (1.71–1.86)	48.1	0.0	0.1 (0.0–0.5)
Haiku 4.5	S2	99.6 (99.0–99.8)	0.99 (0.92–1.06)	9.5	0.2	0.1 (0.0–0.5)
Haiku 4.5	S3	99.4 (98.8–99.7)	0.92 (0.86–0.99)	0.0	0.0	0.1 (0.0–0.5)
Haiku 4.5	S4	99.7 (99.1–99.9)	0.90 (0.83–0.96)	0.0	0.0	0.1 (0.0–0.5)
Haiku 4.5	S3-bin	91.2 (89.4–92.6)	0.21 (0.18–0.23)	0.0	40.0	8.5 (7.1–10.2)
Qwen3.5-9B (local)	S1	99.8 (99.4–100.0)	2.21 (2.12–2.29)	44.3	0.0	0.0 (0.0–0.3)
Qwen3.5-9B (local)	S2	99.6 (99.0–99.8)	0.99 (0.92–1.06)	9.5	0.2	0.1 (0.0–0.5)
Qwen3.5-9B (local)	S3	99.8 (99.4–100.0)	1.23 (1.15–1.31)	0.0	0.0	0.0 (0.0–0.3)
Qwen3.5-9B (local)	S4	99.8 (99.4–100.0)	1.20 (1.13–1.27)	0.0	0.0	0.0 (0.0–0.3)
Qwen3.5-9B (local)	S3-bin	91.2 (89.5–92.7)	0.26 (0.23–0.29)	0.0	49.2	8.8 (7.3–10.5)

Table 3. Harder conditions (notes / clinician), Haiku 4.5 extraction, 559 paired cases; systems as in Table 2 (full grid including Qwen3.5-9B in Supplementary Table S6).

Condition	System	Accuracy, % (95% CI)	Correct if answered, %	Under- triage, % (95% CI)	Over-triage, %	Premature, %	Questions/case
clean / ideal	S1	99.1 (97.9–99.6)	99.3	0.2 (0.0–1.0)	0.7	0.0	1.71 (1.60–1.81)
clean / ideal	S2	99.6 (98.7–99.9)	99.8	0.2 (0.0–1.0)	0.2	0.2	0.92 (0.82–1.01)
clean / ideal	S3	99.1 (97.9–99.6)	99.3	0.2 (0.0–1.0)	0.7	0.0	0.85 (0.75–0.94)
clean / ideal	S4	99.6 (98.7–99.9)	99.8	0.2 (0.0–1.0)	0.2	0.0	0.82 (0.73–0.91)
clean / ideal	S3-bin	91.4 (88.8–93.5)	91.4	8.2 (6.2–10.8)	0.4	37.0	0.20 (0.16–0.24)

Condition	System	Accuracy, % (95% CI)	Correct if answered, %	Under- triage, % (95% CI)	Over-triage, %	Premature, %	Questions/case
clean / noisy	S1	88.2 (85.3–90.6)	97.6	0.5 (0.2–1.6)	8.8	0.0	1.71 (1.60–1.81)
clean / noisy	S2	84.4 (81.2–87.2)	97.7	0.5 (0.2–1.6)	13.1	1.8	0.95 (0.84–1.05)
clean / noisy	S3	88.2 (85.3–90.6)	97.6	0.5 (0.2–1.6)	8.8	0.0	0.88 (0.77–0.97)
clean / noisy	S4	88.7 (85.8–91.1)	98.0	0.5 (0.2–1.6)	8.2	0.0	0.86 (0.76–0.96)
clean / noisy	S3-bin	88.7 (85.8–91.1)	91.2	8.2 (6.2–10.8)	2.5	36.0	0.20 (0.16–0.25)
messy / ideal	S1	98.6 (97.2–99.3)	98.7	0.5 (0.2–1.6)	0.9	0.0	1.92 (1.80–2.04)
messy / ideal	S2	99.8 (99.0– 100.0)	100.0	0.0 (0.0–0.7)	0.2	0.7	0.92 (0.82–1.01)
messy / ideal	S3	98.6 (97.2–99.3)	98.7	0.5 (0.2–1.6)	0.9	0.0	1.04 (0.94–1.15)
messy / ideal	S4	99.1 (97.9–99.6)	99.3	0.5 (0.2–1.6)	0.4	0.0	1.04 (0.94–1.15)
messy / ideal	S3-bin	90.7 (88.0–92.8)	90.7	8.8 (6.7–11.4)	0.5	39.4	0.30 (0.23–0.36)
messy / noisy	S1	85.5 (82.3–88.2)	96.8	0.5 (0.2–1.6)	11.1	0.0	1.92 (1.80–2.04)
messy / noisy	S2	85.0 (81.8–87.7)	97.3	1.1 (0.5–2.3)	12.0	3.4	0.97 (0.87–1.08)
messy / noisy	S3	85.5 (82.3–88.2)	96.8	0.5 (0.2–1.6)	11.1	0.0	1.08 (0.97–1.20)
messy / noisy	S4	85.9 (82.7–88.5)	97.4	0.5 (0.2–1.6)	10.7	0.0	1.11 (1.00–1.23)
messy / noisy	S3-bin	86.8 (83.7–89.3)	90.3	8.6 (6.5–11.2)	3.8	38.3	0.31 (0.25–0.38)

Table 4. Real case reports (MedCalc-Bench Verified, training split sample; test split in Supplementary Table S8).

Extractor	Calculator	n	Code reproduces label, %	Determined from note, % (95% CI)	Truth still possible, %	Category correct, miss- ing=normal, %
Haiku 4.5	Cockcroft- Gault	125	51	88.0 (81.1–92.6)	85	71
Haiku 4.5	CURB-65	85	100	49.4 (39.0–59.8)	93	82
Haiku 4.5	HEART	125	100	12.8 (8.0–19.8)	90	22
Haiku 4.5	PERC	124	100	90.3 (83.8–94.4)	100	94
Haiku 4.5	Wells PE	125	100	19.2 (13.3–27.0)	98	81

Extractor	Calculator	n	Code reproduces label, %	Determined from note, % (95% CI)	Truth still possible, %	Category correct, missing=normal, %
Haiku 4.5	All	584	90	52.1 (48.0–56.1)	93	69
Qwen3.5-9B	Cockcroft-Gault	125	51	84.8 (77.5–90.0)	90	72
Qwen3.5-9B	CURB-65	85	100	41.2 (31.3–51.8)	95	81
Qwen3.5-9B	HEART	125	100	12.0 (7.4–18.9)	93	9
Qwen3.5-9B	PERC	124	100	87.9 (81.0–92.5)	100	91
Qwen3.5-9B	Wells PE	125	100	9.6 (5.6–16.0)	99	78
Qwen3.5-9B	All	584	90	47.4 (43.4–51.5)	95	65

Supplementary material: calculator review items (S1), priors (S2), missingness sensitivity with natural-share reweighting (S3), confidence-echo threshold sweep (S4), calibration (S5), full result grids and paired comparisons (S6–S7), real-note test split (S8), exact prompts (S9).

Supplementary material

S1. Calculator readings awaiting clinical review

Each calculator cites its primary source in code. Where the source was ambiguous or differed from common implementations, the implemented reading and the alternatives are listed below. The author, an emergency physician, reviewed and accepted the implemented readings.

Type:

- **S (source):** resolvable by reading the full text of the primary paper. These cluster on the abstract-only [A] sources, PERC and Wells 2000.
- **C (clinical):** needs clinical judgement.
- **M (modelling):** a modelling choice with no clinical answer.

Priority is face validity: how much a clinician reading a generated case would notice if it were wrong.

- **High:** visible in every case of that calculator.
- **Med:** visible in some cases.
- **Low:** internal only.

For a bounds benchmark most items only need to be *fixed and documented*: the ground truth is self-consistent either way.

#	Type	Priority	Calculator	Item	Implemented	Alternatives / notes
1	S	High	HEART	Troponin bands	≤ normal / 1–3× / > 3×, deliberately following the validation cohorts (Backus 2010/2013, Poldervaart 2013), not the 2008 table	Six 2008: 1–2× / > 2×. Exactly 3× at the band edge: ≥ 3× (Backus 2011, Poldervaart) vs > 3× (MDCalc)
27	C	High	HEART	Troponin assay reference	Troponin is a level relative to the assay's upper reference limit, accepted by the author. Checked: every HEART note states the level relative to the reference limit; the one raw value also gives the reference limit	Real hs-troponin uses sex-specific 99th percentiles. Out of scope while troponin is a ratio; needed for real data

#	Type	Priority	Calculator	Item	Implemented	Alternatives / notes
23	S	High	HEART	Age exactly 65	2 points (age ≥ 65), per the 2008 text (“65 years or older”)	2008 table says “ ≤ 65 ” (typo); some renderings use $45-65 \square 1, > 65 \square 2$ (65 scores 1)
2	S	High	HEART	Age exactly 45	1 point (age ≥ 45)	Poldervaart 2013: $\leq 45 \square 0$
4	C	High	HEART	Smoking recency	≤ 3 months (MDCalc), accepted by the author	< 1 month (Six 2008). No note states a quit date, so the choice affects no case
5	C	High	HEART	Family history	first-degree relative with CAD before 65, accepted by the author	Only 3 notes give an age (mother 54, brother 55, father 58), all clear-cut, so the choice affects no case
6	S	Med	HEART	Obesity	boolean, labelled BMI > 30	not defined in Six 2008; BMI > 30 in Poldervaart 2013
7	S	Med	HEART	TIA as atherosclerotic disease	included (2 points)	Six 2008 lists revascularisation, MI, stroke, PAD. MedCalc-Bench counts TIA as an ordinary risk factor (1 of 100 anchor disagreements)
3	S	Med	HEART	ECG 2 points	significant ST deviation (depression or elevation)	2008 table: ST depression only

#	Type	Priority	Calculator	Item	Implemented	Alternatives / notes
11	S	High	PERC	SaO2	positive if < 95%, on room air (the renderer states “on room air”)	2004 abstract: rule needs > 94%. “Sats normal on 2 L” is normal but PERC-positive in some readings; our notes never render supplemental O2
12	S	Med	PERC	Surgery/trauma	within 4 weeks requiring hospitalisation (2008 abstract)	requiring general anaesthesia (some sources)
13	S	Med	PERC	Hormone use	exogenous oestrogen	any hormone therapy (2004 abstract: “hormone use”)
24	S	Med	Wells	Item definitions	Stated in the parameter labels given to renderer and extractor: malignancy = on treatment, treated within 6 months, or palliative; DVT signs = leg swelling plus pain on deep-vein palpation; previous DVT/PE	Previous DVT/PE should say “objectively diagnosed” (label says “Previous DVT or PE”)
14	S	Med	Wells	Immobilisation duration	“> 3 days” (NICE)	“≥ 3 days” (MDCalc)

#	Type	Priority	Calculator	Item	Implemented	Alternatives / notes
25	S	Med	Wells	4-week window	label: "Immobalisation > 3 days or surgery in the previous 4 weeks" (window read as applying to both)	the window may apply to surgery only
15	S	Med	Wells	PE-likely item	"alternative diagnosis less likely than PE" (NICE)	"PE #1 or equally likely" (MDCalc)
8	C	Med	CURB-65	BUN equivalent	code converts BUN to urea mmol/L and applies > 7 (= BUN > 19.6)	MDCalc: BUN > 19; an audit suggests > 20

#	Type	Priority	Calculator	Item	Implemented	Alternatives / notes
26	C	High	CURB-65 / qSOFA	How mental status is extracted (accepted: GCS 14 □ present)	Bool params, labelled “Confusion (AMT $\leq$ 8 or new disorientation)” and “Altered mentation (GCS < 15)”. There is no GCS parameter: “GCS 14” □ altered mentation present; a vague “drowsy” is left to the extractor’s judgement. Checked: dementia, baseline impairment and vague words (“lethargic”) appear only where mental status is explicitly documented, never where it is undocumented, so no case is ambiguous. Baseline impairment is consistent with each definition (qSOFA counts any abnormal mentation; CURB-65 counts only new disorientation)	add a numeric GCS parameter (would change prompts and cohort)

#	Type	Priority	Calculator	Item	Implemented	Alternatives / notes
9	C	Low	CURB-65 / qSOFA	Confusion vs altered mentation	two separate parameters	Coupling is not needed: each case belongs to exactly one calculator, so one patient never carries both
10	C	Med	qSOFA	Altered mentation threshold	GCS < 15 (Sepsis-3)	GCS ≤ 13 (Seymour derivation model)
16	C	Med	Cockcroft-Gault	Weight	actual body weight (original)	ideal/adjusted weight in obesity; MedCalc-Bench picks by BMI (explains 49% anchor disagreement)
17	M	Low	Cockcroft-Gault	Decision thresholds	30, 60 mL/min (config-overrideable)	FDA 2024 Table 1 also has 90
18	M	Low	Units	Creatinine factor	88.4 μmol/L per mg/dL	88.42 (10,000 / 113.12). Clinically irrelevant; can move a boundary case by < 0.03% of CrCl. No case sits that close in our cohort
19	M	Med	Params	“Stated normal” intervals	see table below	—

#	Type	Priority	Calculator	Item	Implemented	Alternatives / notes
20	M	Low	Params	Creatinine can't be "normal"	Present or Unknown only	The no-straddle guarantee is per parameter and threshold. CG thresholds apply to a function of age, weight and creatinine, so no creatinine interval could guarantee a single CrCl band. For CG, bounds return a set of categories whenever an input is uncertain (100 of 200 CG cases are determined from the note by design)

#	Type	Priority	Calculator	Item	Implemented	Alternatives / notes
21	M	Med	Cohort	Population priors	independent sampling (plus DBP < SBP – 15)	Independence can yield implausible combinations (e.g. established CAD with no risk factors, or age < 40 with ≥ 3 risk factors: 12 HEART cases, 6% of HEART and 1% of the cohort). It doesn't affect scores (atherosclerotic disease scores 2 regardless), and excluding the flagged cases leaves results unchanged (supplementary Table S10). Reported as a limitation; correlated priors are future work

#	Type	Priority	Calculator	Item	Implemented	Alternatives / notes
22	C	Med	Cohort	Medication implying a comorbidity	Only metformin ☐ diabetes is reasonably specific. Lisinopril ☐ hypertension and atorvastatin ☐ hypercholesterolaemia are weak (ACE inhibitors in HF, post-MI and CKD; statins post-ACS and in diabetes). Scoring: for a comorbidity shown only through a medication, both Present and Unknown count as correct; only Absent is an error	treat the implication as probabilistic in the generator

S2. Population priors for the synthetic cohort

Plausible emergency-department populations for each calculator's intended use; parameters sampled independently except diastolic < systolic blood pressure – 15 mmHg. The PERC cohort represents low gestalt pre-test probability. Reviewed by the author.

Calculator	Parameter	Prior
HEART	heart_history	slightly_suspicious 0.45, moderately_suspicious 0.35, highly_suspicious 0.2
HEART	heart_ecg	normal 0.6, nonspecific_repolarization 0.3, significant_st_deviation 0.1
HEART	age	Normal(58, 15) truncated to [18, 100]
HEART	hypertension	P(yes) = 0.4
HEART	hypercholesterolemia	P(yes) = 0.3

Calculator	Parameter	Prior
HEART	diabetes	P(yes) = 0.2
HEART	obesity	P(yes) = 0.25
HEART	smoking	P(yes) = 0.25
HEART	family_history_cad	P(yes) = 0.2
HEART	atherosclerotic_disease	P(yes) = 0.2
HEART	heart_troponin	le_normal 0.75, 1_to_3x_normal 0.15, gt_3x_normal 0.1
CURB-65	confusion	P(yes) = 0.15
CURB-65	urea	Normal(7, 3.5) truncated to [1.5, 40]
CURB-65	resp_rate	Normal(22, 6) truncated to [8, 50]
CURB-65	sbp	Normal(125, 25) truncated to [60, 220]
CURB-65	dbp	Normal(72, 14) truncated to [30, 130]
CURB-65	age	Normal(65, 17) truncated to [18, 100]
qSOFA	resp_rate	Normal(21, 5) truncated to [8, 50]
qSOFA	altered_mentation	P(yes) = 0.2
qSOFA	sbp	Normal(118, 24) truncated to [60, 220]
PERC	age	Normal(38, 12) truncated to [18, 90]
PERC	heart_rate	Normal(88, 15) truncated to [45, 160]
PERC	spo2	Normal(97, 2) truncated to [85, 100]
PERC	unilateral_leg_swelling	P(yes) = 0.05
PERC	hemoptysis	P(yes) = 0.03
PERC	recent_surgery_trauma	P(yes) = 0.05
PERC	prior_vte	P(yes) = 0.05
PERC	hormone_use	P(yes) = 0.12
Wells PE	dvt_signs	P(yes) = 0.12
Wells PE	pe_most_likely	P(yes) = 0.3
Wells PE	heart_rate	Normal(95, 18) truncated to [45, 170]
Wells PE	immobilization_or_surgery	P(yes) = 0.12
Wells PE	prior_vte	P(yes) = 0.12
Wells PE	hemoptysis	P(yes) = 0.05
Wells PE	malignancy	P(yes) = 0.1
Cockcroft-Gault	age	Normal(60, 17) truncated to [18, 100]
Cockcroft-Gault	weight	Normal(75, 17) truncated to [35, 180]
Cockcroft-Gault	creatinine	Normal(1.3, 0.8) truncated to [0.4, 8]
Cockcroft-Gault	sex	male 0.5, female 0.5

S3. Missingness sensitivity (oracle extraction, natural share of undetermined cases)

Fresh cohorts (200 cases per calculator) at each missingness level; stratum results reweighted to the natural share of undetermined cases.

Missingness	System	Accuracy (natural share)	Questions/case (natural share)	Undetermined share
0.1	S1 Ask-all	0.999	0.642	0.252
0.1	S3-bin (missing=normal)	0.964	0.089	0.252
0.1	S3 Bounds	0.999	0.326	0.252

Missingness	System	Accuracy (natural share)	Questions/case (natural share)	Undetermined share
0.1	S4	0.999	0.316	0.252
	Bounds+VOI+echo			
0.3	S1 Ask-all	0.997	1.925	0.61
0.3	S3-bin	0.899	0.282	0.61
	(missing=normal)			
0.3	S3 Bounds	0.997	1.144	0.61
0.3	S4	0.997	1.082	0.61
	Bounds+VOI+echo			
0.5	S1 Ask-all	0.994	3.289	0.824
0.5	S3-bin	0.824	0.483	0.824
	(missing=normal)			
0.5	S3 Bounds	0.994	2.159	0.824
0.5	S4	0.994	1.979	0.824
	Bounds+VOI+echo			

S4. Confidence-echo threshold sweep (S4 policy, clean notes, ideal clinician)

extraction	echo_threshold	accuracy	mean_questions	echo_per_case	echo_caught_errors
haiku__sonnet	0.5	0.995	0.8833	0.005	5
haiku__sonnet	0.8	0.9967	0.895	0.015	9
haiku__sonnet	0.9	0.9967	0.895	0.015	9
haiku__sonnet	0.95	0.9967	0.895	0.015	9
haiku__sonnet	0.99	0.9983	1.705	0.8242	19
haiku__sonnet	0.999	0.9983	1.7075	0.8267	19
qwen_local__sonnet	0.5	0.9983	1.1783	0.0	0
qwen_local__sonnet	0.8	0.9983	1.1783	0.0	0
qwen_local__sonnet	0.9	0.9983	1.1983	0.02	2
qwen_local__sonnet	0.95	0.9983	1.2967	0.1183	5
qwen_local__sonnet	0.99	0.9983	1.8142	0.6358	6
qwen_local__sonnet	0.999	0.9983	1.8142	0.6358	6

S5. Calibration of extraction confidence

Present/absent claims; isotonic and temperature calibration fitted on a seeded 30% development split, evaluated on the rest. Haiku 4.5 (self-reported): ECE 0.014 raw, 0.003 temperature, 0.002 isotonic. Qwen3.5-9B (token log-probabilities): ECE 0.048 raw, 0.033 temperature, 0.002 isotonic.

S6. Full results grid (559 paired cases; both extractors, both note styles, both clinicians)




Figure 5: Reliability of raw extraction confidence (test split).

Condition	System	n	Accuracy, % (95% CI)	Correct when an- swering, %	Under- triage, % (95% CI)	Over- triage, %	Premature commit- ment, %	Questions/cases- (95% CI)	Irrelevant cases- tions, %
Haiku, clean notes, ideal clinician	S1 Ask-all	559	99.1 (97.9– 99.6)	99.3	0.2 (0.0–1.0)	0.7	0.0	1.71 (1.60– 1.81)	50.0
Haiku, clean notes, ideal clinician	S2 LLM agent	559	99.6 (98.7– 99.9)	99.8	0.2 (0.0–1.0)	0.2	0.2	0.92 (0.82– 1.01)	10.1
Haiku, clean notes, ideal clinician	S3 Bounds	559	99.1 (97.9– 99.6)	99.3	0.2 (0.0–1.0)	0.7	0.0	0.85 (0.75– 0.94)	0.0
Haiku, clean notes, ideal clinician	S4 Bounds+VOI+echo	559	99.6 (98.7– 99.9)	99.8	0.2 (0.0–1.0)	0.2	0.0	0.82 (0.73– 0.91)	0.0
Haiku, clean notes, ideal clinician	S3-bin (miss- ing=normal)	559	91.4 (88.8– 93.5)	91.4	8.2 (6.2– 10.8)	0.4	37.0	0.20 (0.16– 0.24)	0.0
Haiku, clean notes, noisy clinician	S1 Ask-all	559	88.2 (85.3– 90.6)	97.6	0.5 (0.2–1.6)	8.8	0.0	1.71 (1.60– 1.81)	48.6
Haiku, clean notes, noisy clinician	S2 LLM agent	559	84.4 (81.2– 87.2)	97.7	0.5 (0.2–1.6)	13.1	1.8	0.95 (0.84– 1.05)	8.5
Haiku, clean notes, noisy clinician	S3 Bounds	559	88.2 (85.3– 90.6)	97.6	0.5 (0.2–1.6)	8.8	0.0	0.88 (0.77– 0.97)	0.0
Haiku, clean notes, noisy clinician	S4 Bounds+VOI+echo	559	88.7 (85.8– 91.1)	98.0	0.5 (0.2–1.6)	8.2	0.0	0.86 (0.76– 0.96)	0.0

Condition	System	n	Accuracy, % (95% CI)	Correct when an- swering, %	Under- triage, % (95% CI)	Over- triage, %	Premature commit- ment, %	Questions/cases- (95% CI)	Irrelevant cases- tions, %
Haiku, clean notes, noisy clinician	S3-bin (miss- ing=normal)	559	88.7 (85.8– 91.1)	91.2	8.2 (6.2– 10.8)	2.5	36.0	0.20 (0.16– 0.25)	0.0
Haiku, messy notes, ideal clinician	S1 Ask-all	559	98.6 (97.2– 99.3)	98.7	0.5 (0.2–1.6)	0.9	0.0	1.92 (1.80– 2.04)	45.6
Haiku, messy notes, ideal clinician	S2 LLM agent	559	99.8 (99.0– 100.0)	100.0	0.0 (0.0–0.7)	0.2	0.7	0.92 (0.82– 1.01)	10.7
Haiku, messy notes, ideal clinician	S3 Bounds	559	98.6 (97.2– 99.3)	98.7	0.5 (0.2–1.6)	0.9	0.0	1.04 (0.94– 1.15)	0.0
Haiku, messy notes, ideal clinician	S4 Bounds+VOI+echo	559	99.1 (97.9– 99.6)	99.3	0.5 (0.2–1.6)	0.4	0.0	1.04 (0.94– 1.15)	0.0
Haiku, messy notes, ideal clinician	S3-bin (miss- ing=normal)	559	90.7 (88.0– 92.8)	90.7	8.8 (6.7– 11.4)	0.5	39.4	0.30 (0.23– 0.36)	0.0
Haiku, messy notes, noisy clinician	S1 Ask-all	559	85.5 (82.3– 88.2)	96.8	0.5 (0.2–1.6)	11.1	0.0	1.92 (1.80– 2.04)	43.6
Haiku, messy notes, noisy clinician	S2 LLM agent	559	85.0 (81.8– 87.7)	97.3	1.1 (0.5–2.3)	12.0	3.4	0.97 (0.87– 1.08)	10.1
Haiku, messy notes, noisy clinician	S3 Bounds	559	85.5 (82.3– 88.2)	96.8	0.5 (0.2–1.6)	11.1	0.0	1.08 (0.97– 1.20)	0.0

Condition	System	n	Accuracy, % (95% CI)	Correct when an- swering, %	Under- triage, % (95% CI)	Over- triage, %	Premature commit- ment, %	Questions/cases- (95% CI)	Irrelevant cases- tions, %
Haiku, messy notes, noisy clinician	S4 Bounds+VOI+echo	559	85.9 (82.7– 88.5)	97.4	0.5 (0.2–1.6)	10.7	0.0	1.11 (1.00– 1.23)	0.0
Haiku, messy notes, noisy clinician	S3-bin (miss- ing=normal)	559	86.8 (83.7– 89.3)	90.3	8.6 (6.5– 11.2)	3.8	38.3	0.31 (0.25– 0.38)	0.0
Qwen, clean notes, ideal clinician	S1 Ask-all	559	99.8 (99.0– 100.0)	100.0	0.0 (0.0–0.7)	0.2	0.0	2.15 (2.04– 2.27)	45.5
Qwen, clean notes, ideal clinician	S2 LLM agent	559	99.6 (98.7– 99.9)	99.8	0.2 (0.0–1.0)	0.2	0.2	0.92 (0.82– 1.01)	10.1
Qwen, clean notes, ideal clinician	S3 Bounds	559	99.8 (99.0– 100.0)	100.0	0.0 (0.0–0.7)	0.2	0.0	1.17 (1.06– 1.28)	0.0
Qwen, clean notes, ideal clinician	S4 Bounds+VOI+echo	559	99.8 (99.0– 100.0)	100.0	0.0 (0.0–0.7)	0.2	0.0	1.14 (1.04– 1.25)	0.0
Qwen, clean notes, ideal clinician	S3-bin (miss- ing=normal)	559	91.6 (89.0– 93.6)	91.6	8.4 (6.4– 11.0)	0.0	46.5	0.25 (0.21– 0.30)	0.0
Qwen, clean notes, noisy clinician	S1 Ask-all	559	84.4 (81.2– 87.2)	97.7	0.5 (0.2–1.6)	12.7	0.0	2.15 (2.04– 2.27)	44.0
Qwen, clean notes, noisy clinician	S2 LLM agent	559	84.4 (81.2– 87.2)	97.7	0.5 (0.2–1.6)	13.1	1.8	0.95 (0.84– 1.05)	8.5

Condition	System	n	Accuracy, % (95% CI)	Correct when an- swering, %	Under- triage, % (95% CI)	Over- triage, %	Premature commit- ment, %	Questions/cases- (95% CI)	Irrelevant questions, %
Qwen, clean notes, noisy clinician	S3 Bounds	559	84.4 (81.2– 87.2)	97.7	0.5 (0.2–1.6)	12.7	0.0	1.21 (1.09– 1.32)	0.0
Qwen, clean notes, noisy clinician	S4 Bounds+VOI+echo	559	84.4 (81.2– 87.2)	97.7	0.5 (0.2–1.6)	12.7	0.0	1.20 (1.09– 1.31)	0.0
Qwen, clean notes, noisy clinician	S3-bin (miss- ing=normal)	559	87.8 (84.9– 90.3)	91.1	8.6 (6.5– 11.2)	3.0	45.3	0.26 (0.22– 0.31)	0.0
Qwen, messy notes, ideal clinician	S1 Ask-all	559	99.5 (98.4– 99.8)	99.6	0.2 (0.0–1.0)	0.4	0.0	2.15 (2.03– 2.28)	45.7
Qwen, messy notes, ideal clinician	S2 LLM agent	559	99.8 (99.0– 100.0)	100.0	0.0 (0.0–0.7)	0.2	0.7	0.92 (0.82– 1.01)	10.7
Qwen, messy notes, ideal clinician	S3 Bounds	559	99.5 (98.4– 99.8)	99.6	0.2 (0.0–1.0)	0.4	0.0	1.17 (1.06– 1.28)	0.0
Qwen, messy notes, ideal clinician	S4 Bounds+VOI+echo	559	99.6 (98.7– 99.9)	99.8	0.0 (0.0–0.7)	0.4	0.0	1.22 (1.10– 1.33)	0.0
Qwen, messy notes, ideal clinician	S3-bin (miss- ing=normal)	559	90.0 (87.2– 92.2)	90.0	9.8 (7.6– 12.6)	0.2	47.6	0.23 (0.18– 0.27)	0.0
Qwen, messy notes, noisy clinician	S1 Ask-all	559	84.3 (81.0– 87.0)	97.5	0.4 (0.1–1.3)	12.7	0.0	2.15 (2.03– 2.28)	43.7

Condition	System	n	Accuracy, % (95% CI)	Correct when an- swering, %	Under- triage, % (95% CI)	Over- triage, %	Premature commit- ment, %	Questions/cases- (95% CI)	Irrelevant cases- tions, %
Qwen, messy notes, noisy clinician	S2 LLM agent	559	85.0 (81.8– 87.7)	97.3	1.1 (0.5–2.3)	12.0	3.4	0.97 (0.87– 1.08)	10.1
Qwen, messy notes, noisy clinician	S3 Bounds	559	84.3 (81.0– 87.0)	97.5	0.4 (0.1–1.3)	12.7	0.0	1.21 (1.09– 1.33)	0.0
Qwen, messy notes, noisy clinician	S4 Bounds+VOI+echo	559	84.3 (81.0– 87.0)	97.1	0.5 (0.2–1.6)	12.7	0.0	1.29 (1.17– 1.41)	0.0
Qwen, messy notes, noisy clinician	S3-bin (miss- ing=normal)	559	86.6 (83.5– 89.2)	89.3	10.2 (8.0– 13.0)	2.7	46.7	0.24 (0.19– 0.28)	0.0

S7. Paired comparisons (exact McNemar for accuracy, Wilcoxon signed-rank for questions; Holm-adjusted within condition)

Full cohort (n = 1,200):

Condition	Comparison	Accuracy A vs B, %	Discordant (A only / B only)	McNemar p (Holm)	Questions A vs B	Wilcoxon p (Holm)
Oracle, clean, ideal	S3 Bounds vs S1 Ask-all	99.8 vs 99.8	0 / 0	1.000	0.92 vs 1.74	<0.001
Oracle, clean, ideal	S3 Bounds vs S2 LLM agent	99.8 vs 99.6	3 / 0	0.750	0.92 vs 0.99	<0.001
Oracle, clean, ideal	S3 Bounds vs S3-bin (miss- ing=normal)	99.8 vs 91.8	99 / 2	<0.001	0.92 vs 0.21	<0.001
Oracle, clean, ideal	S4 Bounds+VOI+echo vs S3 Bounds	99.8 vs 99.8	0 / 0	1.000	0.87 vs 0.92	<0.001
Haiku, clean, ideal	S3 Bounds vs S1 Ask-all	99.4 vs 99.4	0 / 0	1.000	0.92 vs 1.78	<0.001
Haiku, clean, ideal	S3 Bounds vs S2 LLM agent	99.4 vs 99.6	3 / 5	1.000	0.92 vs 0.99	<0.001
Haiku, clean, ideal	S3 Bounds vs S3-bin (miss- ing=normal)	99.4 vs 91.2	101 / 2	<0.001	0.92 vs 0.21	<0.001

Condition	Comparison	Accuracy A vs B, %	Discordant (A only / B only)	McNemar p (Holm)	Questions A vs B	Wilcoxon p (Holm)
Haiku, clean, ideal	S4 Bounds+VOI+echo vs S3 Bounds	99.7 vs 99.4	3 / 0	0.750	0.90 vs 0.92	0.005
Qwen, clean, ideal	S3 Bounds vs S1 Ask-all	99.8 vs 99.8	0 / 0	1.000	1.23 vs 2.21	<0.001
Qwen, clean, ideal	S3 Bounds vs S2 LLM agent	99.8 vs 99.6	3 / 0	0.750	1.23 vs 0.99	<0.001
Qwen, clean, ideal	S3 Bounds vs S3-bin (missing=normal)	99.8 vs 91.2	105 / 2	<0.001	1.23 vs 0.26	<0.001
Qwen, clean, ideal	S4 Bounds+VOI+echo vs S3 Bounds	99.8 vs 99.8	0 / 0	1.000	1.20 vs 1.23	0.004
Haiku, clean, noisy	S3 Bounds vs S1 Ask-all	87.0 vs 87.0	0 / 0	1.000	0.96 vs 1.78	<0.001
Haiku, clean, noisy	S3 Bounds vs S2 LLM agent	87.0 vs 83.5	65 / 23	<0.001	0.96 vs 1.03	<0.001
Haiku, clean, noisy	S3 Bounds vs S3-bin (missing=normal)	87.0 vs 88.0	70 / 82	0.750	0.96 vs 0.21	<0.001
Haiku, clean, noisy	S4 Bounds+VOI+echo vs S3 Bounds	87.2 vs 87.0	3 / 0	0.750	0.94 vs 0.96	0.078

Paired subset (n = 559):

Condition	Comparison	Accuracy A vs B, %	Discordant (A only / B only)	McNemar p (Holm)	Questions A vs B	Wilcoxon p (Holm)
Haiku, clean notes, ideal clinician	S3 Bounds vs S1 Ask-all	99.1 vs 99.1	0 / 0	1.000	0.85 vs 1.71	<0.001
Haiku, clean notes, ideal clinician	S3 Bounds vs S2 LLM agent	99.1 vs 99.6	1 / 4	0.750	0.85 vs 0.92	0.002
Haiku, clean notes, ideal clinician	S3 Bounds vs S3-bin (missing=normal)	99.1 vs 91.4	45 / 2	<0.001	0.85 vs 0.20	<0.001
Haiku, clean notes, ideal clinician	S4 Bounds+VOI+echo vs S3 Bounds	99.6 vs 99.1	3 / 0	0.750	0.82 vs 0.85	0.023
Haiku, clean notes, noisy clinician	S3 Bounds vs S1 Ask-all	88.2 vs 88.2	0 / 0	1.000	0.88 vs 1.71	<0.001
Haiku, clean notes, noisy clinician	S3 Bounds vs S2 LLM agent	88.2 vs 84.4	34 / 13	0.012	0.88 vs 0.95	<0.001

Condition	Comparison	Accuracy A vs B, %	Discordant (A only / B only)	McNemar p (Holm)	Questions A vs B	Wilcoxon p (Holm)
Haiku, clean notes, noisy clinician	S3 Bounds vs S3-bin (missing=normal)	88.2 vs 88.7	31 / 34	1.000	0.88 vs 0.20	<0.001
Haiku, clean notes, noisy clinician	S4 Bounds+VOI+echo vs S3 Bounds	88.7 vs 88.2	3 / 0	0.750	0.86 vs 0.88	0.342
Haiku, messy notes, ideal clinician	S3 Bounds vs S1 Ask-all	98.6 vs 98.6	0 / 0	1.000	1.04 vs 1.92	<0.001
Haiku, messy notes, ideal clinician	S3 Bounds vs S2 LLM agent	98.6 vs 99.8	0 / 7	0.047	1.04 vs 0.92	0.002
Haiku, messy notes, ideal clinician	S3 Bounds vs S3-bin (missing=normal)	98.6 vs 90.7	46 / 2	<0.001	1.04 vs 0.30	<0.001
Haiku, messy notes, ideal clinician	S4 Bounds+VOI+echo vs S3 Bounds	99.1 vs 98.6	3 / 0	0.500	1.04 vs 1.04	0.933
Haiku, messy notes, noisy clinician	S3 Bounds vs S1 Ask-all	85.5 vs 85.5	0 / 0	1.000	1.08 vs 1.92	<0.001
Haiku, messy notes, noisy clinician	S3 Bounds vs S2 LLM agent	85.5 vs 85.0	30 / 27	1.000	1.08 vs 0.97	0.032
Haiku, messy notes, noisy clinician	S3 Bounds vs S3-bin (missing=normal)	85.5 vs 86.8	33 / 40	1.000	1.08 vs 0.31	<0.001
Haiku, messy notes, noisy clinician	S4 Bounds+VOI+echo vs S3 Bounds	85.9 vs 85.5	3 / 1	1.000	1.11 vs 1.08	0.048
Qwen, clean notes, ideal clinician	S3 Bounds vs S1 Ask-all	99.8 vs 99.8	0 / 0	1.000	1.17 vs 2.15	<0.001
Qwen, clean notes, ideal clinician	S3 Bounds vs S2 LLM agent	99.8 vs 99.6	1 / 0	1.000	1.17 vs 0.92	<0.001
Qwen, clean notes, ideal clinician	S3 Bounds vs S3-bin (missing=normal)	99.8 vs 91.6	47 / 1	<0.001	1.17 vs 0.25	<0.001
Qwen, clean notes, ideal clinician	S4 Bounds+VOI+echo vs S3 Bounds	99.8 vs 99.8	0 / 0	1.000	1.14 vs 1.17	0.033
Qwen, clean notes, noisy clinician	S3 Bounds vs S1 Ask-all	84.4 vs 84.4	0 / 0	1.000	1.21 vs 2.15	<0.001
Qwen, clean notes, noisy clinician	S3 Bounds vs S2 LLM agent	84.4 vs 84.4	32 / 32	1.000	1.21 vs 0.95	<0.001

Condition	Comparison	Accuracy A vs B, %	Discordant (A only / B only)	McNemar p (Holm)	Questions A vs B	Wilcoxon p (Holm)
Qwen, clean notes, noisy clinician	S3 Bounds vs S3-bin (missing=normal)	84.4 vs 87.8	34 / 53	0.212	1.21 vs 0.26	<0.001
Qwen, clean notes, noisy clinician	S4	84.4 vs 84.4	0 / 0	1.000	1.20 vs 1.21	0.960
Qwen, messy notes, ideal clinician	Bounds+VOI+echo vs S3 Bounds	99.5 vs 99.5	0 / 0	1.000	1.17 vs 2.15	<0.001
Qwen, messy notes, ideal clinician	S3 Bounds vs S1 Ask-all	99.5 vs 99.8	0 / 2	1.000	1.17 vs 0.92	<0.001
Qwen, messy notes, ideal clinician	S3 Bounds vs S3-bin (missing=normal)	99.5 vs 90.0	54 / 1	<0.001	1.17 vs 0.23	<0.001
Qwen, messy notes, ideal clinician	S4	99.6 vs 99.5	1 / 0	1.000	1.22 vs 1.17	0.006
Qwen, messy notes, ideal clinician	Bounds+VOI+echo vs S3 Bounds					
Qwen, messy notes, noisy clinician	S3 Bounds vs S1 Ask-all	84.3 vs 84.3	0 / 0	1.000	1.21 vs 2.15	<0.001
Qwen, messy notes, noisy clinician	S3 Bounds vs S2 LLM agent	84.3 vs 85.0	31 / 35	1.000	1.21 vs 0.97	<0.001
Qwen, messy notes, noisy clinician	S3 Bounds vs S3-bin (missing=normal)	84.3 vs 86.6	42 / 55	0.892	1.21 vs 0.24	<0.001
Qwen, messy notes, noisy clinician	S4	84.3 vs 84.3	2 / 2	1.000	1.29 vs 1.21	<0.001
Qwen, messy notes, noisy clinician	Bounds+VOI+echo vs S3 Bounds					

S8. Real case reports, all splits and extractors

Split	Extractor	Calculator	n	Code reproduces label, %	Determined from note, % (95% CI)	Truth still possible, %	Category correct, missing=normal, %	Entity agreement, %
train	Haiku 4.5	Cockcroft-Gault	125	51	88.0 (81.1–92.6)	85	71	90
train	Haiku 4.5	CURB-65	85	100	49.4 (39.0–59.8)	93	82	89
train	Haiku 4.5	HEART	125	100	12.8 (8.0–19.8)	90	22	73

Split	Extractor	Calculator	n	Code re- produces label, %	Determined from note, % (95% CI)	Truth still possible, %	Category correct, miss- ing=normal, %	Entity agree- ment, %
train	Haiku 4.5	PERC	124	100	90.3 (83.8– 94.4)	100	94	90
train	Haiku 4.5	Wells PE	125	100	19.2 (13.3– 27.0)	98	81	87
train	Haiku 4.5	All	584	90	52.1 (48.0– 56.1)	93	69	86
train	Qwen3.5- 9B	Cockcroft- Gault	125	51	84.8 (77.5– 90.0)	90	72	92
train	Qwen3.5- 9B	CURB-65	85	100	41.2 (31.3– 51.8)	95	81	92
train	Qwen3.5- 9B	HEART	125	100	12.0 (7.4–18.9)	93	9	74
train	Qwen3.5- 9B	PERC	124	100	87.9 (81.0– 92.5)	100	91	89
train	Qwen3.5- 9B	Wells PE	125	100	9.6 (5.6–16.0)	99	78	88
train	Qwen3.5- 9B	All	584	90	47.4 (43.4– 51.5)	95	65	87
test	Haiku 4.5	Cockcroft- Gault	20	40	95.0 (76.4– 99.1)	100	95	96
test	Haiku 4.5	CURB-65	20	100	55.0 (34.2– 74.2)	100	100	97
test	Haiku 4.5	HEART	20	95	60.0 (38.7– 78.1)	85	45	87
test	Haiku 4.5	PERC	20	100	75.0 (53.1– 88.8)	100	95	99
test	Haiku 4.5	Wells PE	20	100	10.0 (2.8–30.1)	100	85	89
test	Haiku 4.5	All	100	87	59.0 (49.2– 68.1)	97	84	94
test	Qwen3.5- 9B	Cockcroft- Gault	20	40	90.0 (69.9– 97.2)	100	90	96

Split	Extractor	Calculator	n	Code re-produces label, %	Determined from note, % (95% CI)	Truth still possible, %	Category correct, miss-ing=normal, %	Entity agree-ment, %
test	Qwen3.5-9B	CURB-65	20	100	50.0 (29.9–70.1)	95	90	98
test	Qwen3.5-9B	HEART	20	95	30.0 (14.5–51.9)	95	35	77
test	Qwen3.5-9B	PERC	20	100	75.0 (53.1–88.8)	100	95	96
test	Qwen3.5-9B	Wells PE	20	100	10.0 (2.8–30.1)	100	85	89
test	Qwen3.5-9B	All	100	87	51.0 (41.3–60.6)	98	79	91

S10. Sensitivity to implausible synthetic cases

Characteristics were sampled independently, so some cases are clinically implausible. We flagged established atherosclerotic disease with no risk factors, and age under 40 with three or more risk factors or atherosclerotic disease. We then recomputed the main results without those cases (clean notes, ideal clinician).

12 of 1200 cases flagged: heart-0017 (age < 40 with ≥ 3 risk factors or atherosclerotic disease); heart-0023 (age < 40 with ≥ 3 risk factors or atherosclerotic disease); heart-0036 (age < 40 with ≥ 3 risk factors or atherosclerotic disease); heart-0069 (age < 40 with ≥ 3 risk factors or atherosclerotic disease); heart-0087 (established atherosclerotic disease with no risk factors); heart-0104 (established atherosclerotic disease with no risk factors); heart-0128 (established atherosclerotic disease with no risk factors); heart-0132 (established atherosclerotic disease with no risk factors); heart-0135 (age < 40 with ≥ 3 risk factors or atherosclerotic disease); heart-0137 (established atherosclerotic disease with no risk factors); heart-0172 (age < 40 with ≥ 3 risk factors or atherosclerotic disease); heart-0195 (established atherosclerotic disease with no risk factors)

Extraction	Cases	System	n	Accuracy, %	Under-triage, %	Questions/case
Oracle	all	S1 Ask-all	1200	99.8	0.0	1.74
Oracle	all	S2 LLM agent	1200	99.6	0.1	0.99
Oracle	all	S3 Bounds	1200	99.8	0.0	0.92
Oracle	all	S4	1200	99.8	0.0	0.87
Oracle	all	Bounds+VOI+echo				
Oracle	all	S3-bin (miss-ing=normal)	1200	91.8	8.2	0.21
Oracle	excluding flagged	S1 Ask-all	1188	99.8	0.0	1.73
Oracle	excluding flagged	S2 LLM agent	1188	99.6	0.1	0.99
Oracle	excluding flagged	S3 Bounds	1188	99.8	0.0	0.92

Extraction	Cases	System	n	Accuracy, %	Under-triage, %	Questions/case
Oracle	excluding flagged	S4 Bounds+VOI+echo	1188	99.8	0.0	0.87
Oracle	excluding flagged	S3-bin (miss- ing=normal)	1188	91.9	8.1	0.21
Haiku 4.5	all	S1 Ask-all	1200	99.4	0.1	1.78
Haiku 4.5	all	S2 LLM agent	1200	99.6	0.1	0.99
Haiku 4.5	all	S3 Bounds	1200	99.4	0.1	0.92
Haiku 4.5	all	S4 Bounds+VOI+echo	1200	99.7	0.1	0.90
Haiku 4.5	all	S3-bin (miss- ing=normal)	1200	91.2	8.5	0.21
Haiku 4.5	excluding flagged	S1 Ask-all	1188	99.4	0.1	1.77
Haiku 4.5	excluding flagged	S2 LLM agent	1188	99.6	0.1	0.99
Haiku 4.5	excluding flagged	S3 Bounds	1188	99.4	0.1	0.93
Haiku 4.5	excluding flagged	S4 Bounds+VOI+echo	1188	99.7	0.1	0.90
Haiku 4.5	excluding flagged	S3-bin (miss- ing=normal)	1188	91.3	8.3	0.21
Qwen3.5-9B	all	S1 Ask-all	1200	99.8	0.0	2.21
Qwen3.5-9B	all	S2 LLM agent	1200	99.6	0.1	0.99
Qwen3.5-9B	all	S3 Bounds	1200	99.8	0.0	1.23
Qwen3.5-9B	all	S4 Bounds+VOI+echo	1200	99.8	0.0	1.20
Qwen3.5-9B	all	S3-bin (miss- ing=normal)	1200	91.2	8.8	0.26
Qwen3.5-9B	excluding flagged	S1 Ask-all	1188	99.8	0.0	2.20
Qwen3.5-9B	excluding flagged	S2 LLM agent	1188	99.6	0.1	0.99
Qwen3.5-9B	excluding flagged	S3 Bounds	1188	99.8	0.0	1.23
Qwen3.5-9B	excluding flagged	S4 Bounds+VOI+echo	1188	99.8	0.0	1.20
Qwen3.5-9B	excluding flagged	S3-bin (miss- ing=normal)	1188	91.4	8.6	0.26

S9. Exact prompts

Prompt versions: renderer v3, extractor x1, agent a1. The example fact sheet is for one CURB-65 case with a trap; <the rendered note> stands for the note text.

Renderer (system)

You write realistic clinical notes for a research dataset. Every patient is fictional; the notes a

Renderer (user), example CURB-65 case

Write one clinical note in US English (en-US).

Setting: Emergency department. Adult with suspected community-acquired pneumonia.

Style guide:

- US emergency department or inpatient documentation conventions.
- Typical sections: Chief complaint, HPI, PMH, Medications, Allergies, Social history, Exam (vitals first), Results, Assessment and plan. Not every section is required.
- Common US abbreviations are fine (HPI, PMH, SOB, CP, ROS, HR, RR, BP, SpO2, NKDA).
- Units: US conventional (mg/dL, lb or kg, °F or °C) except where the fact sheet gives a unit.
- Terse clinical style; sentence fragments are normal.

Fact sheet (each item is mandatory):

- [confusion] STATE: the patient is alert and fully oriented, not confused.
- [urea] STATE (number exactly): serum urea 8 mmol/L. Also mention a Serum urea (mmol/L) of 6.6 fr
- [resp_rate] DO NOT MENTION Respiratory rate (/min). Include nothing from which it could be infer
- [sbp] DO NOT MENTION Systolic blood pressure (mmHg). Include nothing from which it could be infe
- [dbp] DO NOT MENTION Diastolic blood pressure (mmHg). Include nothing from which it could be inf
- [age] DO NOT MENTION Age (years). Include nothing from which it could be inferred (no value, no

Rules:

- Write numbers exactly as given in the fact sheet (same digits and decimals), with the unit given
- Express each item in natural clinical language, as a clinician would chart it; do not copy the f
- Every STATE item must appear explicitly in the note, including negative ones (e.g. "no diabetes"
- Follow each item exactly. Do not add information about any fact-sheet item beyond what it says,
- No blanket or complete statements that would cover a DO NOT MENTION item: no "Medications: none"
- You may add other realistic details (presenting complaint, unrelated history, unrelated examinatio
- Do not name any clinical score or rule (HEART score, CURB-65, CURB65, qSOFA, PERC, Wells, Cockcr
- Do not refer to these instructions, the fact sheet, or the note being synthetic.
- Output only the note, 150–350 words, plain text (simple section headings allowed).

Judge (system)

You audit synthetic clinical notes against a list of parameters. For each parameter, decide only f

Judge (user)

Note:

```
<note>  
<the rendered note>  
</note>
```

Parameters:

- confusion: Confusion (AMT <= 8 or new disorientation to person/place/time) [yes/no finding]
- urea: Serum urea (mmol/L) [numeric (mmol/L)]
- resp_rate: Respiratory rate (/min) [numeric (/min)]
- sbp: Systolic blood pressure (mmHg) [numeric (mmHg)]
- dbp: Diastolic blood pressure (mmHg) [numeric (mmHg)]
- age: Age (years) [numeric (years)]

For every parameter, return one object:

`{"param": "<id>", "status": "stated" | "negated_or_normal" | "not_mentioned" | "implied", "level":`

- `stated`: the value or finding is explicitly documented (for yes/no findings: present).
- `negated_or_normal`: explicitly absent, denied, or documented as normal without a value.
- `not_mentioned`: nothing in the note documents or implies it.
- `implied`: not explicitly documented, but inferable from the note.

Return a JSON array with exactly one object per parameter, in the order listed, and nothing else.

Extractor (system)

You extract clinical parameters from a clinical note into a fixed JSON structure. You report only

Extractor (user)

Note:

`<note>`

`<the rendered note>`

`</note>`

Parameters:

- `confusion`: Confusion (AMT $\leq$ 8 or new disorientation to person/place/time) [yes/no finding]
- `urea`: Serum urea (mmol/L) [numeric; unit tokens: mmol/L, urea_mg/dL, bun_mg/dL]
- `resp_rate`: Respiratory rate (/min) [numeric; unit tokens: /min]
- `sbp`: Systolic blood pressure (mmHg) [numeric; unit tokens: mmHg]
- `dbp`: Diastolic blood pressure (mmHg) [numeric; unit tokens: mmHg]
- `age`: Age (years) [numeric; unit tokens: years]

For every parameter give:

- `"status"`:
 - `"present"`: the note states the finding, or states a value/level for it.
 - `"absent"`: the note explicitly says the finding is absent or denied, or that the parameter is not present.
 - `"unknown"`: the note does not document it. Do not infer absence from silence: if the note does not mention the parameter, it is unknown.
- `"value"`: for yes/no findings use null; for graded parameters one of the listed levels; for numeric parameters the value as written.
- `"unit"`: for numeric parameters, the unit as written, mapped to one of the listed unit tokens; otherwise null.
- `"evidence"`: the shortest exact quote from the note (copied character for character) that supports the status and value.
- `"confidence"`: your probability (0 to 1) that status and value are correct.

If the note gives several values for the same parameter (e.g. an earlier reading or a previous visit),

S2 agent (system)

You are a clinical decision-support agent working with an emergency clinician. You determine a patient's

S2 agent (first user turn)

Calculator: CURB-65

Decision categories: low, moderate, high

Clinical note:

`<note>`

`<the rendered note>`

`</note>`

Calculator parameters (ids you can ask about or pass to calculate):

- confusion: Confusion (AMT $\leq$ 8 or new disorientation to person/place/time) [yes/no]
- urea: Serum urea (mmol/L) [number in mmol/L]
- resp_rate: Respiratory rate (/min) [number in /min]
- sbp: Systolic blood pressure (mmHg) [number in mmHg]
- dbp: Diastolic blood pressure (mmHg) [number in mmHg]
- age: Age (years) [number in years]

Actions (reply with one JSON object per turn):

1. {"action": "ask", "parameter": "<id>", "question": "<your question to the clinician>"}
The clinician answers from the patient (or says the value is not available).
2. {"action": "calculate", "values": {"<id>": <value>, ...}}
Runs the calculator. Give every parameter: yes/no as true/false, graded ones as the level name,
3. {"action": "answer", "category": "<one of: low, moderate, high, cannot_determine>"}
Your final decision category. Use "cannot_determine" only if it truly cannot be determined.