CATCH-UP IMMUNIZATION

DR OSAMA HUSSEIN, MD

PEDIATRIC & NEONATOLOGY CONSULTANT

PORT SAID NEONATOLOGY SOCIETY

Hepatitis B vaccine

PNS
PORT SAID
NEONATOLOGY SOCIETY

Minimal interval

4 wk | 8 wk | 8 wk

24 wk

0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 4yr | 6yr | 8yr | 10yr | 12yr | 14yr | 16yr | 18yr

Dose 1 | Dose 2 | Dose 3 | Dose 4

Maximum interval

As soon as possible

Weight < 2000 g: Delay vaccination until 30 days of age, if born to an HBsAg- negative mother.



CDC

ACIP
Advisory Committee on Immunization Practices

BCG

Minimal interval

0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 4yr | 6yr | 8yr | 10yr | 12yr | 14yr | 16yr | 18yr

Dose

Maximum interval

Catch up




Infants below 3 months of age may be vaccinated with BCG vaccine.

Children 3 months to 3 years of age whose tuberculin pure protein derivative (TB-PPD)- or Bacilli Calmette Guérin pure protein derivative (BCG-PPD) test is negative and who have not been vaccinated should be provided with catch-up vaccination.

Children 4 years of age and older cannot receive BCG catch-up vaccination.

Children who have been vaccinated with BCG cannot receive BCG catch-up vaccination, even in the absence of a BCG scar.

Rotavirus vaccine






ACIP stands for Advisory Committee on Immunization Practices

Penta vaccine

Td

التطعيم الخماسى





CDC

ACIP
Advisory Committee on Immunization Practices

Pneumococcal vaccine

Minimal interval

4-8 wk

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Dose 1 | Dose 2 | Dose 3 | Dose 4

Maximum interval



ACIP stands for Advisory Committee on Immunization Practices

Pneumococcal vaccine

Minimal interval

Maximum interval

4 wk

8 wk

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Dose 1

Dose 2

Dose 3



CDC

ACIP stands for Advisory Committee on Immunization Practices

Pneumococcal vaccine

Minimal interval

8 wk

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Dose 1

Dose 2

Maximum interval



ACIP stands for Advisory Committee on Immunization Practices

Pneumococcal vaccine

Minimal interval

Maximum interval

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Dose 1



ACIP
Advisory Committee on Immunization Practices

MMR

Minimal interval

4
wk

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Dose 1

Dose 2


Maximum interval



CDC

ACIP
Advisory Committee on Immunization Practices


Varicella





ACIP
Advisory Committee on Immunization Practices

Hepatitis A





ACIP
Advisory Committee on Immunization Practices

Meningococcal vaccine ACWY

PNS
PORT SAID
NEONATOLOGY SOCIETY

Minimal interval

8 wk

0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 4yr | 6yr | 8yr | 10yr | 12yr | 14yr | 16yr | 18yr

Dose 1

Dose 2

Maximum interval



CDC

ACIP
Advisory Committee on Immunization Practices

Meningococcal vaccine ACWY

PNS
PORT SAID
NEONATOLOGY SOCIETY

Minimal interval

8 wk

0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 4yr | 6yr | 8yr | 10yr | 12yr | 14yr | 16yr | 18yr

Dose 1

Dose 2

Maximum interval



CDC

ACIP
Advisory Committee on Immunization Practices

Meningococcal vaccine ACWY

PNS
PORT SAID
NEONATOLOGY SOCIETY

0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	4yr	6yr	8yr	10yr	12yr	14yr	16yr	18yr

Dose 1

Maximum interval



CDC

ACIP
Advisory Committee on Immunization Practices

Meningococcal vaccine ACWY

Minimal interval

12 wk

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Dose 1

Dose 2

Maximum interval



Meningococcal vaccine ACWY

Minimal interval

12 wk

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Dose 1

Dose 2

Maximum interval



CDC

ACIP
Advisory Committee on Immunization Practices

Meningococcal vaccine ACWY

PNS
PORT SAID
NEONATOLOGY SOCIETY

Minimal interval

Maximum interval

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Dose 1



CDC

ACIP
Advisory Committee on Immunization Practices

Flu vaccine

Minimal interval

4 wk

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Dose 1

Dose 2

Maximum interval



ACIP
Advisory Committee on Immunization Practices

Flu vaccine

Minimal interval

Maximum interval

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

Booster



ACIP stands for Advisory Committee on Immunization Practices

Flu vaccine

Minimal interval

Maximum interval

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 4yr 6yr 8yr 10yr 12yr 14yr 16yr 18yr

1Dose



ACIP
Advisory Committee on Immunization Practices

Reference societies

1. Centre for International Child Health, Department of Paediatrics, Imperial College, London, UK
2. Immunobiology Section, Great Ormond Street Institute of Child Health (ICH), London, UK
3. National Centre for Immunisation Research and Surveillance, Australia,
4. Precision Vaccines Program, Division of Infectious Diseases, Boston Children's Hospital, Boston, MA, USA
5. Harvard Medical School, Boston, MA, USA
6. Centers for Disease Control and Prevention (CDC)
7. Advisory Committee on Immunization Practices


Thank
you
PNS
PORT SAID
NEONATOLOGY SOCIETY
AYMAN
ABU ELHUSSIN
ok.com/AymanAbuElhussin.PH