CATCH UP VACCINES

🌹 Vaccinations:

Birth: HepB
2,4,6 months: DTaP, Rota, Hib, PCV, IPV, HepB
At **6 months** (+) **BCG & OPV**
9 months: MCV4, Measles
12 months: MCV4, MMR, OPV, PCV13
18 months: Varicella, DTaP, HepA, OPV, HiB, MMR
24 months: HepA
4-6 years: MMR, OPV, Varicella, DTaP
(+)
HPV at 11 & 12 years (+Tdap)
MCV4 at 18 years
Annual flu at 6 months

🌹 Catch up vaccine:

⏳ Hepatitis B

Total 4 doses (birth,2,4,6)
Catch-up 3 doses
(1-2): 1 month, (2-3): 2 months
Minimum for the last dose: 24 months

⏳ IPV

Total doses = 3 -> Catch up = 3
(1-2): 1 month, (2-3): 6 months

⏳ DTP

DTaP, Tdap, Td
Total doses = 6 -> Catchup = 4
(1-2): 1 month, (2-3): 1 month
(3-4) 6 months
Note:
After completion of the series, he will need a 2nd booster unless the first one was > 4Y of age.
Td instead of DTP for children >7Y

⏳ MMR, Varicella, HepA

Minimum age (12M)
Total doses: (3,2,2) -> Catchup: 2 doses
(MMR, Varicella, HepA) = 1 month, 3 months, 6 months intervals

⏳ Rota
Total doses 3 doses (2,4,6)
Minimum first dose 15weeks
Minimum last dose 8 months
One month interval between the doses

⏳ HiB
Total doses = 4 -> Catch Up: Depends
(1-2) & (2-3): 1 month, (3-4): 2 months
Child (< 1Y) -> Full series
Child (1-5Y) -> Single Dose
Child (> 5Y) -> No vaccines

⏳ PCV13: Pneumococcal
Total doses = 4 -> Catch up: Depends
< 2Y -> 2 doses (2 months interval)
> 2Y -> Single dose
> 5Y -> No vaccines

⏳ MCV4: Meningococcal
No minimum age
Total doses = 3 -> Catch up = 2 (3 months interval)

🌹 Steps:

1. Write **all vaccines** from birth to pre-school
2. Write all the **missed vaccines** (since the last dose received, if ever)
3. Delete the **repeated** ones
4. Start with **Rota**: if he's older than **8 months** -> No need to give it
5. Then with **BCG**: If he's older than **12 months** -> Given only if he's high risk (NTBCP guidelines) with PPD test
6. No need to give **OPV**
7. Write down all the remaining vaccines, and check the total numbers received (**DTaP= 4 doses**), (**HepB** & **IPV** = **3 doses**), (**MMR**, **varicella**, **HepA**, **MCV4** = **2 doses**)
8. For **PCV13 & HiB**, depends on the **age** of the patient:
   - If he's **older than 5Y** -> **Don't give**
   - **Between (1-5Y) in HiB** or **(2-5Y) in PCV13** -> Give **Single** dose
   - **Younger than 1Y** in **HiB** -> Give **Full series (4)**
   - **Younger than 2Y** in **PCV13** -> Give **(2) doses only**

Hanin