

ADR Identification Basics

Analyze patient case scenarios to identify potential Adverse Drug Reactions (ADRs). Review symptoms, medication history, and timelines to determine whether the reaction is drug-related and identify the suspected drug.

ID: task1

ADR Identification Basics

Task 1 – Pharmacovigilance Case Assessment

Objective: Identify a potential Adverse Drug Reaction (ADR).

Key factors: suspected drug • symptoms • time relationship • dechallenge

Case: 55-year-old patient treated with amoxicillin.

Case Scenario

Patient: 55-year-old

Indication: Bacterial infection

Suspected drug: Amoxicillin

After 2 days of treatment: generalized skin rash + itching

No previous history of similar symptoms

After stopping amoxicillin: symptoms gradually improved

ADR Identification

Suspected ADR: Generalized skin rash and itching

Suspected drug: Amoxicillin

Time relationship: Reaction appeared 2 days after starting treatment

Dechallenge: Symptoms improved after discontinuation

Interpretation: The reaction is potentially drug-related and should be evaluated as a suspected ADR.

Why Is This Considered a Suspected ADR?

1. Temporal relationship – symptoms followed drug exposure.
2. Clinical presentation – rash and itching are compatible with a possible drug reaction.
3. No previous similar symptoms – supports a possible new drug-related event.
4. Positive dechallenge – symptoms improved after the suspected drug was stopped.

Important: These findings support suspicion; they do not by themselves prove causality.

Final Assessment

Drug: Amoxicillin

Event: Generalized skin rash + itching

Onset: 2 days after treatment started

Dechallenge: Improved after stopping the drug

Conclusion: Suspected ADR – requires appropriate evaluation/reporting according to the applicable pharmacovigilance procedure.