

ADR Identification Basics

Task 1 – Pharmacovigilance Case Assessment

Objective: Identify a potential Adverse Drug Reaction (ADR) by reviewing the suspected drug, symptoms, timeline and response after withdrawal.

Case Scenario

Patient	55-year-old
Indication	Bacterial infection
Suspected drug	Amoxicillin
Reaction	Generalized skin rash and itching
Onset	2 days after starting treatment
Previous history	No history of similar symptoms
After discontinuation	Symptoms gradually improved

ADR Identification

- **Suspected ADR:** Generalized skin rash and itching.
- **Suspected drug:** Amoxicillin.
- **Time relationship:** The reaction occurred 2 days after treatment began.
- **Dechallenge:** Symptoms improved after amoxicillin was discontinued.
- **Assessment:** The event is potentially drug-related and should be evaluated as a suspected ADR.

Why does the case support an ADR suspicion?

- **Temporal relationship:** The symptoms followed exposure to the drug.
- **Clinical presentation:** Rash and itching can occur as drug-related reactions.
- **No previous similar symptoms:** This supports a possible new drug-related event.
- **Positive dechallenge:** Improvement after stopping the suspected drug supports the association.

- **Important:** These observations support suspicion but do not independently prove causality.

Final Answer for Submission

Amoxicillin is the suspected drug, and generalized skin rash with itching is the suspected ADR. The reaction occurred 2 days after starting therapy and improved after discontinuation, supporting a possible drug-related association. Therefore, this should be treated as a suspected ADR and evaluated/reported according to the applicable pharmacovigilance procedure.