

CURVED KOSZUL DUALITY FOR ALGEBRAS OVER UNITAL OPERADS

Najib Idrissi

May 2019 @ Higher Algebras in Topology, MPIM Bonn



Université
de Paris



GOAL

Goal

Find resolutions of “algebras”.

Goal

Find resolutions of “algebras”.

Why?

- Compute derived invariants : derived tensor product, derived mapping space...
- Define homotopy algebras over operads.

GOAL

Goal

Find resolutions of “algebras”.

Why?

- Compute derived invariants : derived tensor product, derived mapping space...
- Define homotopy algebras over operads.

Tool of choice: Koszul duality.

Starting data: quadratic algebra $A = F(E)/(R)$, $R \subset E \otimes E$

QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra $A = F(E)/(R)$, $R \subset E \otimes E$

\rightsquigarrow Koszul dual A^i : cofree coalgebra on ΣE with “corelations” $\Sigma^2 R$

QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra $A = F(E)/(R)$, $R \subset E \otimes E$

\rightsquigarrow Koszul dual $A^!$: cofree coalgebra on ΣE with “corelations” $\Sigma^2 R$

(Usually easier to understand $A^! = F(E^*)/(R^\perp)$)

QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra $A = F(E)/(R)$, $R \subset E \otimes E$

\rightsquigarrow Koszul dual $A^!$: cofree coalgebra on ΣE with “corelations” $\Sigma^2 R$

(Usually easier to understand $A^! = F(E^*)/(R^\perp)$)

Examples

1. $A = F(E)$, $R = 0 \implies A^! = 1 \oplus E^*$ with trivial multiplication;

QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra $A = F(E)/(R)$, $R \subset E \otimes E$

\rightsquigarrow Koszul dual $A^!$: cofree coalgebra on ΣE with “corelations” $\Sigma^2 R$

(Usually easier to understand $A^! = F(E^*)/(R^\perp)$)

Examples

1. $A = F(E)$, $R = 0 \implies A^! = 1 \oplus E^*$ with trivial multiplication;
2. $A = S(E) = F(E)/(xy - yx) \implies A^! = F(E^*)/(x^*y^* + y^*x^*) = \Lambda(E^*)$.

QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra $A = F(E)/(R)$, $R \subset E \otimes E$

\rightsquigarrow Koszul dual $A^!$: cofree coalgebra on ΣE with “corelations” $\Sigma^2 R$

(Usually easier to understand $A^! = F(E^*)/(R^\perp)$)

Examples

1. $A = F(E)$, $R = 0 \implies A^! = 1 \oplus E^*$ with trivial multiplication;
2. $A = S(E) = F(E)/(xy - yx) \implies A^! = F(E^*)/(x^*y^* + y^*x^*) = \Lambda(E^*)$.

\implies Koszul complex $K_A := (A \otimes A^i, d_\kappa)$;

QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra $A = F(E)/(R)$, $R \subset E \otimes E$

\rightsquigarrow Koszul dual $A^!$: cofree coalgebra on ΣE with “corelations” $\Sigma^2 R$

(Usually easier to understand $A^! = F(E^*)/(R^\perp)$)

Examples

1. $A = F(E)$, $R = 0 \implies A^! = 1 \oplus E^*$ with trivial multiplication;
2. $A = S(E) = F(E)/(xy - yx) \implies A^! = F(E^*)/(x^*y^* + y^*x^*) = \Lambda(E^*)$.

\implies Koszul complex $K_A := (A \otimes A^i, d_\kappa)$; A is Koszul if K_A is acyclic

Example

$F(E)$ and $S(E)$ are both Koszul.

QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where $BA = (F^c(\Sigma \bar{A}), d_B)$ and $\Omega C = (F(\Sigma^{-1} \bar{C}), d_\Omega)$.

QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where $BA = (F^c(\Sigma\bar{A}), d_B)$ and $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$.

Canonical morphism $\Omega BA \xrightarrow{\sim} A$ is always a cofibrant resolution...

QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where $BA = (F^c(\Sigma\bar{A}), d_B)$ and $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$.

Canonical morphism $\Omega BA \xrightarrow{\sim} A$ is always a cofibrant resolution...but big!

QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where $BA = (F^c(\Sigma\bar{A}), d_B)$ and $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$.

Canonical morphism $\Omega BA \xrightarrow{\sim} A$ is always a cofibrant resolution...but big!

A quadratic $\implies \exists$ canonical morphism $\Omega A^i \rightarrow A$

QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where $BA = (F^c(\Sigma \bar{A}), d_B)$ and $\Omega C = (F(\Sigma^{-1} \bar{C}), d_\Omega)$.

Canonical morphism $\Omega BA \xrightarrow{\sim} A$ is always a cofibrant resolution...but big!

A quadratic $\implies \exists$ canonical morphism $\Omega A^i \rightarrow A$

Theorem (Priddy '70s)

A is Koszul $\iff \Omega A^i \xrightarrow{\sim} A$.

QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where $BA = (F^c(\Sigma\bar{A}), d_B)$ and $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$.

Canonical morphism $\Omega BA \xrightarrow{\sim} A$ is always a cofibrant resolution...but big!

A quadratic $\implies \exists$ canonical morphism $\Omega A^i \rightarrow A$

Theorem (Priddy '70s)

A is Koszul $\iff \Omega A^i \xrightarrow{\sim} A$.

Much smaller resolution!

Examples

$A = F(E) \implies \Omega A^i = A = F(E)$ versus $\Omega BA = F(F^c(F(E)))$

$A = S(E) \implies \Omega A^i = F(\Lambda^c(E))$ versus $\Omega BA = F(F^c(S(E)))$.

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: curved dg-coalgebra

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: curved dg-coalgebra

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: curved dg-coalgebra

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;
- linear $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$ is a coderivation;

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: **curved dg-coalgebra**

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;
- linear $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$ is a coderivation;
- constant $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$ s.t. $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$ and $\theta d = 0$.

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: **curved dg-coalgebra**

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;
- linear $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$ is a coderivation;
- constant $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$ s.t. $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$ and $\theta d = 0$.

Example

$$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y])$$

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: **curved dg**-coalgebra

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;
- linear $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$ is a coderivation;
- constant $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$ s.t. $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$ and $\theta d = 0$.

Example

$$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y]) \rightsquigarrow qA = F(\mathfrak{g})/(xy - yx) = S(\mathfrak{g})$$

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: **curved dg**-coalgebra

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;
- linear $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$ is a coderivation;
- constant $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$ s.t. $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$ and $\theta d = 0$.

Example

$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y]) \rightsquigarrow qA = F(\mathfrak{g})/(xy - yx) = S(\mathfrak{g}) \rightsquigarrow qA^i = S^c(\Sigma\mathfrak{g})$; d_{A^i} = coderivation induced by $d(x \wedge y) = [x, y]$

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: **curved dg-coalgebra**

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;
- linear $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$ is a coderivation;
- constant $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$ s.t. $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$ and $\theta d = 0$.

Example

$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y]) \rightsquigarrow qA = F(\mathfrak{g})/(xy - yx) = S(\mathfrak{g}) \rightsquigarrow qA^i = S^c(\Sigma\mathfrak{g})$; d_{A^i} = coderivation induced by $d(x \wedge y) = [x, y] \rightsquigarrow C_*^{CE}(\mathfrak{g})$

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: **curved dg-coalgebra**

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;
- linear $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$ is a coderivation;
- constant $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$ s.t. $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$ and $\theta d = 0$.

Example

$$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y]) \rightsquigarrow qA = F(\mathfrak{g})/(xy - yx) = S(\mathfrak{g}) \rightsquigarrow qA^i = S^c(\Sigma\mathfrak{g}); d_{A^i} = \text{coderivation induced by } d(x \wedge y) = [x, y] \rightsquigarrow C_*^{CE}(\mathfrak{g})$$

Bar/cobar adjunction: $\text{semi.aug.algebras} \rightleftharpoons \text{curved dg-coalgebras}$.

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: **curved dg-coalgebra**

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;
- linear $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$ is a coderivation;
- constant $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$ s.t. $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$ and $\theta d = 0$.

Example

$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y]) \rightsquigarrow qA = F(\mathfrak{g})/(xy - yx) = S(\mathfrak{g}) \rightsquigarrow qA^i = S^c(\Sigma \mathfrak{g})$; d_{A^i} = coderivation induced by $d(x \wedge y) = [x, y] \rightsquigarrow C_*^{CE}(\mathfrak{g})$

Bar/cobar adjunction: semi.aug.algebras \rightleftarrows curved dg-coalgebras.

Theorem (Polischuck, Positselski)

If qA is Koszul then $\Omega A^i \xrightarrow{\sim} A$ is a cofibrant resolution.

QLC ALGEBRAS – CURVED KD

Quadratic-linear-constant algebra: $A = uF(E)/(R)$ with $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual $A^i = (qA^i, d_{A^i}, \theta_{A^i})$: **curved dg-coalgebra**

- quadratic $\rightsquigarrow qA := F(E)/(qR)$ where $qR := \text{proj}_{E^{\otimes 2}}(R)$;
- linear $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$ is a coderivation;
- constant $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$ s.t. $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$ and $\theta d = 0$.

Example

$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y]) \rightsquigarrow qA = F(\mathfrak{g})/(xy - yx) = S(\mathfrak{g}) \rightsquigarrow qA^i = S^c(\Sigma \mathfrak{g})$; d_{A^i} = coderivation induced by $d(x \wedge y) = [x, y] \rightsquigarrow C_*^{CE}(\mathfrak{g})$

Bar/cobar adjunction: semi.aug.algebras \rightleftarrows curved dg-coalgebras.

Theorem (Polischuck, Positselski)

If qA is Koszul then $\Omega A^i \xrightarrow{\sim} A$ is a cofibrant resolution.


Goal: do this for more general types of algebras (e.g. Poisson algebras). ^{4/15}

What are “more general types of algebras”?

OPERADS

What are “more general types of algebras”?


Operad $\mathbf{P} = \{P(n)\}_{n \geq 0}$: combinatorial object that encodes a type of algebra.



OPERADS

What are “more general types of algebras”?

Operad $\mathbf{P} = \{P(n)\}_{n \geq 0}$: combinatorial object that encodes a type of algebra.




Examples

The “three graces”: **Ass** = associative algebras; **Com** = commutative algebras; **Lie** = Lie algebras.

OPERADS

What are “more general types of algebras”?

Operad $\mathbf{P} = \{P(n)\}_{n \geq 0}$: combinatorial object that encodes a type of algebra.



Examples


The “three graces”: **Ass** = associative algebras; **Com** = commutative algebras; **Lie** = Lie algebras.

E_n = homotopy associative and commutative (for $n \geq 2$) algebras.

OPERADS

What are “more general types of algebras”?

Operad $\mathbf{P} = \{\mathbf{P}(n)\}_{n \geq 0}$: combinatorial object that encodes a type of algebra.



Examples

The “three graces”: **Ass** = associative algebras; **Com** = commutative algebras; **Lie** = Lie algebras.

E_n = homotopy associative and commutative (for $n \geq 2$) algebras.

$e_n := H_*(E_n)$, $n \geq 2$ = Poisson n -algebras.

Quadratic operad: $\mathbf{P} = \mathbf{FOp}(E)/(R)$ where E is a generating set of operations and $R \subset E \circ E$ is a set of quadratic relations.

Quadratic operad: $\mathbf{P} = \mathbf{FOp}(E)/(R)$ where E is a generating set of operations and $R \subset E \circ E$ is a set of quadratic relations.

Examples

$\mathbf{Com} = \mathbf{FOp}(\mu)/(\mu(\mu(x, y), z) = \mu(x, \mu(y, z)))$ is quadratic.

Quadratic operad: $\mathbf{P} = \mathbf{FOp}(E)/(R)$ where E is a generating set of operations and $R \subset E \circ E$ is a set of quadratic relations.

Examples

$\mathbf{Com} = \mathbf{FOp}(\mu)/(\mu(\mu(x, y), z) = \mu(x, \mu(y, z)))$ is quadratic.

Formally similar definitions: Koszul dual cooperad $\mathbf{P}^i = \mathbf{FOp}^c(\Sigma E, \Sigma^2 R)$ and its linear dual $\mathbf{P}^! = \mathbf{FOp}(E^*)/(R^\perp)$.

KD FOR QUADRATIC OPERADS

Quadratic operad: $\mathbf{P} = \mathbf{FOp}(E)/(R)$ where E is a generating set of operations and $R \subset E \circ E$ is a set of quadratic relations.

Examples

$\mathbf{Com} = \mathbf{FOp}(\mu)/(\mu(\mu(x, y), z) = \mu(x, \mu(y, z)))$ is quadratic.

Formally similar definitions: Koszul dual cooperad $\mathbf{P}^! = \mathbf{FOp}^c(\Sigma E, \Sigma^2 R)$ and its linear dual $\mathbf{P}^! = \mathbf{FOp}(E^*)/(R^\perp)$.

Examples

$\mathbf{Ass}^! = \mathbf{Ass}$; $\mathbf{Com}^! = \mathbf{Lie}$, $\mathbf{Lie}^! = \mathbf{Com}$; $e_n^! = e_n\{-n\}$.

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftarrows \{\text{aug.operads}\} : B$$

KOSZUL RESOLUTIONS FOR QUADRATIC OPERADS

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftharpoons \{\text{aug.operads}\} : B$$

Canonical morphism $\Omega B\mathbf{P} \xrightarrow{\sim} \mathbf{P}$ always a resolution, but very big

KOSZUL RESOLUTIONS FOR QUADRATIC OPERADS

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftharpoons \{\text{aug.operads}\} : B$$

Canonical morphism $\Omega BP \xrightarrow{\sim} P$ always a resolution, but very big

Theorem (Ginzburg–Kapranov '94, Getzler–Jones '94, Getzler '95...)

If P is quadratic and Koszul, then $P_\infty := \Omega P^i \xrightarrow{\sim} P$.

KOSZUL RESOLUTIONS FOR QUADRATIC OPERADS

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftharpoons \{\text{aug.operads}\} : B$$

Canonical morphism $\Omega BP \xrightarrow{\sim} P$ always a resolution, but very big

Theorem (Ginzburg–Kapranov '94, Getzler–Jones '94, Getzler '95...)

If P is quadratic and **Koszul**, then $P_\infty := \Omega P^i \xrightarrow{\sim} P$.

In this case, P_∞ -algebras = “homotopy P -algebras”.

KOSZUL RESOLUTIONS FOR QUADRATIC OPERADS

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftarrows \{\text{aug.operads}\} : B$$

Canonical morphism $\Omega BP \xrightarrow{\sim} P$ always a resolution, but very big

Theorem (Ginzburg–Kapranov '94, Getzler–Jones '94, Getzler '95...)

If P is quadratic and **Koszul**, then $P_\infty := \Omega P^i \xrightarrow{\sim} P$.

In this case, P_∞ -algebras = “homotopy P -algebras”.

Examples

$\text{Ass}_\infty = A_\infty$ -algebras, $\text{Com}_\infty = C_\infty$ -algebras, $\text{Lie}_\infty = L_\infty$ -algebras...

$P = \text{FOp}(E)/(R)$ Koszul \rightsquigarrow bar/cobar adjunction:

$$\Omega_\kappa : \{\text{coaug. } P\text{-coalgebras}\} \rightleftarrows \{\text{aug. } P\text{-algebras}\} : B_\kappa$$

$\mathbf{P} = \mathbf{FOp}(E)/(R)$ Koszul \rightsquigarrow bar/cobar adjunction:

$$\Omega_{\kappa} : \{\text{coaug. } \mathbf{P}\text{-coalgebras}\} \rightleftarrows \{\text{aug. } \mathbf{P}\text{-algebras}\} : B_{\kappa}$$

\rightsquigarrow resolution of \mathbf{P} -algebras: $\Omega_{\kappa} B_{\kappa}(-)$, but very big.

$\mathbf{P} = \mathbf{FOp}(E)/(R)$ Koszul \rightsquigarrow bar/cobar adjunction:

$$\Omega_{\kappa} : \{\text{coaug. } \mathbf{P}\text{-coalgebras}\} \rightleftarrows \{\text{aug. } \mathbf{P}\text{-algebras}\} : B_{\kappa}$$

\rightsquigarrow resolution of \mathbf{P} -algebras: $\Omega_{\kappa} B_{\kappa}(-)$, but very big.

Monogenic \mathbf{P} -algebras: $A = \mathbf{P}(V)/(S)$, $S \subset E(V)$

$\mathbf{P} = \mathbf{FOp}(E)/(R)$ Koszul \rightsquigarrow bar/cobar adjunction:

$$\Omega_{\kappa} : \{\text{coaug. } \mathbf{P}\text{-coalgebras}\} \rightleftarrows \{\text{aug. } \mathbf{P}\text{-algebras}\} : B_{\kappa}$$

\rightsquigarrow resolution of \mathbf{P} -algebras: $\Omega_{\kappa} B_{\kappa}(-)$, but very big.

Monogenic \mathbf{P} -algebras: $A = \mathbf{P}(V)/(S)$, $S \subset E(V)$ (\mathbf{P} binary \implies monogenic = quadratic)

KD FOR MONOGENIC ALGEBRAS

$\mathbf{P} = \mathbf{FOp}(E)/(R)$ Koszul \rightsquigarrow bar/cobar adjunction:

$$\Omega_\kappa : \{\text{coaug. } \mathbf{P}\text{-coalgebras}\} \rightleftarrows \{\text{aug. } \mathbf{P}\text{-algebras}\} : B_\kappa$$

\rightsquigarrow resolution of \mathbf{P} -algebras: $\Omega_\kappa B_\kappa(-)$, but very big.

Monogenic \mathbf{P} -algebras: $A = \mathbf{P}(V)/(S)$, $S \subset E(V)$ (\mathbf{P} binary \implies monogenic = quadratic)

Koszul dual:

$$A^i := \mathbf{P}(\Sigma V, \Sigma^2 S), \quad A^! = \mathbf{P}(V^*)/(S^\perp).$$

Theorem (Millès '12)

If \mathbf{P} is quadratic Koszul and if A is a **Koszul** monogenic algebra, then $\Omega_\kappa A^i \xrightarrow{\sim} A$ is a resolution of A .

KD FOR MONOGENIC ALGEBRAS

$\mathbf{P} = \mathbf{FOp}(E)/(R)$ Koszul \rightsquigarrow bar/cobar adjunction:

$$\Omega_\kappa : \{\text{coaug. } \mathbf{P}\text{-coalgebras}\} \rightleftarrows \{\text{aug. } \mathbf{P}\text{-algebras}\} : B_\kappa$$

\rightsquigarrow resolution of \mathbf{P} -algebras: $\Omega_\kappa B_\kappa(-)$, but very big.

Monogenic \mathbf{P} -algebras: $A = \mathbf{P}(V)/(S)$, $S \subset E(V)$ (\mathbf{P} binary \implies monogenic = quadratic)

Koszul dual:

$$A^i := \mathbf{P}(\Sigma V, \Sigma^2 S), \quad A^! = \mathbf{P}(V^*)/(S^\perp).$$

Theorem (Millès '12)

If \mathbf{P} is quadratic Koszul and if A is a **Koszul** monogenic algebra, then $\Omega_\kappa A^i \xrightarrow{\sim} A$ is a resolution of A .

$\mathbf{P} = \mathbf{Ass}$: recovers the classical Koszul duality of associative algebras.

CURVED KD FOR QLC OPERADS

Extension to operads with quadratic-linear-constant relations:

CURVED KD FOR QLC OPERADS

Extension to operads with quadratic-linear-constant relations:

Example

$$u\text{Com} = \text{FOp}(\mu, \mathfrak{i}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(x, \mathfrak{i}) = x)$$

CURVED KD FOR QLC OPERADS

Extension to operads with quadratic-linear-constant relations:

Example

$$u\text{Com} = \text{FOp}(\mu, \mathfrak{!}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(x, \mathfrak{!}) = x)$$

Koszul dual **curved** cooperad: $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A^i}, \theta_{A^i})$

CURVED KD FOR QLC OPERADS

Extension to operads with quadratic-linear-constant relations:

Example

$$u\text{Com} = \text{FOp}(\mu, \mathfrak{!}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(x, \mathfrak{!}) = x)$$

Koszul dual **curved** cooperad: $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A^i}, \theta_{A^i})$

- quadratic $\rightsquigarrow qu\mathbf{P}$: “quadratrization” of $u\mathbf{P}$;

CURVED KD FOR QLC OPERADS

Extension to operads with quadratic-linear-constant relations:

Example

$$u\text{Com} = \text{FOp}(\mu, \mathfrak{!}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(x, \mathfrak{!}) = x)$$

Koszul dual **curved** cooperad: $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A^i}, \theta_{A^i})$

- quadratic $\rightsquigarrow qu\mathbf{P}$: “quadratzation” of $u\mathbf{P}$;
- linear $\rightsquigarrow d_{A^i} : qu\mathbf{P}^i \rightarrow qu\mathbf{P}^i$ coderivation;

CURVED KD FOR QLC OPERADS

Extension to operads with quadratic-linear-constant relations:

Example

$$u\text{Com} = \text{FOp}(\mu, \mathfrak{!}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(x, \mathfrak{!}) = x)$$

Koszul dual **curved** cooperad: $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A_i}, \theta_{A_i})$

- quadratic $\rightsquigarrow qu\mathbf{P}$: “quadratzation” of $u\mathbf{P}$;
- linear $\rightsquigarrow d_{A_i} : qu\mathbf{P}^i \rightarrow qu\mathbf{P}^i$ coderivation;
- constants $\rightsquigarrow \theta_{A_i} : qu\mathbf{P}^i \rightarrow \mathbb{R} \text{ id}$ s.t. $d^2 = (\theta \circ \text{id} \mp \text{id} \circ \theta)\Delta$ and $\theta d = 0$

CURVED KD FOR QLC OPERADS

Extension to operads with quadratic-linear-constant relations:

Example

$$u\text{Com} = \text{FOp}(\mu, \mathfrak{!}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(x, \mathfrak{!}) = x)$$

Koszul dual **curved** cooperad: $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A_i}, \theta_{A_i})$

- quadratic $\rightsquigarrow qu\mathbf{P}$: “quadratzation” of $u\mathbf{P}$;
- linear $\rightsquigarrow d_{A_i} : qu\mathbf{P}^i \rightarrow qu\mathbf{P}^i$ coderivation;
- constants $\rightsquigarrow \theta_{A_i} : qu\mathbf{P}^i \rightarrow \mathbb{R} \text{ id}$ s.t. $d^2 = (\theta \circ \text{id} \mp \text{id} \circ \theta)\Delta$ and $\theta d = 0$

Bar/cobar extends to the curved setting

CURVED KD FOR QLC OPERADS

Extension to operads with quadratic-linear-constant relations:

Example

$$u\text{Com} = \text{FOp}(\mu, \mathfrak{I}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(x, \mathfrak{I}) = x)$$

Koszul dual **curved** cooperad: $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A_i}, \theta_{A_i})$

- quadratic $\rightsquigarrow qu\mathbf{P}$: “quadratzation” of $u\mathbf{P}$;
- linear $\rightsquigarrow d_{A_i} : qu\mathbf{P}^i \rightarrow qu\mathbf{P}^i$ coderivation;
- constants $\rightsquigarrow \theta_{A_i} : qu\mathbf{P}^i \rightarrow \mathbb{R} \text{ id}$ s.t. $d^2 = (\theta \circ \text{id} \mp \text{id} \circ \theta)\Delta$ and $\theta d = 0$

Bar/cobar extends to the curved setting

Theorem (Hirsh–Millès ’12)

If $qu\mathbf{P}$ is Koszul, then $u\mathbf{P}_\infty := \Omega(u\mathbf{P}^i) \xrightarrow{\sim} u\mathbf{P}$: resolution of $u\mathbf{P}$

$uP = \mathbf{FOp}(E \oplus \mathfrak{l})/(R)$: “unital version” of a quadratic operad P

CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \text{FOp}(E \oplus \mathfrak{I})/(R)$: “unital version” of a quadratic operad P

$A = uP(V)/(S)$: algebra w/ QLC relations $S \subset E(V) \oplus V \oplus \mathfrak{I}$

CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \mathbf{FOp}(E \oplus \mathfrak{I})/(R)$: “unital version” of a quadratic operad P

$A = uP(V)/(S)$: algebra w/ QLC relations $S \subset E(V) \oplus V \oplus \mathfrak{I}$

Koszul dual: curved P_i -coalgebra $A^i = (qA^i, d_{A^i}, \theta_{A^i})$

CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \mathbf{FOp}(E \oplus \mathfrak{I})/(R)$: “unital version” of a quadratic operad P

$A = uP(V)/(S)$: algebra w/ QLC relations $S \subset E(V) \oplus V \oplus \mathfrak{I}$

Koszul dual: curved P_i -coalgebra $A^i = (qA^i, d_{A^i}, \theta_{A^i})$

- quadratic $\rightsquigarrow qA = P(V)/(qS)$: “quadratization” of A ;

CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \text{FOp}(E \oplus \mathfrak{I})/(R)$: “unital version” of a quadratic operad P

$A = uP(V)/(S)$: algebra w/ QLC relations $S \subset E(V) \oplus V \oplus \mathfrak{I}$

Koszul dual: curved P_i -coalgebra $A^i = (qA^i, d_{A^i}, \theta_{A^i})$

- quadratic $\rightsquigarrow qA = P(V)/(qS)$: “quadratization” of A ;
- linear $\rightsquigarrow d_{A^i}$: coderivation;

CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \text{FOp}(E \oplus \mathbb{I})/(R)$: “unital version” of a quadratic operad P
 $A = uP(V)/(S)$: algebra w/ QLC relations $S \subset E(V) \oplus V \oplus \mathbb{I}$

Koszul dual: curved P_i -coalgebra $A^i = (qA^i, d_{A^i}, \theta_{A^i})$

- quadratic $\rightsquigarrow qA = P(V)/(qS)$: “quadratization” of A ;
- linear $\rightsquigarrow d_{A^i}$: coderivation;
- constant $\rightsquigarrow \theta : qA^i \rightarrow \mathbb{R}\mathbb{I}$ (+ relations)

CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \text{FOp}(E \oplus \mathbb{I})/(R)$: “unital version” of a quadratic operad P
 $A = uP(V)/(S)$: algebra w/ QLC relations $S \subset E(V) \oplus V \oplus \mathbb{I}$

Koszul dual: curved P^i -coalgebra $A^i = (qA^i, d_{A^i}, \theta_{A^i})$

- quadratic $\rightsquigarrow qA = P(V)/(qS)$: “quadratization” of A ;
- linear $\rightsquigarrow d_{A^i}$: coderivation;
- constant $\rightsquigarrow \theta : qA^i \rightarrow \mathbb{R}\mathbb{I}$ (+ relations)

Generalization of bar/cobar adjunction:

$$\Omega_\kappa : \{\text{curved } P^i\text{-coalgebras}\} \rightleftarrows \{\text{semi.aug. } uP\text{-algebras}\} : B_\kappa$$

CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \text{FOp}(E \oplus \mathbb{I})/(R)$: “unital version” of a quadratic operad P
 $A = uP(V)/(S)$: algebra w/ QLC relations $S \subset E(V) \oplus V \oplus \mathbb{I}$

Koszul dual: curved P_i -coalgebra $A_i = (qA_i, d_{A_i}, \theta_{A_i})$

- quadratic $\rightsquigarrow qA = P(V)/(qS)$: “quadratization” of A ;
- linear $\rightsquigarrow d_{A_i}$: coderivation;
- constant $\rightsquigarrow \theta : qA_i \rightarrow \mathbb{R}\mathbb{I}$ (+ relations)

Generalization of bar/cobar adjunction:

$$\Omega_\kappa : \{\text{curved } P_i\text{-coalgebras}\} \rightleftarrows \{\text{semi.aug. } uP\text{-algebras}\} : B_\kappa$$

Theorem (I. '18)

If qA is Koszul then $\Omega_\kappa A_i \xrightarrow{\sim} A$ is a resolution.

APPLICATION 1: FACTORIZATION HOMOLOGY

M : framed n -manifold, A : uE_n -algebra

APPLICATION 1: FACTORIZATION HOMOLOGY

M : framed n -manifold, A : uE_n -algebra

Goal

Compute $\int_M A = \operatorname{hocolim}_{(D^n) \sqcup k \hookrightarrow M} A^{\otimes k}$.

APPLICATION 1: FACTORIZATION HOMOLOGY

M : framed n -manifold, A : uE_n -algebra

Goal

Compute $\int_M A = \operatorname{hocolim}_{(D^n) \sqcup k \hookrightarrow M} A^{\otimes k}$.

Theorem (Francis 2015)

$\int_M A \simeq E_M \circ_{uE_n}^{\mathbb{L}} A = \operatorname{hocolim}_{\mathbb{L}} (E_M \circ uE_n \circ A \rightrightarrows E_M \circ A)$, where:

$$uE_n(k) = \operatorname{Emb}^{\operatorname{fr}}(\underbrace{\mathbb{R}^n \sqcup \cdots \sqcup \mathbb{R}^n}_{k \times}, \mathbb{R}^n); \quad E_M(k) = \operatorname{Emb}^{\operatorname{fr}}(\underbrace{\mathbb{R}^n \sqcup \cdots \sqcup \mathbb{R}^n}_{k \times}, M).$$

(\exists version for unframed manifolds.)

If we work over \mathbb{R} and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

If we work over \mathbb{R} and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

Theorem (Kontsevich '99; Tamarkin '03 ($n = 2$); Lambrechts–Volić '14; Petersen '14 ($n = 2$); Fresse–Willwacher '15)

The operad $C_*(uE_n)$ is formal: $C_*(uE_n) \simeq u\mathbf{e}_n := H_*(uE_n)$.

CHAINS OF FACTORIZATION HOMOLOGY OVER \mathbb{R}

If we work over \mathbb{R} and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

Theorem (Kontsevich '99; Tamarkin '03 ($n = 2$); Lambrechts–Volić '14; Petersen '14 ($n = 2$); Fresse–Willwacher '15)

The operad $C_*(uE_n)$ is formal: $C_*(uE_n) \simeq u\mathbf{e}_n := H_*(uE_n)$.

Theorem (I.)

M closed, simply connected, smooth, $\dim M \geq 4 \implies$ explicit model of $C_*(E_M)$ as a right $C_*(uE_n)$ -module: Lambrechts–Stanley model \mathbf{LS}_M .

CHAINS OF FACTORIZATION HOMOLOGY OVER \mathbb{R}

If we work over \mathbb{R} and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

Theorem (Kontsevich '99; Tamarkin '03 ($n = 2$); Lambrechts–Volić '14; Petersen '14 ($n = 2$); Fresse–Willwacher '15)

The operad $C_*(uE_n)$ is formal: $C_*(uE_n) \simeq u\mathbf{e}_n := H_*(uE_n)$.

Theorem (I.)

M closed, simply connected, smooth, $\dim M \geq 4 \implies$ explicit model of $C_*(E_M)$ as a right $C_*(uE_n)$ -module: Lambrechts–Stanley model \mathbf{LS}_M .

Upshot: $C_*(\int_M A) \simeq \mathbf{LS}_M \circ_{u\mathbf{e}_n}^{\mathbb{L}} \tilde{A}$

CHAINS OF FACTORIZATION HOMOLOGY OVER \mathbb{R}

If we work over \mathbb{R} and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

Theorem (Kontsevich '99; Tamarkin '03 ($n = 2$); Lambrechts–Volić '14; Petersen '14 ($n = 2$); Fresse–Willwacher '15)

The operad $C_*(uE_n)$ is formal: $C_*(uE_n) \simeq u\mathbf{e}_n := H_*(uE_n)$.

Theorem (I.)

M closed, simply connected, smooth, $\dim M \geq 4 \implies$ explicit model of $C_*(E_M)$ as a right $C_*(uE_n)$ -module: Lambrechts–Stanley model \mathbf{LS}_M .

Upshot: $C_*(\int_M A) \simeq \mathbf{LS}_M \circ_{u\mathbf{e}_n}^{\mathbb{L}} \tilde{A}$

\implies we need to resolve A as a $u\mathbf{e}_n$ -algebra.

WEYL ALGEBRA $\text{Poly}(T^*\mathbb{R}^d[1 - n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

WEYL ALGEBRA $\text{Poly}(T^*\mathbb{R}^d[1 - n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of $H_*(u\mathbf{E}_n)$: free symmetric algebra and $\{x_i, \xi_j\} = \delta_{ij}$

WEYL ALGEBRA $\text{Poly}(T^*\mathbb{R}^d[1 - n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of $H_*(u\mathbf{E}_n)$: free symmetric algebra and $\{x_i, \xi_j\} = \delta_{ij}$

\implies quadratic-(linear-)constant presentation

WEYL ALGEBRA $\text{Poly}(T^*\mathbb{R}^d[1 - n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of $H_*(u\mathbf{E}_n)$: free symmetric algebra and $\{x_i, \xi_j\} = \delta_{ij}$

\implies quadratic-(linear-)constant presentation

Quadratization $qA = S(x_i, \xi_j)$ free symmetric algebra + zero bracket

WEYL ALGEBRA $\text{Poly}(T^*\mathbb{R}^d[1 - n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of $H_*(u\mathbf{E}_n)$: free symmetric algebra and $\{x_i, \xi_j\} = \delta_{ij}$

\implies quadratic-(linear-)constant presentation

Quadratization $qA = S(x_i, \xi_j)$ free symmetric algebra + zero bracket

Koszul dual: $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_j)$ cofree symmetric coalgebra + trivial cobracket

WEYL ALGEBRA $\text{Poly}(T^*\mathbb{R}^d[1 - n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of $H_*(u\mathbf{E}_n)$: free symmetric algebra and $\{x_i, \xi_j\} = \delta_{ij}$

\implies quadratic-(linear-)constant presentation

Quadratization $qA = S(x_i, \xi_j)$ free symmetric algebra + zero bracket

Koszul dual: $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_j)$ cofree symmetric coalgebra + trivial cobracket
- $d = 0$ (no linear terms in the relations)

WEYL ALGEBRA $\text{Poly}(T^*\mathbb{R}^d[1 - n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of $H_*(u\mathbf{E}_n)$: free symmetric algebra and $\{x_i, \xi_j\} = \delta_{ij}$

\implies quadratic-(linear-)constant presentation

Quadratization $qA = S(x_i, \xi_j)$ free symmetric algebra + zero bracket

Koszul dual: $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_j)$ cofree symmetric coalgebra + trivial cobracket
- $d = 0$ (no linear terms in the relations)
- curvature: $\theta(\bar{x}_i \wedge \bar{\xi}_j) = -\delta_{ij}$, zero otherwise.

WEYL ALGEBRA $\text{Poly}(T^*\mathbb{R}^d[1 - n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of $H_*(u\mathbf{E}_n)$: free symmetric algebra and $\{x_i, \xi_j\} = \delta_{ij}$

\implies quadratic-(linear-)constant presentation

Quadratization $qA = S(x_i, \xi_j)$ free symmetric algebra + zero bracket

Koszul dual: $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_j)$ cofree symmetric coalgebra + trivial cobracket
- $d = 0$ (no linear terms in the relations)
- curvature: $\theta(\bar{x}_i \wedge \bar{\xi}_j) = -\delta_{ij}$, zero otherwise.

\implies “small” resolution $Q_A := \Omega_\kappa A^i = (SLS^c(\bar{x}_i, \bar{\xi}_j), d) \xrightarrow{\sim} A$

WEYL ALGEBRA $\text{Poly}(T^*\mathbb{R}^d[1 - n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of $H_*(u\mathbf{E}_n)$: free symmetric algebra and $\{x_i, \xi_j\} = \delta_{ij} \mathbf{1}$

\implies quadratic-(linear-)constant presentation

Quadratization $qA = S(x_i, \xi_j)$ free symmetric algebra + zero bracket

Koszul dual: $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_j)$ cofree symmetric coalgebra + trivial cobracket
- $d = 0$ (no linear terms in the relations)
- curvature: $\theta(\bar{x}_i \wedge \bar{\xi}_j) = -\delta_{ij}$, zero otherwise.

\implies “small” resolution $Q_A := \Omega_\kappa A^i = (SLS^c(\bar{x}_i, \bar{\xi}_j), d) \xrightarrow{\sim} A$

(If we had applied curved KD at the level of operads instead:

$$\Omega_\kappa B_\kappa A \supset (\underbrace{SL}_{\text{cobar}} \underbrace{S^c L^c}_{\text{bar}} \underbrace{S(x_i, \xi_j)}_A, d), + \text{ resolution of the unit...})$$

COMPUTATION OF $\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

We can also compute

$$\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) \simeq \mathbf{LS}_M \circ_{ue_n} (SLS^c(\bar{x}_i, \bar{\xi}_j), d)$$

COMPUTATION OF $\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

We can also compute

$$\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) \simeq \mathbf{LS}_M \circ_{ue_n} (SLS^c(\bar{x}_i, \bar{\xi}_j), d)$$

A bit of homological algebra + explicit description of \mathbf{LS}_M :

Theorem (I. '18, see also Markarian '17, Döppenschmitt '18)

$$\int_M \text{Poly}(T^*\mathbb{R}^d[1-n]) \simeq C_*^{CE}(\Omega^{n-*}(M) \otimes \mathbb{R}\langle 1, x_i, \xi_j \rangle) \simeq \mathbb{R}.$$

COMPUTATION OF $\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

We can also compute

$$\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) \simeq \mathbf{LS}_M \circ_{ue_n} (SLS^c(\bar{x}_i, \bar{\xi}_j), d)$$

A bit of homological algebra + explicit description of \mathbf{LS}_M :

Theorem (I. '18, see also Markarian '17, Döppenschmitt '18)

$$\int_M \text{Poly}(T^*\mathbb{R}^d[1-n]) \simeq C_*^{CE}(\Omega^{n-*}(M) \otimes \mathbb{R}\langle 1, x_i, \xi_j \rangle) \simeq \mathbb{R}.$$

Intuition: quantum observable with values in $A \rightsquigarrow$ “expectation” lives in $\int_M A$, should be a number.

APPLICATION 2: DERIVED ENVELOPING ALGEBRA

Operad \mathbf{P} + \mathbf{P} -algebra $A \implies$ notion of A -modules

APPLICATION 2: DERIVED ENVELOPING ALGEBRA

Operad \mathbf{P} + \mathbf{P} -algebra $A \implies$ notion of A -modules

Examples

$\mathbf{P} = \mathbf{Ass} \rightarrow (A, A)$ bimodules; $\mathbf{P} = \mathbf{Com} \rightarrow A$ -modules; $\mathbf{P} = \mathbf{Lie} \rightarrow$ representations of the Lie algebra.

APPLICATION 2: DERIVED ENVELOPING ALGEBRA

Operad \mathbf{P} + \mathbf{P} -algebra $A \implies$ notion of A -modules

Examples

$\mathbf{P} = \mathbf{Ass} \rightarrow (A, A)$ bimodules; $\mathbf{P} = \mathbf{Com} \rightarrow A$ -modules; $\mathbf{P} = \mathbf{Lie} \rightarrow$ representations of the Lie algebra.

\exists an associative algebra $U_{\mathbf{P}}(A)$ s.t. left $U_{\mathbf{P}}(A)$ -modules = A -modules

APPLICATION 2: DERIVED ENVELOPING ALGEBRA

Operad \mathbf{P} + \mathbf{P} -algebra $A \implies$ notion of A -modules

Examples

$\mathbf{P} = \mathbf{Ass} \rightarrow (A, A)$ bimodules; $\mathbf{P} = \mathbf{Com} \rightarrow A$ -modules; $\mathbf{P} = \mathbf{Lie} \rightarrow$ representations of the Lie algebra.

\exists an associative algebra $U_{\mathbf{P}}(A)$ s.t. left $U_{\mathbf{P}}(A)$ -modules = A -modules

Proposition

For $A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n])$, the derived enveloping algebra $U_{\text{uen}}^{\mathbb{L}}(A)$ is q.iso to the underived one.

APPLICATION 2: DERIVED ENVELOPING ALGEBRA

Operad \mathbf{P} + \mathbf{P} -algebra $A \implies$ notion of **A-modules**

Examples

$\mathbf{P} = \mathbf{Ass} \rightarrow (A, A)$ bimodules; $\mathbf{P} = \mathbf{Com} \rightarrow A$ -modules; $\mathbf{P} = \mathbf{Lie} \rightarrow$ representations of the Lie algebra.

\exists an associative algebra $U_{\mathbf{P}}(A)$ s.t. left $U_{\mathbf{P}}(A)$ -modules = A -modules

Proposition

For $A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n])$, the derived enveloping algebra $U_{\text{ue}_n}^{\mathbb{L}}(A)$ is q.iso to the underived one.

Explicit description: if $A = S(\Sigma^{1-n}\mathfrak{g})$, then

$$U_{\text{ue}_n}(A) = A \otimes U_{\text{cLie}_n}(\Sigma^{1-n}\mathfrak{g}),$$

with $X_f \in U_{\text{cLie}_n}(\Sigma^{1-n}\mathfrak{g})$ for $f \in \mathfrak{g}$ satisfying $X_{\bullet} = 0$, $X_{fg} = fX_g \pm gX_f$,
 $X_fg = \{f, g\} \pm gX_f$, $X_{\{f, g\}} = [X_f, X_g]$, $dX_f = X_{df}$.

THANK YOU FOR YOUR ATTENTION!

These slides: <https://idrissi.eu>