

# CURVED KOSZUL DUALITY FOR ALGEBRAS OVER UNITAL OPERADS

---

Najib Idrissi

May 2019 @ Higher Algebras in Topology, MPIM Bonn



Université  
de Paris



# GOAL

## Goal

Find resolutions of “algebras”.

## Goal

Find resolutions of “algebras”.

Why?

- Compute derived invariants : derived tensor product, derived mapping space...
- Define homotopy algebras over operads.

# GOAL

## Goal

Find resolutions of “algebras”.

Why?

- Compute derived invariants : derived tensor product, derived mapping space...
- Define homotopy algebras over operads.

Tool of choice: Koszul duality.

Starting data: quadratic algebra  $A = F(E)/(R)$ ,  $R \subset E \otimes E$

## QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra  $A = F(E)/(R)$ ,  $R \subset E \otimes E$

$\rightsquigarrow$  Koszul dual  $A^i$ : cofree coalgebra on  $\Sigma E$  with “corelations”  $\Sigma^2 R$

## QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra  $A = F(E)/(R)$ ,  $R \subset E \otimes E$

$\rightsquigarrow$  Koszul dual  $A^!$ : cofree coalgebra on  $\Sigma E$  with “corelations”  $\Sigma^2 R$

(Usually easier to understand  $A^! = F(E^*)/(R^\perp)$ )

## QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra  $A = F(E)/(R)$ ,  $R \subset E \otimes E$

$\rightsquigarrow$  Koszul dual  $A^!$ : cofree coalgebra on  $\Sigma E$  with “corelations”  $\Sigma^2 R$

(Usually easier to understand  $A^! = F(E^*)/(R^\perp)$ )

### Examples

1.  $A = F(E)$ ,  $R = 0 \implies A^! = 1 \oplus E^*$  with trivial multiplication;



# QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra  $A = F(E)/(R)$ ,  $R \subset E \otimes E$

$\rightsquigarrow$  Koszul dual  $A^!$ : cofree coalgebra on  $\Sigma E$  with “corelations”  $\Sigma^2 R$

(Usually easier to understand  $A^! = F(E^*)/(R^\perp)$ )

## Examples

1.  $A = F(E)$ ,  $R = 0 \implies A^! = 1 \oplus E^*$  with trivial multiplication;
2.  $A = S(E) = F(E)/(xy - yx) \implies A^! = F(E^*)/(x^*y^* + y^*x^*) = \Lambda(E^*)$ .

# QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra  $A = F(E)/(R)$ ,  $R \subset E \otimes E$

$\rightsquigarrow$  Koszul dual  $A^!$ : cofree coalgebra on  $\Sigma E$  with “corelations”  $\Sigma^2 R$

(Usually easier to understand  $A^! = F(E^*)/(R^\perp)$ )

## Examples

1.  $A = F(E)$ ,  $R = 0 \implies A^! = 1 \oplus E^*$  with trivial multiplication;
2.  $A = S(E) = F(E)/(xy - yx) \implies A^! = F(E^*)/(x^*y^* + y^*x^*) = \Lambda(E^*)$ .

$\implies$  Koszul complex  $K_A := (A \otimes A^i, d_\kappa(\Sigma e) = e)$ ;

# QUADRATIC ALGEBRAS – KOSZUL DUALS

Starting data: quadratic algebra  $A = F(E)/(R)$ ,  $R \subset E \otimes E$

$\rightsquigarrow$  Koszul dual  $A^!$ : cofree coalgebra on  $\Sigma E$  with “corelations”  $\Sigma^2 R$

(Usually easier to understand  $A^! = F(E^*)/(R^\perp)$ )

## Examples

1.  $A = F(E)$ ,  $R = 0 \implies A^! = 1 \oplus E^*$  with trivial multiplication;
2.  $A = S(E) = F(E)/(xy - yx) \implies A^! = F(E^*)/(x^*y^* + y^*x^*) = \Lambda(E^*)$ .

$\implies$  Koszul complex  $K_A := (A \otimes A^i, d_\kappa(\Sigma e) = e)$ ;  $A$  is Koszul if  $K_A$  is acyclic

## Example

$F(E)$  and  $S(E)$  are both Koszul.

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where  $BA = (F^c(\Sigma\bar{A}), d_B)$  and  $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$ .

## QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where  $BA = (F^c(\Sigma\bar{A}), d_B)$  and  $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$ .

Canonical morphism  $\Omega BA \xrightarrow{\sim} A$  is always a cofibrant resolution...

## QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where  $BA = (F^c(\Sigma\bar{A}), d_B)$  and  $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$ .

Canonical morphism  $\Omega BA \xrightarrow{\sim} A$  is always a cofibrant resolution...but big!

## QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where  $BA = (F^c(\Sigma\bar{A}), d_B)$  and  $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$ .

Canonical morphism  $\Omega BA \xrightarrow{\sim} A$  is always a cofibrant resolution...but big!

$A$  quadratic  $\implies \exists$  canonical morphism  $\Omega A^i \rightarrow A$

# QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where  $BA = (F^c(\Sigma\bar{A}), d_B)$  and  $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$ .

Canonical morphism  $\Omega BA \xrightarrow{\sim} A$  is always a cofibrant resolution...but big!

$A$  quadratic  $\implies \exists$  canonical morphism  $\Omega A^i \rightarrow A$

## Theorem (Priddy '70s)

$A$  is Koszul  $\iff \Omega A^i \xrightarrow{\sim} A$ .



# QUADRATIC ALGEBRAS – KOSZUL RESOLUTIONS

Bar/cobar adjunction:

$$\Omega : \{\text{coaug.coalgebras}\} \rightleftarrows \{\text{aug.algebras}\} : B$$

where  $BA = (F^c(\Sigma\bar{A}), d_B)$  and  $\Omega C = (F(\Sigma^{-1}\bar{C}), d_\Omega)$ .

Canonical morphism  $\Omega BA \xrightarrow{\sim} A$  is always a cofibrant resolution...but big!

$A$  quadratic  $\implies \exists$  canonical morphism  $\Omega A^i \rightarrow A$

## Theorem (Priddy '70s)

$A$  is Koszul  $\iff \Omega A^i \xrightarrow{\sim} A$ .

Much smaller resolution!

## Examples

$A = F(E) \implies \Omega A^i = A = F(E)$  versus  $\Omega BA = FF^c F(E)$

$A = S(E) \implies \Omega A^i = F\Lambda^c(E)$  versus  $\Omega BA = FF^c S(E)$ .

Quadratic-linear-constant algebra:  $A = uF(E)/(R)$  with  $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Quadratic-linear-constant algebra:  $A = uF(E)/(R)$  with  $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$ : curved dg-coalgebra

Quadratic-linear-constant algebra:  $A = uF(E)/(R)$  with  $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$ : curved dg-coalgebra

$$r = \underbrace{r_{(2)}}_{\in qR} + \underbrace{r_{(1)}}_{d(r_{(2)})} + \underbrace{r_{(0)}}_{\theta(r_{(2)})} \quad 1 \in R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1.$$

- quadratic  $\rightsquigarrow qA := F(E)/(qR)$  where  $qR := \text{proj}_{E^{\otimes 2}}(R)$ ;

Quadratic-linear-constant algebra:  $A = uF(E)/(R)$  with  $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$ : curved dg-coalgebra

$$r = \underbrace{r_{(2)}}_{\in qR} + \underbrace{r_{(1)}}_{d(r_{(2)})} + \underbrace{r_{(0)}}_{\theta(r_{(2)})} \quad 1 \in R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1.$$

- quadratic  $\rightsquigarrow qA := F(E)/(qR)$  where  $qR := \text{proj}_{E^{\otimes 2}}(R)$ ;
- linear  $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$  is a coderivation;

Quadratic-linear-constant algebra:  $A = uF(E)/(R)$  with  $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$ : curved dg-coalgebra

$$r = \underbrace{r_{(2)}}_{\in qR} + \underbrace{r_{(1)}}_{d(r_{(2)})} + \underbrace{r_{(0)}}_{\theta(r_{(2)})} \quad 1 \in R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1.$$

- quadratic  $\rightsquigarrow qA := F(E)/(qR)$  where  $qR := \text{proj}_{E^{\otimes 2}}(R)$ ;
- linear  $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$  is a coderivation;
- constant  $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$  s.t.  $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$  and  $\theta d = 0$ .

Quadratic-linear-constant algebra:  $A = uF(E)/(R)$  with  $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$ : curved dg-coalgebra

$$r = \underbrace{r_{(2)}}_{\in qR} + \underbrace{r_{(1)}}_{d(r_{(2)})} + \underbrace{r_{(0)}}_{\theta(r_{(2)})} \quad 1 \in R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1.$$

- quadratic  $\rightsquigarrow qA := F(E)/(qR)$  where  $qR := \text{proj}_{E^{\otimes 2}}(R)$ ;
- linear  $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$  is a coderivation;
- constant  $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$  s.t.  $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$  and  $\theta d = 0$ .

## Example

$$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y])$$

Quadratic-linear-constant algebra:  $A = uF(E)/(R)$  with  $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$ : curved dg-coalgebra

$$r = \underbrace{r_{(2)}}_{\in qR} + \underbrace{r_{(1)}}_{d(r_{(2)})} + \underbrace{r_{(0)}}_{\theta(r_{(2)})} \quad 1 \in R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1.$$

- quadratic  $\rightsquigarrow qA := F(E)/(qR)$  where  $qR := \text{proj}_{E^{\otimes 2}}(R)$ ;
- linear  $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$  is a coderivation;
- constant  $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$  s.t.  $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$  and  $\theta d = 0$ .

## Example

$$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y]) \rightsquigarrow qA = F(\mathfrak{g})/(xy - yx) = S(\mathfrak{g})$$



**Quadratic-linear-constant** algebra:  $A = uF(E)/(R)$  with  $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$ : **curved dg-coalgebra**

$$r = \underbrace{r_{(2)}}_{\in qR} + \underbrace{r_{(1)}}_{d(r_{(2)})} + \underbrace{r_{(0)}}_{\theta(r_{(2)})} \quad 1 \in R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1.$$

- quadratic  $\rightsquigarrow qA := F(E)/(qR)$  where  $qR := \text{proj}_{E^{\otimes 2}}(R)$ ;
- linear  $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$  is a coderivation;
- constant  $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$  s.t.  $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$  and  $\theta d = 0$ .

## Example

$$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y]) \rightsquigarrow qA = F(\mathfrak{g})/(xy - yx) = S(\mathfrak{g})$$

$d_{A^i}$  = coderivation induced by  $d(x \wedge y) = [x, y]$

Quadratic-linear-constant algebra:  $A = uF(E)/(R)$  with  $R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1$

Koszul dual  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$ : curved dg-coalgebra

$$r = \underbrace{r_{(2)}}_{\in qR} + \underbrace{r_{(1)}}_{d(r_{(2)})} + \underbrace{r_{(0)}}_{\theta(r_{(2)})} \quad 1 \in R \subset E^{\otimes 2} \oplus E \oplus \mathbb{R}1.$$

- quadratic  $\rightsquigarrow qA := F(E)/(qR)$  where  $qR := \text{proj}_{E^{\otimes 2}}(R)$ ;
- linear  $\rightsquigarrow d_{A^i} : qA^i \rightarrow qA^i$  is a coderivation;
- constant  $\rightsquigarrow \theta_{A^i} : qA^i \rightarrow \mathbb{R}$  s.t.  $d^2 = (\theta \otimes \text{id} \mp \text{id} \otimes \theta)\Delta$  and  $\theta d = 0$ .

## Example

$$A = U(\mathfrak{g}) = uF(\mathfrak{g})/(xy - yx - [x, y]) \rightsquigarrow qA = F(\mathfrak{g})/(xy - yx) = S(\mathfrak{g})$$

$$d_{A^i} = \text{coderivation induced by } d(x \wedge y) = [x, y] \rightsquigarrow A^i = C_*^{CE}(\mathfrak{g})$$

Bar/cobar adjunction:

$$\Omega : \{\text{curved dg-coalgebras}\} \rightleftarrows \{\text{semi.aug.algebras}\} : B$$

where  $BA = (F^c(\Sigma\bar{A}), d_2 + d_1, \theta)$  and  $\Omega(C) = (uF(\Sigma^{-1}C), d_2 + d_1 + d_0)$ .

Bar/cobar adjunction:

$$\Omega : \{\text{curved dg-coalgebras}\} \rightleftarrows \{\text{semi.aug.algebras}\} : B$$

where  $BA = (F^c(\Sigma \bar{A}), d_2 + d_1, \theta)$  and  $\Omega(C) = (uF(\Sigma^{-1}C), d_2 + d_1 + d_0)$ .

## Theorem (Polischuck, Positselski)

If  $qA$  is Koszul then  $\Omega A^i \xrightarrow{\sim} A$  is a cofibrant resolution.

Bar/cobar adjunction:

$$\Omega : \{\text{curved dg-coalgebras}\} \rightleftarrows \{\text{semi.aug.algebras}\} : B$$

where  $BA = (F^c(\Sigma\bar{A}), d_2 + d_1, \theta)$  and  $\Omega(C) = (uF(\Sigma^{-1}C), d_2 + d_1 + d_0)$ .

## Theorem (Polischuck, Positselski)

If  $qA$  is Koszul then  $\Omega A^i \xrightarrow{\sim} A$  is a cofibrant resolution.


Goal: do this for more general types of unital algebras.

What are “more general types of algebras”?

# OPERADS

What are “more general types of algebras”?


**Operad**  $\mathbf{P} = \{P(n)\}_{n \geq 0}$ : combinatorial object that encodes a type of algebra.



# OPERADS

What are “more general types of algebras”?

**Operad**  $\mathbf{P} = \{P(n)\}_{n \geq 0}$ : combinatorial object that encodes a type of algebra.



## Examples


The “three graces”: **Ass** = associative algebras; **Com** = commutative algebras; **Lie** = Lie algebras.



# OPERADS

What are “more general types of algebras”?

**Operad**  $\mathbf{P} = \{P(n)\}_{n \geq 0}$ : combinatorial object that encodes a type of algebra.



## Examples


The “three graces”: **Ass** = associative algebras; **Com** = commutative algebras; **Lie** = Lie algebras.

$E_n$  = homotopy associative and commutative (for  $n \geq 2$ ) algebras.

# OPERADS

What are “more general types of algebras”?

**Operad**  $\mathbf{P} = \{\mathbf{P}(n)\}_{n \geq 0}$ : combinatorial object that encodes a type of algebra.



## Examples

The “three graces”: **Ass** = associative algebras; **Com** = commutative algebras; **Lie** = Lie algebras.

$E_n$  = homotopy associative and commutative (for  $n \geq 2$ ) algebras.

$e_n := H_*(E_n)$ ,  $n \geq 2$  = Poisson  $n$ -algebras.

**Quadratic operad:**  $\mathbf{P} = \mathbf{FOp}(E)/(R)$  where  $E$  is a generating set of operations and  $R \subset E \circ E$  is a set of quadratic relations.

**Quadratic operad:**  $\mathbf{P} = \mathbf{FOp}(E)/(R)$  where  $E$  is a generating set of operations and  $R \subset E \circ E$  is a set of quadratic relations.

### Examples

$\mathbf{Com} = \mathbf{FOp}(\mu)/(\mu(\mu(x, y), z) = \mu(x, \mu(y, z)))$  is quadratic.

**Quadratic operad:**  $\mathbf{P} = \mathbf{FOP}(E)/(R)$  where  $E$  is a generating set of operations and  $R \subset E \circ E$  is a set of quadratic relations.

### Examples

$\mathbf{Com} = \mathbf{FOP}(\mu)/(\mu(\mu(x, y), z) = \mu(x, \mu(y, z)))$  is quadratic.

Formally similar definitions: Koszul dual cooperad  $\mathbf{P}^i = \mathbf{FOP}^c(\Sigma E, \Sigma^2 R)$  and its linear dual  $\mathbf{P}^! = \mathbf{FOP}(E^*)/(R^\perp)$ .

## KD FOR QUADRATIC OPERADS

**Quadratic operad:**  $\mathbf{P} = \mathbf{FOP}(E)/(R)$  where  $E$  is a generating set of operations and  $R \subset E \circ E$  is a set of quadratic relations.

### Examples

$\mathbf{Com} = \mathbf{FOP}(\mu)/(\mu(\mu(x, y), z) = \mu(x, \mu(y, z)))$  is quadratic.

Formally similar definitions: Koszul dual cooperad  $\mathbf{P}^! = \mathbf{FOP}^c(\Sigma E, \Sigma^2 R)$  and its linear dual  $\mathbf{P}^! = \mathbf{FOP}(E^*)/(R^\perp)$ .

### Examples

$\mathbf{Ass}^! = \mathbf{Ass}$ ;  $\mathbf{Com}^! = \mathbf{Lie}$ ,  $\mathbf{Lie}^! = \mathbf{Com}$ ;  $e_n^! = e_n\{-n\}$ .

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftarrows \{\text{aug.operads}\} : B$$

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftarrows \{\text{aug.operads}\} : B$$

Canonical morphism  $\Omega B\mathbf{P} \xrightarrow{\sim} \mathbf{P}$  always a resolution, but very big



# KOSZUL RESOLUTIONS FOR QUADRATIC OPERADS

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftharpoons \{\text{aug.operads}\} : B$$

Canonical morphism  $\Omega BP \xrightarrow{\sim} P$  always a resolution, but very big

**Theorem (Ginzburg–Kapranov '94, Getzler–Jones '94, Getzler '95...)**

If  $P$  is quadratic and Koszul, then  $P_\infty := \Omega P^i \xrightarrow{\sim} P$ .

# KOSZUL RESOLUTIONS FOR QUADRATIC OPERADS

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftharpoons \{\text{aug.operads}\} : B$$

Canonical morphism  $\Omega BP \xrightarrow{\sim} P$  always a resolution, but very big

**Theorem (Ginzburg–Kapranov '94, Getzler–Jones '94, Getzler '95...)**

If  $P$  is quadratic and **Koszul**, then  $P_\infty := \Omega P^i \xrightarrow{\sim} P$ .

In this case,  $P_\infty$ -algebras = “homotopy  $P$ -algebras”.

# KOSZUL RESOLUTIONS FOR QUADRATIC OPERADS

Formally similar definitions: bar/cobar adjunction

$$\Omega : \{\text{coaug.cooperads}\} \rightleftarrows \{\text{aug.operads}\} : B$$

Canonical morphism  $\Omega BP \xrightarrow{\sim} P$  always a resolution, but very big

**Theorem (Ginzburg–Kapranov '94, Getzler–Jones '94, Getzler '95...)**

If  $P$  is quadratic and **Koszul**, then  $P_\infty := \Omega P^i \xrightarrow{\sim} P$ .

In this case,  $P_\infty$ -algebras = “homotopy  $P$ -algebras”.

## Examples

$\text{Ass}_\infty = A_\infty$ -algebras,  $\text{Com}_\infty = C_\infty$ -algebras,  $\text{Lie}_\infty = L_\infty$ -algebras...

$\mathbf{P} = \mathbf{FOp}(E)/(R)$  Koszul  $\rightsquigarrow$  bar/cobar adjunction:

$$\Omega_{\kappa} : \{\text{coaug. } \mathbf{Pi}\text{-coalgebras}\} \rightleftharpoons \{\text{aug. } \mathbf{P}\text{-algebras}\} : B_{\kappa},$$

where  $\Omega_{\kappa}C = (\mathbf{P}(\Sigma^{-1}\bar{C}), d)$  and  $B_{\kappa}A = (\mathbf{Pi}(\Sigma\bar{A}), d)$ .

$\mathbf{P} = \mathbf{FOp}(E)/(R)$  Koszul  $\rightsquigarrow$  bar/cobar adjunction:

$$\Omega_\kappa : \{\text{coaug. } \mathbf{Pi}\text{-coalgebras}\} \rightleftarrows \{\text{aug. } \mathbf{P}\text{-algebras}\} : B_\kappa,$$

where  $\Omega_\kappa C = (\mathbf{P}(\Sigma^{-1}\bar{C}), d)$  and  $B_\kappa A = (\mathbf{Pi}(\Sigma\bar{A}), d)$ .

$\rightsquigarrow$  resolution of  $\mathbf{P}$ -algebras:  $\Omega_\kappa B_\kappa(-)$ , but very big.

# BIG RESOLUTION OF OPERADIC ALGEBRAS

$\mathbf{P} = \mathbf{FOp}(E)/(R)$  Koszul  $\rightsquigarrow$  bar/cobar adjunction:

$$\Omega_{\kappa} : \{\text{coaug. } \mathbf{Pi}\text{-coalgebras}\} \rightleftarrows \{\text{aug. } \mathbf{P}\text{-algebras}\} : B_{\kappa},$$

where  $\Omega_{\kappa}C = (\mathbf{P}(\Sigma^{-1}\bar{C}), d)$  and  $B_{\kappa}A = (\mathbf{Pi}(\Sigma\bar{A}), d)$ .

$\rightsquigarrow$  resolution of  $\mathbf{P}$ -algebras:  $\Omega_{\kappa}B_{\kappa}(-)$ , but very big.

## Example

For a Lie algebra  $\mathfrak{g}$ ,  $\Omega_{\kappa}B_{\kappa}\mathfrak{g} = (L(C_{*-1}^{CE}(\mathfrak{g})), d)$ .

Recall  $\mathbf{P} = \mathbf{FOp}(E)/(R)$ .

Monogenic  $\mathbf{P}$ -algebras:  $A = \mathbf{P}(V)/(S)$ ,  $S \subset E(V)$ .

Recall  $\mathbf{P} = \mathbf{F}\mathbf{Op}(E)/(R)$ .

Monogenic  $\mathbf{P}$ -algebras:  $A = \mathbf{P}(V)/(S)$ ,  $S \subset E(V)$ .

(Monogenic = quadratic for binary  $\mathbf{P}$ )



## KD FOR MONOGENIC OPERADIC ALGEBRAS

Recall  $\mathbf{P} = \mathbf{FOp}(E)/(R)$ .

Monogenic  $\mathbf{P}$ -algebras:  $A = \mathbf{P}(V)/(S)$ ,  $S \subset E(V)$ .

(Monogenic = quadratic for binary  $\mathbf{P}$ )

Koszul dual:  $A^! := \mathbf{P}^i(\Sigma V, \Sigma^2 S)$ ,  $A^! = \mathbf{P}(V^*)/(S^\perp)$ .

Recall  $\mathbf{P} = \mathbf{FOp}(E)/(R)$ .

Monogenic  $\mathbf{P}$ -algebras:  $A = \mathbf{P}(V)/(S)$ ,  $S \subset E(V)$ .

(Monogenic = quadratic for binary  $\mathbf{P}$ )

Koszul dual:  $A^i := \mathbf{P}^i(\Sigma V, \Sigma^2 S)$ ,  $A^! = \mathbf{P}(V^*)/(S^\perp)$ .

Koszul complex:  $K_A = (A \otimes A^i, d_\kappa(\Sigma v) = v)$ .

# KD FOR MONOGENIC OPERADIC ALGEBRAS

Recall  $\mathbf{P} = \mathbf{FOp}(E)/(R)$ .

Monogenic  $\mathbf{P}$ -algebras:  $A = \mathbf{P}(V)/(S)$ ,  $S \subset E(V)$ .

(Monogenic = quadratic for binary  $\mathbf{P}$ )

Koszul dual:  $A^! := \mathbf{P}^!(\Sigma V, \Sigma^2 S)$ ,  $A^! = \mathbf{P}(V^*)/(S^\perp)$ .

Koszul complex:  $K_A = (A \otimes A^i, d_\kappa(\Sigma v) = v)$ .

## Theorem (Millès '12)

If  $\mathbf{P}$  is quadratic Koszul and if  $A$  is a **Koszul** monogenic algebra, then  $\Omega_\kappa A^i \xrightarrow{\sim} A$  is a resolution of  $A$ .

# KD FOR MONOGENIC OPERADIC ALGEBRAS

Recall  $\mathbf{P} = \mathbf{FOp}(E)/(R)$ .

Monogenic  $\mathbf{P}$ -algebras:  $A = \mathbf{P}(V)/(S)$ ,  $S \subset E(V)$ .

(Monogenic = quadratic for binary  $\mathbf{P}$ )

Koszul dual:  $A^! := \mathbf{P}^!(\Sigma V, \Sigma^2 S)$ ,  $A^! = \mathbf{P}(V^*)/(S^\perp)$ .

Koszul complex:  $K_A = (A \otimes A^i, d_\kappa(\Sigma v) = v)$ .

## Theorem (Millès '12)

If  $\mathbf{P}$  is quadratic Koszul and if  $A$  is a **Koszul** monogenic algebra, then  $\Omega_\kappa A^i \xrightarrow{\sim} A$  is a resolution of  $A$ .

## Example

$\mathbf{P} = \mathbf{Ass}$ : recovers the classical Koszul duality of associative algebras.

## CURVED KD FOR QLC OPERADS

Operads with QLC relations  $u\mathbf{P} = \mathbf{FOp}(E)/(R)$ ,  $R \subset E \circ E \oplus E \oplus \mathbb{R} \text{id}$

## CURVED KD FOR QLC OPERADS

Operads with QLC relations  $u\mathbf{P} = \mathbf{FOp}(E)/(R)$ ,  $R \subset E \circ E \oplus E \oplus \mathbb{R} \text{id}$

Koszul dual **curved** cooperad:  $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A_i}, \theta_{A_i})$

## CURVED KD FOR QLC OPERADS

Operads with QLC relations  $u\mathbf{P} = \mathbf{FOp}(E)/(R)$ ,  $R \subset E \circ E \oplus E \oplus \mathbb{R} \text{id}$

Koszul dual **curved** cooperad:  $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A_i}, \theta_{A_i})$

- quadratic  $\rightsquigarrow qu\mathbf{P}$ : “quadratrization” of  $u\mathbf{P}$ ;

## CURVED KD FOR QLC OPERADS

Operads with QLC relations  $u\mathbf{P} = \mathbf{FOp}(E)/(R)$ ,  $R \subset E \circ E \oplus E \oplus \mathbb{R} \text{id}$

Koszul dual **curved** cooperad:  $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A_i}, \theta_{A_i})$

- quadratic  $\rightsquigarrow qu\mathbf{P}$ : “quadratzation” of  $u\mathbf{P}$ ;
- linear  $\rightsquigarrow d_{A_i} : qu\mathbf{P}^i \rightarrow qu\mathbf{P}^i$  coderivation;



## CURVED KD FOR QLC OPERADS

Operads with QLC relations  $u\mathbf{P} = \mathbf{FOp}(E)/(R)$ ,  $R \subset E \circ E \oplus E \oplus \mathbb{R} \text{id}$

Koszul dual **curved** cooperad:  $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A_i}, \theta_{A_i})$

- quadratic  $\rightsquigarrow qu\mathbf{P}$ : “quadratrization” of  $u\mathbf{P}$ ;
- linear  $\rightsquigarrow d_{A_i} : qu\mathbf{P}^i \rightarrow qu\mathbf{P}^i$  coderivation;
- constants  $\rightsquigarrow \theta_{A_i} : qu\mathbf{P}^i \rightarrow \mathbb{R} \text{id}$  s.t.  $d^2 = (\theta \circ \text{id} \mp \text{id} \circ \theta)\Delta$  and  $\theta d = 0$

## CURVED KD FOR QLC OPERADS

Operads with QLC relations  $u\mathbf{P} = \mathbf{FOp}(E)/(R)$ ,  $R \subset E \circ E \oplus E \oplus \mathbb{R} \text{id}$

Koszul dual **curved** cooperad:  $u\mathbf{P}^i = (qu\mathbf{P}^i, d_{A_i}, \theta_{A_i})$

- quadratic  $\rightsquigarrow qu\mathbf{P}$ : “quadratzation” of  $u\mathbf{P}$ ;
- linear  $\rightsquigarrow d_{A_i} : qu\mathbf{P}^i \rightarrow qu\mathbf{P}^i$  coderivation;
- constants  $\rightsquigarrow \theta_{A_i} : qu\mathbf{P}^i \rightarrow \mathbb{R} \text{id}$  s.t.  $d^2 = (\theta \circ \text{id} \mp \text{id} \circ \theta)\Delta$  and  $\theta d = 0$

### Example

$$u\mathbf{Com} = \mathbf{FOp}(\mu, \mathfrak{!}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(\mathfrak{!}, x) = x)$$

## CURVED KD FOR QLC OPERADS

Operads with QLC relations  $uP = \mathbf{FOp}(E)/(R)$ ,  $R \subset E \circ E \oplus E \oplus \mathbb{R} \text{id}$

Koszul dual **curved** cooperad:  $uP^i = (quP^i, d_{Ai}, \theta_{Ai})$

- quadratic  $\rightsquigarrow quP$ : “quadratzation” of  $uP$ ;
- linear  $\rightsquigarrow d_{Ai} : quP^i \rightarrow quP^i$  coderivation;
- constants  $\rightsquigarrow \theta_{Ai} : quP^i \rightarrow \mathbb{R} \text{id}$  s.t.  $d^2 = (\theta \circ \text{id} \mp \text{id} \circ \theta)\Delta$  and  $\theta d = 0$

### Example

$$u\text{Com} = \mathbf{FOp}(\mu, \mathfrak{!}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(\mathfrak{!}, x) = x)$$

$$u\text{Com}^i = (\text{Com}^i \oplus \mathfrak{!}^c, d = 0, \theta(\mu^c \circ_1 \mathfrak{!}^c) = -1)$$

## CURVED KD FOR QLC OPERADS

Operads with QLC relations  $uP = \mathbf{FOp}(E)/(R)$ ,  $R \subset E \circ E \oplus E \oplus \mathbb{R} \text{id}$

Koszul dual **curved** cooperad:  $uP^i = (quP^i, d_{A_i}, \theta_{A_i})$

- quadratic  $\rightsquigarrow quP$ : “quadratzation” of  $uP$ ;
- linear  $\rightsquigarrow d_{A_i} : quP^i \rightarrow quP^i$  coderivation;
- constants  $\rightsquigarrow \theta_{A_i} : quP^i \rightarrow \mathbb{R} \text{id}$  s.t.  $d^2 = (\theta \circ \text{id} \mp \text{id} \circ \theta)\Delta$  and  $\theta d = 0$

### Example

$$u\text{Com} = \mathbf{FOp}(\mu, \mathfrak{!}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(\mathfrak{!}, x) = x)$$

$$u\text{Com}^i = (\text{Com}^i \oplus \mathfrak{!}^c, d = 0, \theta(\mu^c \circ_1 \mathfrak{!}^c) = -1)$$

Bar/cobar extends to the curved setting

## CURVED KD FOR QLC OPERADS

Operads with QLC relations  $uP = \mathbf{FOp}(E)/(R)$ ,  $R \subset E \circ E \oplus E \oplus \mathbb{R} \text{id}$

Koszul dual **curved** cooperad:  $uP^i = (quP^i, d_{A_i}, \theta_{A_i})$

- quadratic  $\rightsquigarrow quP$ : “quadratzation” of  $uP$ ;
- linear  $\rightsquigarrow d_{A_i} : quP^i \rightarrow quP^i$  coderivation;
- constants  $\rightsquigarrow \theta_{A_i} : quP^i \rightarrow \mathbb{R} \text{id}$  s.t.  $d^2 = (\theta \circ \text{id} \mp \text{id} \circ \theta)\Delta$  and  $\theta d = 0$

### Example

$$u\text{Com} = \mathbf{FOp}(\mu, \mathfrak{!}) / (\mu(\mu(x, y), z) = \mu(x, \mu(y, z)), \mu(\mathfrak{!}, x) = x)$$

$$u\text{Com}^i = (\text{Com}^i \oplus \mathfrak{!}^c, d = 0, \theta(\mu^c \circ_1 \mathfrak{!}^c) = -1)$$

Bar/cobar extends to the curved setting

### Theorem (Hirsh–Millès ’12)

If  $quP$  is Koszul, then  $uP_\infty := \Omega(uP^i) \xrightarrow{\sim} uP$ : resolution of  $uP$

$uP = \mathbf{FOp}(E \oplus \mathfrak{I})/(R + R')$ : unital version of quadratic  $P = \mathbf{FOp}(E)/(R)$

## CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \mathbf{FOp}(E \oplus \mathfrak{I})/(R + R')$ : unital version of quadratic  $P = \mathbf{FOp}(E)/(R)$

$A = uP(V)/(S)$ : algebra w/ QLC relations  $S \subset E(V) \oplus V \oplus \mathfrak{I}$

## CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \mathbf{FOp}(E \oplus \mathfrak{I})/(R + R')$ : unital version of quadratic  $P = \mathbf{FOp}(E)/(R)$

$A = uP(V)/(S)$ : algebra w/ QLC relations  $S \subset E(V) \oplus V \oplus \mathfrak{I}$

**Koszul dual**: curved  $P_i$ -coalgebra  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$



## CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \text{FOp}(E \oplus \mathfrak{I})/(R + R')$ : unital version of quadratic  $P = \text{FOp}(E)/(R)$

$A = uP(V)/(S)$ : algebra w/ QLC relations  $S \subset E(V) \oplus V \oplus \mathfrak{I}$

**Koszul dual**: curved  $P_i$ -coalgebra  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$

- quadratic  $\rightsquigarrow qA = P(V)/(qS)$ : “quadratization” of  $A$ ;

## CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \text{FOp}(E \oplus \mathfrak{I})/(R + R')$ : unital version of quadratic  $P = \text{FOp}(E)/(R)$

$A = uP(V)/(S)$ : algebra w/ QLC relations  $S \subset E(V) \oplus V \oplus \mathfrak{I}$

**Koszul dual**: curved  $P_i$ -coalgebra  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$

- quadratic  $\rightsquigarrow qA = P(V)/(qS)$ : “quadratization” of  $A$ ;
- linear  $\rightsquigarrow d_{A^i}$ : coderivation;

## CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \text{FOp}(E \oplus \mathbb{I})/(R + R')$ : unital version of quadratic  $P = \text{FOp}(E)/(R)$

$A = uP(V)/(S)$ : algebra w/ QLC relations  $S \subset E(V) \oplus V \oplus \mathbb{I}$

**Koszul dual**: curved  $P_i$ -coalgebra  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$

- quadratic  $\rightsquigarrow qA = P(V)/(qS)$ : “quadratization” of  $A$ ;
- linear  $\rightsquigarrow d_{A^i}$ : coderivation;
- constant  $\rightsquigarrow \theta : qA^i \rightarrow \mathbb{R}\mathbb{I}$  (+ relations)

## CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \mathbf{FOp}(E \oplus \mathbb{I})/(R + R')$ : unital version of quadratic  $P = \mathbf{FOp}(E)/(R)$

$A = uP(V)/(S)$ : algebra w/ QLC relations  $S \subset E(V) \oplus V \oplus \mathbb{I}$

**Koszul dual**: curved  $P^i$ -coalgebra  $A^i = (qA^i, d_{A^i}, \theta_{A^i})$

- quadratic  $\rightsquigarrow qA = P(V)/(qS)$ : “quadratization” of  $A$ ;
- linear  $\rightsquigarrow d_{A^i}$ : coderivation;
- constant  $\rightsquigarrow \theta : qA^i \rightarrow \mathbb{R}\mathbb{I}$  (+ relations)

Generalization of bar/cobar adjunction:

$$\Omega_\kappa : \{\text{curved } P^i\text{-coalgebras}\} \rightleftarrows \{\text{semi.aug. } uP\text{-algebras}\} : B_\kappa$$

# CURVED KD FOR ALGEBRAS OVER BINARY UNITAL OPERADS

$uP = \mathbf{FOp}(E \oplus \mathbb{I})/(R + R')$ : unital version of quadratic  $P = \mathbf{FOp}(E)/(R)$   
 $A = uP(V)/(S)$ : algebra w/ QLC relations  $S \subset E(V) \oplus V \oplus \mathbb{I}$

**Koszul dual**: curved  $P_i$ -coalgebra  $A_i = (qA_i, d_{A_i}, \theta_{A_i})$

- quadratic  $\rightsquigarrow qA = P(V)/(qS)$ : “quadratization” of  $A$ ;
- linear  $\rightsquigarrow d_{A_i}$ : coderivation;
- constant  $\rightsquigarrow \theta : qA_i \rightarrow \mathbb{R}\mathbb{I}$  (+ relations)

Generalization of bar/cobar adjunction:

$$\Omega_\kappa : \{\text{curved } P_i\text{-coalgebras}\} \rightleftarrows \{\text{semi.aug. } uP\text{-algebras}\} : B_\kappa$$

## Theorem (I. '18)

If  $qA$  is Koszul then  $\Omega_\kappa A_i \xrightarrow{\sim} A$  is a resolution.

## APPLICATION 1: FACTORIZATION HOMOLOGY

$M$ : framed  $n$ -manifold,  $A$ :  $uE_n$ -algebra

## APPLICATION 1: FACTORIZATION HOMOLOGY

$M$ : framed  $n$ -manifold,  $A$ :  $uE_n$ -algebra

### Goal

Compute  $\int_M A = \operatorname{hocolim}_{(D^n) \sqcup k \hookrightarrow M} A^{\otimes k}$ .

# APPLICATION 1: FACTORIZATION HOMOLOGY

$M$ : framed  $n$ -manifold,  $A$ :  $uE_n$ -algebra

## Goal

Compute  $\int_M A = \operatorname{hocolim}_{(D^n) \sqcup k \hookrightarrow M} A^{\otimes k}$ .

## Theorem (Francis 2015)

$\int_M A \simeq E_M \circ_{uE_n}^{\mathbb{L}} A = \operatorname{hocolim}_{\mathbb{L}} (E_M \circ uE_n \circ A \rightrightarrows E_M \circ A)$ , where:

$$uE_n(k) = \operatorname{Emb}^{\operatorname{fr}}(\underbrace{\mathbb{R}^n \sqcup \cdots \sqcup \mathbb{R}^n}_{k \times}, \mathbb{R}^n); \quad E_M(k) = \operatorname{Emb}^{\operatorname{fr}}(\underbrace{\mathbb{R}^n \sqcup \cdots \sqcup \mathbb{R}^n}_{k \times}, M).$$

( $\exists$  version for unframed manifolds.)



If we work over  $\mathbb{R}$  and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

## CHAINS OF FACTORIZATION HOMOLOGY OVER $\mathbb{R}$

If we work over  $\mathbb{R}$  and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

Theorem (Kontsevich '99; Tamarkin '03 ( $n = 2$ ); Lambrechts–Volić '14; Petersen '14 ( $n = 2$ ); Fresse–Willwacher '15)

The operad  $C_*(uE_n)$  is formal:  $C_*(uE_n) \simeq ue_n := H_*(uE_n)$ .

# CHAINS OF FACTORIZATION HOMOLOGY OVER $\mathbb{R}$

If we work over  $\mathbb{R}$  and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

**Theorem (Kontsevich '99; Tamarkin '03 ( $n = 2$ ); Lambrechts–Volić '14; Petersen '14 ( $n = 2$ ); Fresse–Willwacher '15)**

The operad  $C_*(uE_n)$  is formal:  $C_*(uE_n) \simeq ue_n := H_*(uE_n)$ .

**Theorem (I.)**

$M$  closed, simply connected, smooth,  $\dim M \geq 4 \implies$

Lambrechts–Stanley model of  $C_*(E_M)$  as a right  $C_*(uE_n)$ -module:

$$LS_M = C_*^{CE}(\mathcal{M}^{n-*} \otimes \mathbf{Lie}_n[1-n]) + \text{action of } \mathbf{Com}.$$

# CHAINS OF FACTORIZATION HOMOLOGY OVER $\mathbb{R}$

If we work over  $\mathbb{R}$  and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

**Theorem (Kontsevich '99; Tamarkin '03 ( $n = 2$ ); Lambrechts–Volić '14; Petersen '14 ( $n = 2$ ); Fresse–Willwacher '15)**

The operad  $C_*(uE_n)$  is formal:  $C_*(uE_n) \simeq ue_n := H_*(uE_n)$ .

## **Theorem (I.)**

$M$  closed, simply connected, smooth,  $\dim M \geq 4 \implies$

Lambrechts–Stanley model of  $C_*(E_M)$  as a right  $C_*(uE_n)$ -module:

$$LS_M = C_*^{CE}(\mathcal{M}^{n-*} \otimes \mathbf{Lie}_n[1-n]) + \text{action of } \mathbf{Com}.$$

Upshot:  $C_*(\int_M A) \simeq LS_M \circ_{ue_n}^{\mathbb{L}} \tilde{A}$

# CHAINS OF FACTORIZATION HOMOLOGY OVER $\mathbb{R}$

If we work over  $\mathbb{R}$  and we just want chains:

$$C_*(\int_M A) \simeq C_*(E_M) \circ_{C_*(uE_n)}^{\mathbb{L}} C_*(A).$$

**Theorem (Kontsevich '99; Tamarkin '03 ( $n = 2$ ); Lambrechts–Volić '14; Petersen '14 ( $n = 2$ ); Fresse–Willwacher '15)**

The operad  $C_*(uE_n)$  is formal:  $C_*(uE_n) \simeq u\mathbf{e}_n := H_*(uE_n)$ .

## **Theorem (I.)**

$M$  closed, simply connected, smooth,  $\dim M \geq 4 \implies$

Lambrechts–Stanley model of  $C_*(E_M)$  as a right  $C_*(uE_n)$ -module:

$$\mathbf{LS}_M = C_*^{CE}(\mathcal{M}^{n-*} \otimes \mathbf{Lie}_n[1-n]) + \text{action of } \mathbf{Com}.$$

Upshot:  $C_*(\int_M A) \simeq \mathbf{LS}_M \circ_{u\mathbf{e}_n}^{\mathbb{L}} \tilde{A}$

$\implies$  we need to resolve  $A$  as a  $u\mathbf{e}_n$ -algebra.

## WEYL ALGEBRA $\mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

## WEYL ALGEBRA $\mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of  $u\mathbf{e}_n$ : free symmetric algebra and  $\{x_i, \xi_j\} = \delta_{ij}$

## WEYL ALGEBRA $\mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of  $u\mathbf{e}_n$ : free symmetric algebra and  $\{x_i, \xi_j\} = \delta_{ij}$

$\implies$  quadratic-(linear-)constant presentation



## WEYL ALGEBRA $\mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of  $u\mathbf{e}_n$ : free symmetric algebra and  $\{x_i, \xi_j\} = \delta_{ij}$

$\implies$  quadratic-(linear-)constant presentation

Quadratization  $qA = S(x_i, \xi_j)$  free symmetric algebra + zero bracket

## WEYL ALGEBRA $\mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of  $u\mathbf{e}_n$ : free symmetric algebra and  $\{x_i, \xi_j\} = \delta_{ij}$

$\implies$  quadratic-(linear-)constant presentation

Quadratization  $qA = S(x_i, \xi_j)$  free symmetric algebra + zero bracket

Koszul dual:  $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_i)$  cofree symmetric coalgebra + trivial cobracket

## WEYL ALGEBRA $\mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of  $u\mathbf{e}_n$ : free symmetric algebra and  $\{x_i, \xi_j\} = \delta_{ij}$

$\implies$  quadratic-(linear-)constant presentation

Quadratization  $qA = S(x_i, \xi_j)$  free symmetric algebra + zero bracket

Koszul dual:  $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_j)$  cofree symmetric coalgebra + trivial cobracket
- $d = 0$

## WEYL ALGEBRA $\mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of  $u\mathbf{e}_n$ : free symmetric algebra and  $\{x_i, \xi_j\} = \delta_{ij}$

$\implies$  quadratic-(linear-)constant presentation

Quadratization  $qA = S(x_i, \xi_j)$  free symmetric algebra + zero bracket

Koszul dual:  $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_j)$  cofree symmetric coalgebra + trivial cobracket
- $d = 0$
- curvature:  $\theta(\bar{x}_i \wedge \bar{\xi}_j) = -\delta_{ij}$ .

## WEYL ALGEBRA $\mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of  $u\mathbf{e}_n$ : free symmetric algebra and  $\{x_i, \xi_j\} = \delta_{ij}$

$\implies$  quadratic-(linear-)constant presentation

Quadratization  $qA = S(x_i, \xi_j)$  free symmetric algebra + zero bracket

Koszul dual:  $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_j)$  cofree symmetric coalgebra + trivial cobracket
- $d = 0$
- curvature:  $\theta(\bar{x}_i \wedge \bar{\xi}_j) = -\delta_{ij}$ .

$\implies$  “small” resolution  $Q_A := \Omega_\kappa A^i = (SLS^c(\bar{x}_i, \bar{\xi}_j), d) \xrightarrow{\sim} A$

# WEYL ALGEBRA $\mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

$$A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) = S(x_1, \dots, x_d, \xi_1, \dots, \xi_d)$$

Action of  $u\mathbf{e}_n$ : free symmetric algebra and  $\{x_i, \xi_j\} = \delta_{ij}$

$\implies$  quadratic-(linear-)constant presentation

Quadratization  $qA = S(x_i, \xi_j)$  free symmetric algebra + zero bracket

Koszul dual:  $A^i = (qA^i, d, \theta)$

- $qA^i = S^c(\bar{x}_i, \bar{\xi}_j)$  cofree symmetric coalgebra + trivial cobracket
- $d = 0$
- curvature:  $\theta(\bar{x}_i \wedge \bar{\xi}_j) = -\delta_{ij}$ .

$\implies$  “small” resolution  $Q_A := \Omega_\kappa A^i = (SLS^c(\bar{x}_i, \bar{\xi}_j), d) \xrightarrow{\sim} A$

(If we had applied curved KD at the level of operads instead:

$$\Omega_\kappa B_\kappa A \supset (\underbrace{SL}_{\text{cobar}} \underbrace{S^c L^c}_{\text{bar}} \underbrace{S(x_i, \xi_j)}_A, d), + \text{ resolution of the unit...})$$

## COMPUTATION OF $\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

We can also compute

$$\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) \simeq \mathbf{LS}_M \circ_{ue_n} (SLS^c(\bar{x}_i, \bar{\xi}_j), d)$$

## COMPUTATION OF $\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

We can also compute

$$\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) \simeq \text{LS}_M \circ_{ue_n} (SLS^c(\bar{x}_i, \bar{\xi}_j), d)$$

Theorem (I. '18, see also Markarian '17, Döppenschmitt '18)

$$\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) \simeq C_*^{CE}(\mathcal{M}^{n-*} \otimes \mathbb{R}\langle 1, x_i, \xi_j \rangle) \simeq \mathbb{R}.$$



## COMPUTATION OF $\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n])$

We can also compute

$$\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) \simeq \text{LS}_M \circ_{ue_n} (SLS^c(\bar{x}_i, \bar{\xi}_j), d)$$

**Theorem (I. '18, see also Markarian '17, Döppenschmitt '18)**

$$\int_M \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1-n]) \simeq C_*^{CE}(\mathcal{M}^{n-*} \otimes \mathbb{R}\langle 1, x_i, \xi_j \rangle) \simeq \mathbb{R}.$$

Intuition: quantum observable with values in  $A \rightsquigarrow$  “expectation” lives in  $\int_M A$ , should be a number.

## APPLICATION 2: DERIVED ENVELOPING ALGEBRA

Operad  $\mathbf{P}$  +  $\mathbf{P}$ -algebra  $A \implies$  notion of  $A$ -modules

## APPLICATION 2: DERIVED ENVELOPING ALGEBRA

Operad  $\mathbf{P}$  +  $\mathbf{P}$ -algebra  $A \implies$  notion of  $A$ -modules

### Examples

$\mathbf{P} = \mathbf{Ass} \rightarrow (A, A)$  bimodules;  $\mathbf{P} = \mathbf{Com} \rightarrow A$ -modules;  $\mathbf{P} = \mathbf{Lie} \rightarrow$  representations of the Lie algebra.

## APPLICATION 2: DERIVED ENVELOPING ALGEBRA

Operad  $\mathbf{P}$  +  $\mathbf{P}$ -algebra  $A \implies$  notion of  $A$ -modules

### Examples

$\mathbf{P} = \mathbf{Ass} \rightarrow (A, A)$  bimodules;  $\mathbf{P} = \mathbf{Com} \rightarrow A$ -modules;  $\mathbf{P} = \mathbf{Lie} \rightarrow$  representations of the Lie algebra.

$\exists$  an associative algebra  $U_{\mathbf{P}}(A)$  s.t. left  $U_{\mathbf{P}}(A)$ -modules =  $A$ -modules

## APPLICATION 2: DERIVED ENVELOPING ALGEBRA

Operad  $\mathbf{P}$  +  $\mathbf{P}$ -algebra  $A \implies$  notion of  $A$ -modules

### Examples

$\mathbf{P} = \mathbf{Ass} \rightarrow (A, A)$  bimodules;  $\mathbf{P} = \mathbf{Com} \rightarrow A$ -modules;  $\mathbf{P} = \mathbf{Lie} \rightarrow$  representations of the Lie algebra.

$\exists$  an associative algebra  $U_{\mathbf{P}}(A)$  s.t. left  $U_{\mathbf{P}}(A)$ -modules =  $A$ -modules

### Proposition

For  $A = \mathcal{O}_{\text{poly}}(T^*\mathbb{R}^d[1 - n])$ , the derived enveloping algebra  $U_{\text{ue}_n}^{\mathbb{L}}(A)$  is q.iso to the underived one.

**THANK YOU FOR YOUR ATTENTION!**

These slides: <https://idrissi.eu>