

# Quantum chromodynamics on your smartphone using 'SeekYouToo' : A workman guide

May 1, 2020

**Abstract.** Monte Carlo studies of non-abelian gauge theories emerged in the early eighties now establishing a standard tool of modern lattice quantum field theory. Due to the dramatic increase of mobile computational power in the 2010s it is now possible to carry out very basic simulations on smart phones. The author describes an according cross-platform implementation of  $SU_2$  gluon dynamics and discusses some didactical aspects.

## A basic sketch of quantum chromodynamics

### Electrodynamics by concepts

Quantum chromodynamics (QCD) describes the strong interaction of quarks and gluons. Quarks make up hadrons such as the protons and neutron forming themselves the nuclei of atoms. Gluons glue together the quarks implementing the strong force. Quarks have spin 1/2 and thus are fermions (i.e. particles with half odd integer spin) in contrast to gluon carrying spin 1 and being bosons (particles with integer spin). Spin plays an essential statistical role : Fermions obey the Pauli exclusion principle stating that two identical fermions cannot occupy the same quantum state simultaneously. Thus electrons live in distinct shells around the nuclei of atoms. Gluons play the same role as photons, being the carriers of electromagnetic force, yet gluons can also interact with themselves.

Electrodynamics describes the electromagnetics fields. Historically Gauss, Ampere and Faraday formulated distinct equations for the electric and magnetic fields,  $\mathbf{E}$  and  $\mathbf{B}$ .

The magnetic field has no monopoles and thus divergenceless as expressed in Gauss's law  $\nabla \cdot \mathbf{B} = 0$  of the magnetic field. As the divergence of a curl always vanishes the magnetic field can be written as the curl of its potential  $\mathbf{B} = \nabla \times \mathbf{A}$ .

Faradays law states a changing magnetic field induces a voltage in a coil. That is the path integral of the electric field around a closed loop equal to the negative rate of the change of the magnetic flux through the area enclosed by the loop:  $\oint_C \mathbf{E} \cdot d\mathbf{r} = -\partial \Phi / \partial t$ . Put in vector notation we have  $\nabla \times \mathbf{E} = -\partial \mathbf{B} / \partial t$ .

Using the magnetic potential  $\mathbf{A}$  we have  $\nabla \times (\mathbf{E} + \partial\mathbf{A}/\partial t) = 0$ . When the curl of a vector field vanishes the field can always be written as the gradient of a scalar, giving  $\mathbf{E} = -\nabla V - \partial\mathbf{A}/\partial t$  with  $V$  being the electric potential.

As the electric and magnetic field only depends on the changes of the recordings potentials we assume it is possible that we can change the potentials and obtains the same fields. If we make the transformation  $\mathbf{A} \rightarrow \mathbf{A} + \nabla\lambda$  the magnetic field  $\mathbf{B}$  does not change since the curl of a gradient vanishes,  $\mathbf{B} = \nabla \times \mathbf{A} + \nabla \times \nabla\lambda = \nabla \times \mathbf{A}$ . However the electric field is changed,  $\mathbf{E} = -\nabla V - \partial\mathbf{A}/\partial t - \nabla\partial\lambda/\partial t = -\nabla(V + \partial\lambda/\partial t) - \partial\mathbf{A}/\partial t$ .

If another change  $V \rightarrow V - \partial\lambda/\partial t$  is made also  $\mathbf{E}$  remains unchanged. A partial choice of the potentials is called a gauge and can be done in many ways.

Gauss's law for the electric field  $\nabla \cdot \mathbf{E} = \rho/\epsilon_0$ , and Ampere's law  $\nabla \times \mathbf{B} = \mu_0\mathbf{J} + \mu_0\epsilon_0\partial\mathbf{E}/\partial t$  have not been mentioned yet completing the set of the of the "four macroscopic Maxwell equations". Using an appropriate gauge condition, the Lorenz (Ludvig) gauge  $\nabla \cdot \mathbf{A} + \mu_0\epsilon_0(\partial V/\partial t) = 0$  lets one write the inhomogeneous Maxwell equations in a concise fashion

$$\begin{aligned}\square^2 V &= \mu_0\epsilon_0 \frac{\partial V}{\partial t} - \nabla^2 V = \frac{\rho}{\epsilon_0} \\ \square^2 \mathbf{A} &= \mu_0\epsilon_0 \frac{\partial \mathbf{A}}{\partial t} - \nabla^2 \mathbf{A} = \mu_0\mathbf{J}\end{aligned}$$

Though the electric potential relates to static charges and the magnetic one to moving charges the expressional symmetry of this appealing formulation suggests that the distinction between electric and magnetic fields are a question about point of view and that the electric and magnetic fields form one field, the electromagnetic field.

Maxwell carried out a relativistic formulation taking into account Lorentz (Hendrik) transformations. That is, he analysed how the fields transform when changing from a coordinate frame in Minkoskian spacetime to some other frame beeing boosted at constant velocity relative to the original frame. These transformation can be represented by 4x4 real matrices  $\Lambda$  (parametrized by the velocity  $v$ ) such that a transformed spacetime-vector has the same scalar (using the Minkowski metric) as the original vector. The transformation of the electric and magnetic field can be written in a concise form letting the Lorentz group matrices  $\Lambda$  act on the electromagnetic field tensor

$$F_{\mu\nu} = \begin{bmatrix} 0 & -E_x/c & -E_y/c & -E_z/c \\ E_x/c & 0 & -B_z & B_y \\ E_y/c & B_z & 0 & -B_x \\ E_z/c & -B_y & B_x & 0 \end{bmatrix}$$

Using the four-vector potential given by  $A_\mu = (V/c, A_x, A_y, A_z)$  and the four-current defined as  $J^\nu = (c\rho, J_x, J_y, J_z)$  the inhomogeneous Maxwell equation can be written in the very compact form

$$\partial_\mu F^{\mu\nu} = \mu_0 J^\nu$$

with

$$F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$$

Note that in the 4-vector notation greek indices run from 0 to 3 and the Einstein summation convention is applied when an index is repeated both lowered and raised, that is for two 4-vectors  $X^\mu$  and  $Y_\mu$  the expression  $X^\mu Y_\mu$  means  $X^\mu Y_\mu = \sum_{\mu=0}^3 X^\mu Y_\mu$ .

Also note that a second order tensor (a matrix)  $X_{\mu\nu}$  with only lowered indices is called covariant and that according contravariant tensor (raised indices) is build by applying the metric tensor  $\eta$

$$X^{\mu\nu} = \eta^{\mu\alpha} \eta^{\nu\beta} X_{\alpha\beta}; \quad \eta^{\mu\nu} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

One important feature of the 4-vector notation to be mentioned is that the product of a contravariant and a covariant tensor  $X^\mu X_\mu$  is invariant under Lorentz transformation.

It is easy to show that the Maxwell equation is the Euler-Lagrange equation of the Lagrangian

$$L = \mathcal{L}_{field} + \mathcal{L}_{interaction} = -\frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - J^\mu A_\mu.$$

## Gauge theories and symmetries

Noether's theorem states that for every differentiable global symmetry of the action of a Lagrangian system there is a corresponding conservation law. For instance time translations give conservation of energy and spatial translations yield conservation of energy. If we have for example a complex-valued scalar field  $\phi$  being invariant under some  $U(1)$  transformation,  $\phi(x) \rightarrow e^{i\alpha} \phi(x)$  where  $\alpha$  does not depend on position  $x$  one can construct an according conserved Noether current. If however we have a position-dependent  $\alpha(x)$  and the Lagrangian involves derivatives we obtain unwanted term destroying the invariance, because at each point we have different transformation roles. In order to have obtain local symmetry we tweak the partial derivative introducing a new gauge potential  $A_\mu$ : The covariant derivative

$$D_\mu = \partial_\mu - igA_\mu$$

eliminates the terms introduced by the partial derivative, arguing as follows. We assume that  $A_\mu$  transforms as  $A_\mu \rightarrow A_\mu + (1/g)\partial_\mu \alpha(x)$ . By comparing to the gauge discussed for the electromagnetic field,  $V \rightarrow V - \partial\lambda/\partial t$  and  $\mathbf{A} \rightarrow \mathbf{A} + \nabla\lambda$ , we see that  $\lambda$  is equivalent to  $\alpha(x)/g$  with  $g$  being the coupling constant (being  $e$  in the case of electrodynamics). Using some algebra we see

that the covariant derivative applied to the phase transformed field together with the gauge transformation gives us an expression in which the covariant derivative itself remains does not depend on the phase:

$$\begin{aligned}
D_\mu &\rightarrow \left[ \partial_\mu + ig \left( A_\mu - \frac{1}{g} \partial_\mu \alpha(x) \right) \right] e^{i\alpha(x)} \phi = \\
&= e^{i\alpha(x)} \partial_\mu \phi \\
&\quad + i e^{i\alpha(x)} \phi \partial_\mu \alpha(x) + ig A_\mu e^{i\alpha(x)} \phi \\
&\quad - i e^{i\alpha(x)} \phi \partial_\mu \alpha(x) \\
&= e^{i\alpha(x)} (\partial_\mu - ig A_\mu) \phi \\
&= e^{i\alpha(x)} D_\mu \phi
\end{aligned}$$

As the derivative  $D_\mu$  transforms the same way as the field  $\phi$  it is the name “covariant” suits pretty well.

Using covariant derivatives we can thus construct meaningful Lagrangians preserving local symmetry. The simple real gauge theory is quantum electrodynamics (QED) augmenting electrodynamics by fermions (for instance electrons) described by Dirac spinors  $\psi$ :

$$\begin{aligned}
\mathcal{L} &= \mathcal{L}_{Dirac} + \mathcal{L}_{Maxwell} + \mathcal{L}_{interaction} \\
&= \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - e \bar{\psi} \gamma^\mu \psi A_\mu
\end{aligned}$$

Using the covariant derivative the Lagrangian reads

$$\mathcal{L} = \bar{\psi} (i\gamma^\mu D_\mu - m) \psi - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$$

and one obtains as the equation of motions for the fermionic wave function  $\psi$  and the electromagnetic potential  $A_\mu$

$$\begin{aligned}
(i\gamma^\mu D_\mu - m) \psi &= 0 \\
\partial_\mu F^{\mu\nu} &= e \bar{\psi} \gamma^\nu \psi
\end{aligned}$$

The symmetry group of QED is  $U(1)$ , that is the Lagrangian is invariant under  $\psi \rightarrow e^{i\alpha(x)} \psi$  and the according gauge field is the electromagnetic potential.

$U(1)$  is a simple non-trivial symmetry group and we now turn to the case to more involved case that the groups in question are non-abelian. Yang and Mills therefore considered transformations of the form  $\phi \rightarrow \exp(i\alpha^a t^a) \phi$  where ...

## Quantum chromodynamics

$$\mathcal{L}_{electromagnetic} = F_{\mu\nu}F^{\mu\nu}$$

$$\mathcal{L}_{gluonic} = G_{\mu\nu}G^{\mu\nu}$$