

In [1]:

```
# Import modules
# Make sure you have installed netCDF4, xarray, and nc-time-axis
import numpy as np
import pandas as pd
import xarray as xr
from matplotlib import pyplot as plt
%matplotlib inline
```

In [2]:

```
# Prerequisites

# Load modulesimport numpy as np

import matplotlib as mpl
import matplotlib.pyplot as plt
import matplotlib.gridspec as gridspec
```

In [3]:







```
# 打开文件
ds= xr.open_dataset("NOAA_NCDC_ERSST_v3b_SST.nc")
ds
```

Out[3]:



xarray.Dataset

► Dimensions: (lat: 89, lon: 180, time: 684)

▼ Coordinates:

lat	(lat)	float32	-88.0 -86.0 -84.0 ... 86.0 88.0	 
lon	(lon)	float32	0.0 2.0 4.0 ... 354.0 356.0 35...	 
time	(time)	datetime64[ns]	1960-01-15 ... 2016-12-15	 

▼ Data variables:

sst	(time, lat, lon)	float32	...	 
-----	------------------	---------	-----	---

▼ Attributes:

Conventions :	IRIDL
source :	https://iridl.ldeo.columbia.edu/SOURCES/.NOAA/.NCDC/.ERSST/.version3b/.sst/
history :	extracted and cleaned by Ryan Abernathey for Research Computing in Earth Science

In [4]:







```
#1. 1
#选择区域
ds_r = ds.sel(lat=slice(-5,5), lon=slice(190,240))
ds_r
```

Out[4]:

xarray.Dataset

► Dimensions: (lat: 5, lon: 26, time: 684)

▼ Coordinates:

lat	(lat)	float32	-4.0 -2.0 0.0 2.0 4.0	 
lon	(lon)	float32	190.0 192.0 194.0 ... 238.0 2...	 
time	(time)	datetime64[ns]	1960-01-15 ... 2016-12-15	 

▼ Data variables:

sst	(time, lat, lon)	float32	...	 
------------	------------------	---------	-----	---

▼ Attributes:

Conventions : IRIDL

source : <https://iridl.ldeo.columbia.edu/SOURCES/.NOAA/.NCDC/.ERSST/.version3b/.sst/>

history : extracted and cleaned by Ryan Abernathey for Research Computing in Earth Science

In []:

In [5]:

```
#计算每月sst的均值
ds_monthly=ds_r.sst.groupby(ds.time.dt.month).mean(dim=['lat', 'lon', 'time'])
# ds_monthly=ds.sst.groupby(ds.time.dt.year.month).mean()
ds_monthly
```

Out[5]:

xarray.DataArray 'sst' (month: 12)

```
array([26.56812, 26.742603, 27.239906, 27.694029, 27.79552, 27.598068,
       27.199274, 26.824581, 26.7382, 26.717516, 26.693666, 26.61345 ],
      dtype=float32)
```

▼ Coordinates:

month	(month)	int64	1 2 3 4 5 6 7 8 9 10 11 12	 
--------------	---------	-------	----------------------------	---

► Attributes: (0)

In [6]:

```

#1.1
#借鉴https://www.cnblogs.com/traditional/p/13776180.html
#计算Niño 3.4
ds_rol = ds_monthly.rolling(month=3, center=True).mean()


ds_rol
#计算异常值作图
sst_an=ds_monthly-ds_rol
sst_an
sst_an.plot()

#1.2
#借鉴https://www.cnblogs.com/Gaoqiking/p/11069517.html
# plt.plot(ds_monthly.month, sst_an)
# plt.fill_between(ds_monthly.month.to_numpy(), sst_an, where=(sst_an>0), color='red')
# plt.fill_between(ds_monthly.month.to_numpy(), sst_an, where=(sst_an<0), color='blue')
# 按照月分组后，年不见了不清楚如何将每年的各个月分组进行计算

```

Out[6]:

[<matplotlib.lines.Line2D at 0x1e88876b520>]



In [7]:







```
ds = xr.open_dataset("CERES_EBAF-TOA_200003-201701.nc", engine="netcdf4")
ds
#2. 1
#绘制长、短和太阳辐射波的均值2D图
# long=ds.toa_lw_all_mon.mean(dim='time')
# long.plot()
# # short=ds.toa_sw_all_mon.mean(dim='time')
# short.plot()
# solar=ds.solar_mon.mean(dim='time')
# solar.plot()
```

Out[7]:












xarray.Dataset

► Dimensions: (lon: 360, time: 203, lat: 180)

▼ Coordinates:

lon	(lon)	float32	0.5 1.5 2.5 ... 357.5 358.5 35...	 
time	(time)	datetime64[ns]	2000-03-15 ... 2017-01-15	 
lat	(lat)	float32	-89.5 -88.5 -87.5 ... 88.5 89.5	 

▼ Data variables:

toa_sw_all_mon	(time, lat, lon)	float32	...	 
toa_lw_all_mon	(time, lat, lon)	float32	...	 
toa_net_all_mon	(time, lat, lon)	float32	...	 
toa_sw_clr_mon	(time, lat, lon)	float32	...	 
toa_lw_clr_mon	(time, lat, lon)	float32	...	 
toa_net_clr_mon	(time, lat, lon)	float32	...	 
toa_cre_sw_mon	(time, lat, lon)	float32	...	 
toa_cre_lw_mon	(time, lat, lon)	float32	...	 
toa_cre_net_mon	(time, lat, lon)	float32	...	 
solar_mon	(time, lat, lon)	float32	...	 
cldarea_total_d...	(time, lat, lon)	float32	...	 
cldpress_total_...	(time, lat, lon)	float32	...	 
cldtemp_total_d...	(time, lat, lon)	float32	...	 
cldtau_total_da...	(time, lat, lon)	float32	...	 

▼ Attributes:

title :	CERES EBAF (Energy Balanced and Filled) TOA Fluxes. Monthly Av erages and 07/2005 to 06/2015 Climatology.
institution :	NASA/LaRC (Langley Research Center) Hampton, Va
Conventions :	CF-1.4
comment :	Data is from East to West and South to North.
Version :	Edition 4.0; Release Date March 7, 2017
Fill_Value :	Fill Value is -999.0
DOI :	10.5067/TERRA+AQUA/CERES/EBAF-TOA_L3B.004.0
Production_Files :	List of files used in creating the present Master netCDF file: /homedir/nloeb/ebaf/monthly_means/adj_fluxes/deliverable/sw*.gz /homedir/nloeb/ebaf/monthly_means/adj_fluxes/deliverable/lw*.gz


```
/homedir/nloeb/ebaf/monthly_means/adj_fluxes/deliverable/net*.gz  
/homedir/nloeb/ebaf/monthly_means/adj_fluxes/deliverable/solflx*.gz  
/homedir/nloeb/ebaf/monthly_means/out_glob.dat
```

In [8]:

```
#2.1  
#一开始不太理解，向谢栋学姐请教相关问题  
#绘制长、短和太阳辐射波的均值2D图  
long=ds.toa_lw_all_mon.mean(dim='time')  
long.plot()  
# short=ds.toa_sw_all_mon.mean(dim='time')  
# short.plot()  
# solar=ds.solar_mon.mean(dim='time')  
# solar.plot()
```

Out[8]:

<matplotlib.collections.QuadMesh at 0x1e888f3efd0>




In [37]:

```
short=ds.toa_sw_all_mon.mean(dim='time')  
short.plot()
```

Out[37]:

<matplotlib.collections.QuadMesh at 0x2ca3dde8c40>




In [60]:

```
solar=ds.solar_mon.mean(dim='time')  
solar.plot()
```

Out[60]:

<matplotlib.collections.QuadMesh at 0x188a1b6b850>




In [69]:

```
# ds1=ds.solar_mon.mean(dim="time")-ds.toa_lw_all_mon.mean(dim="time")-ds.toa_sw_all_mon.mean(dim="t
#TOA net=solar-long-short
add=solar-long-short
add.plot()

# ds1.plot()
```

Out[69]:

<matplotlib.collections.QuadMesh at 0x188a54dd910>




In [9]:

```
#绘制TOA net的图
toa_net=ds.toa_net_all_mon.mean(dim='time')
toa_net.plot()
#TOA的图与三种波长相叠加的图相同
```

Out[9]:

```
<matplotlib.collections.QuadMesh at 0x1e889339d30>
```



In [10]:

```
#2.2
#太阳光发射的
launch=ds.solar_mon.mean(dim='time')
#创建加权
weights=np.cos(np.deg2rad(ds.lat))
# launch
w_solar=solar.weighted(weights)
w_long=long.weighted(weights)
w_short=short.weighted(weights)

# 计算总的均值
w_solar_mean=w_solar.mean()
w_long_mean=w_long.mean()
w_short_mean=w_short.mean()
print(w_solar_mean, w_long_mean, w_short_mean)
```

```
-----
NameError                                Traceback (most recent call last)
<ipython-input-10-882cbe10a16e> in <module>
      5 weights=np.cos(np.deg2rad(ds.lat))
      6 # launch
----> 7 w_solar=solar.weighted(weights)
      8 w_long=long.weighted(weights)
      9 w_short=short.weighted(weights)
```


```
NameError: name 'solar' is not defined
```


In [87]:

```
#2.3  
#以经度为基准进行求和并绘图  
ra_sum=ds.toa_cre_net_mon.sum(dim='lon')  
ra_sum.transpose().plot()
```

Out[87]:

<matplotlib.collections.QuadMesh at 0x188a8921d30>







In [11]:

```
#2.4
#计算平均云面积
cldarea = ds.cldarea_total_daynight_mon.mean(dim='time')
high_area = (cldarea>=75)
low_area = (cldarea<=25)

#计算长波和短波的平均辐射值
long_v=ds.toa_lw_all_mon.mean(dim='time')
short_v=ds.toa_sw_all_mon.mean(dim='time')

#绘图
long_v.where(high_area).plot()
plt.show()
long_v.where(low_area).plot()
plt.show()
short_v.where(low_area).plot()
plt.show()
short_v.where(high_area).plot()
plt.show()
```







In [12]:

```
#2.5
#计算短波和长波全球的平均值
long_t=ds.toa_lw_all_mon.mean(dim='time')
short_t=ds.toa_sw_all_mon.mean(dim='time')

#计算云面积的平均值
cldarea = ds.cldarea_total_daynight_mon.mean(dim='time')

#绘图
long_v.where(cldarea).plot()
plt.show()

short_v.where(cldarea).plot()
plt.show()
```



In [13]:

打开文件







```
ds = xr.open_dataset("sst.mnmean.nc")
ds
```

Out[13]:





xarray.Dataset

► Dimensions: (lat: 89, lon: 180, time: 2014, nbnds: 2)

▼ Coordinates:

lat	(lat)	float32	88.0 86.0 84.0 ... -86.0 -88.0		
lon	(lon)	float32	0.0 2.0 4.0 ... 354.0 356.0 ...		
time	(time)	datetime64[ns]	1854-01-01 ... 2021-10-01		

▼ Data variables:

time_bnds	(time, nbnds)	float64	...		
sst	(time, lat, lon)	float32	...		


► Attributes: (37)

In [14]:

```
#没有找到正确的方法下载带有连续时间的数据文件，所以应用了之前老师上传的一个数据文件
#3.1
ds.sst.mean(dim=['lat', 'lon']).transpose().plot()
```


Out[14]:

[<matplotlib.lines.Line2D at 0x1e88ac56ee0>]




In [15]:

```
#3.2  
ds.sst.mean(dim='lon').plot()  
plt.show()
```




In [16]:

```
#3.2  
ds.sst.mean(dim='lat').plot()  
plt.show()
```




In [17]:

```
#3.2
ds.sst.mean(dim='time').plot()
plt.show()
```



In [18]:

```
#3.2
data = ds.sst.sel(time=slice('2013-01-01', '2014-12-31'))
data1 = data.where(data < 10)
data1.plot()
plt.show()
```



In [19]:

```
ds.sst.groupby('time.month').mean().mean(dim='lon').transpose().plot.contourf()  
plt.show()
```

