

Table 2. Adjusted per-SD ordinal odds ratios for morphology and aSAH severity

Proportional-odds ordinal regression, one index per model; OR per 1 SD, OR > 1 = higher severity

Predictor	No.	Complete-case OR (95% CI), <i>p</i>	Multiple imputation OR (95% CI), <i>p</i>	MI FDR
modified Fisher				
Size ratio	99	2.04 (1.16–3.60), 0.013	1.67 (1.17–2.39), 0.004	0.046
Aspect ratio	97	1.26 (0.75–2.14), 0.384	1.46 (1.01–2.10), 0.042	0.115
Max diameter	99	1.13 (0.64–2.00), 0.671	1.33 (0.93–1.89), 0.113	0.183
Volume (log)	98	1.16 (0.69–1.93), 0.582	1.35 (0.96–1.88), 0.082	0.148
Neck width	99	1.03 (0.68–1.55), 0.904	1.09 (0.81–1.48), 0.576	0.712
Height/width ratio	99	0.90 (0.60–1.34), 0.606	1.01 (0.74–1.38), 0.941	0.941
Neck/parent ratio	99	1.51 (0.95–2.39), 0.080	1.39 (0.99–1.96), 0.058	0.122
Hunt & Hess				
Size ratio	99	1.56 (1.01–2.39), 0.043	1.54 (1.10–2.16), 0.011	0.047
Aspect ratio	97	1.37 (0.92–2.03), 0.123	1.71 (1.22–2.40), 0.002	0.038
Max diameter	99	1.47 (0.86–2.53), 0.162	1.52 (1.08–2.12), 0.016	0.055
Volume (log)	98	1.41 (0.85–2.34), 0.185	1.61 (1.14–2.27), 0.007	0.047
Neck width	99	1.02 (0.67–1.55), 0.932	1.08 (0.79–1.46), 0.634	0.740
Height/width ratio	99	0.97 (0.66–1.42), 0.878	1.14 (0.84–1.54), 0.402	0.528
Neck/parent ratio	99	1.23 (0.80–1.90), 0.341	1.27 (0.94–1.72), 0.127	0.190
WFNS				
Size ratio	103	1.82 (1.16–2.87), 0.010	1.38 (1.00–1.91), 0.049	0.115
Aspect ratio	101	1.46 (0.98–2.18), 0.061	1.50 (1.10–2.05), 0.010	0.047
Max diameter	103	1.69 (0.97–2.95), 0.062	1.34 (0.96–1.87), 0.085	0.148
Volume (log)	102	1.45 (0.86–2.42), 0.162	1.42 (1.00–2.01), 0.049	0.115
Neck width	103	1.03 (0.66–1.61), 0.885	1.05 (0.75–1.48), 0.772	0.811
Height/width ratio	103	1.05 (0.71–1.55), 0.805	1.07 (0.77–1.49), 0.682	0.754
Neck/parent ratio	103	1.26 (0.80–1.99), 0.313	1.20 (0.86–1.68), 0.280	0.392

Values are proportional-odds (cumulative) odds ratios per 1 SD from separate single-index models, adjusted for age, sex, smoking, hypertension, diabetes, statin use, and aneurysm circulation. Multiple imputation: MICE, *m* = 20. Benjamini–Hochberg FDR controls across the 21 ordinal tests (7 indices × 3 grades); **bold** = FDR < 0.05. The continuous outcome (admission GCS) was analysed separately by linear regression. SD, standard deviation.