

# ViReport v0.0.1

Niema Moshiri

2020-03-20

## 1 Input Dataset

The analysis was conducted on a dataset containing 867 sequences. The average sequence length was 29814.42, with a standard deviation of 242.885. The earliest sample date was 2019-12-24, the median sample date was 2020-03-01, and the most recent sample date was 2020-03-13.




Figure 1: Distribution of input sequence lengths

## 2 Preprocessed Dataset

The input dataset was preprocessed such that sequences were given safe names: non-letters/digits in sequence IDs were converted to underscores. After preprocessing, the dataset contained 867 sequences. The average sequence length was 29814.42, with a standard deviation of 242.885. The earliest sample date was 2019-12-24, the median sample date was 2020-03-01, and the most recent sample date was 2020-03-13.

## 3 Multiple Sequence Alignment

Multiple sequence alignment was performed using MAFFT (Katoh & Standley, 2013) in automatic mode. There were 30142 positions (1032 invariant) and 769 unique sequences in the multiple sequence alignment. Pairwise distances were computed from the multiple sequence alignment using the tn93 tool of HIV-TRACE (Pond et al., 2018). The average pairwise sequence distance was 0.000303, with a standard deviation of 0.000186.




Figure 2: Distribution of input sample dates

Across the positions of the multiple sequence alignment that had non-zero Shannon entropy, the minimum Shannon entropy was 0.0131, the maximum Shannon entropy was 2.122, and the average Shannon entropy was 0.0899, with a standard deviation of 0.119.

## 4 Phylogenetic Inference

A maximum-likelihood phylogeny was inferred under the General Time-Reversible (GTR) model (Tavare, 1986) using FastTree 2 (Price et al., 2010) using a Gamma20-based likelihood. The inferred phylogeny was MinVar-rooted using FastRoot (Mai et al., 2017). Pairwise distances were computed from the phylogeny using TreeSwift (Moshiri, 2020). The maximum pairwise phylogenetic distance (i.e., tree diameter) was 0.00581, and the average pairwise phylogenetic distance was 0.00148, with a standard deviation of 0.000989.

## 5 Phylogenetic Dating

The rooted phylogeny was dated using treedater (Volz & Frost, 2017). The height of the dated tree was 96.145 days, so given that the most recent sample was collected on 2020-03-13, the estimated time of the most recent common ancestor (tMRCA) was 2019-12-07.

## 6 Ancestral Sequence Reconstruction

Ancestral sequence reconstruction was performed using TreeTime (Sagulenko et al., 2018).

## 7 Transmission Clustering

Transmission clustering was performed using TreeN93 (Moshiri, 2018) using pairwise phylogenetic distances. The total number of singletons (i.e., non-clustered individuals) was 146, and the total number of clusters




Figure 3: Distribution of preprocessed sequence lengths

(excluding singletons) was 43. The average cluster size (excluding singletons) was 16.488, with a standard deviation of 32.133, and the maximum and minimum cluster sizes were 139 and 2, respectively.

## 8 Citations

- Katoh K., Standley D.M. (2013). "MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability". *Molecular Biology and Evolution*. 30(4), 772-780.
- Le S.Q., Gascuel O. (2008). "An Improved General Amino Acid Replacement Matrix". *Molecular Biology and Evolution*. 25(7), 1307-1320.
- Mai U., Sayyari E., Mirarab S. (2017). "Minimum Variance Rooting of Phylogenetic Trees and Implications for Species Tree Reconstruction". *PLoS ONE*. 12(8), e0182238.
- Moshiri N. (2018). "TreeN93: a non-parametric distance-based method for inferring viral transmission clusters". *bioRxiv*.
- Moshiri N. (2020). "TreeSwift: a massively scalable Python tree package". *SoftwareX*. In press.
- Moshiri N. (2020). "ViReport" (<https://github.com/niemasd/ViReport>).
- Pond S.L.K., Weaver S., Leigh Brown A.J., Wertheim J.O. (2018). "HIV-TRACE (TRANsmiission Cluster Engine): a Tool for Large Scale Molecular Epidemiology of HIV-1 and Other Rapidly Evolving Pathogens". *Molecular Biology and Evolution*. 35(7), 1812-1819.
- Price M.N., Dehal P.S., Arkin A.P. (2010). "FastTree 2 – Approximately Maximum-Likelihood Trees for Large Alignments". *PLoS ONE*. 5(3), e9490.
- Sagulenko P., Puller V., Neher R.A. (2018). "TreeTime: Maximum-likelihood phylodynamic analysis". *Virus Evolution*. 4(1), vex042.




Figure 4: Distribution of preprocessed sample dates

- Tavaré S. (1986). "Some Probabilistic and Statistical Problems in the Analysis of DNA Sequences". Lectures on Mathematics in the Life Sciences. 17, 57-86.
- Volz E.M., Frost S.D.W. (2017). "Scalable relaxed clock phylogenetic dating". Virus Evolution. 3(2), vex025.




Figure 5: Distribution of pairwise sequence distances




Figure 6: Shannon entropy across the positions of the multiple sequence alignment. A significance threshold was computed using Tukey's Rule:  $1.5 \times$  the interquartile range added to the third quartile, which was 0.194. The significance threshold is shown as a red dashed line, and significant points are shown in red.



Figure 7: Rooted phylogenetic tree in unit of expected per-site mutations




Figure 8: Distribution of pairwise phylogenetic distances



Figure 9: Dated phylogenetic tree in unit of years






Figure 10: Distribution of cluster sizes (excluding singletons)