

DETECTING ECCENTRICITY: WHAT IS THE CURRENT EVIDENCE?

NIHAR GUPTA

Max Planck Institute for Gravitational Physics and University of Maryland

GW Eccentricity Corfu, Greece September 2026



TABLE OF CONTENTS

01 What creates eccentric binary black holes and why are we interested in them?

02 How do we determine what constitutes a discovery?

03 Group project: let's make a prior together!

04 Parameter estimation on real data, going event-by-event. Going into details!

05 Population properties of eccentric binaries and their application to individual events

06 Eccentric searches, are we missing eccentric events?

07 Conclusions, what can we take away from current observations?

WHAT IS DISCOVERY: WHAT IS BAYES?

Bayes/Laplace told us that


$$p(\overset{\text{Parameters}}{\theta} | \underset{\text{Data}}{d}, \underset{\text{Model}}{M}) = \frac{p(d|\theta, M)p(\theta|M)}{p(d|M)}$$

WHAT IS DISCOVERY: WHAT IS BAYES?

Bayes/Laplace told us that

$$p(\overset{\text{Parameters}}{\theta} | \underset{\text{Data}}{d}, \underset{\text{Model}}{M}) = \frac{p(d|\theta, M) \overset{\text{prior}}{p(\theta|M)}}{p(d|M)}$$

The **prior** is **model dependent** and encodes our belief about the parameters **before seeing the data**



Mandau & Dickinson 2014

WHAT IS DISCOVERY: WHAT IS BAYES?


Bayes/Laplace told us that

Parameters

$$p(\theta | d, M) = \frac{p(d|\theta, M)p(\theta|M)}{p(d|M)}$$

Data Model


The prior is **model dependent** and encodes our belief about the parameters before seeing the data



Mandau & Dickinson 2014


The **likelihood** is **model dependent** and encodes our belief about the **data generating process**

Signal model



SXS Collaboration 2025


Noise Model



LVK Collaboration 2020


GW200208_22

This is a ~66 solar mass BBH with 14 cycles in band.
Bayes factor ~ 17, 3, — (uniform, log-uni, expert prior)



GW200208_22

This is a ~66 solar mass BBH with 14 cycles in band.
Bayes factor ~ 17, 3, — (uniform, log-uni, expert prior)




BUT IS IT AN EVENT?

LVK 2021 found that the FAR is 5 per year. The p-astro is 70%.

GW200208_22


This is a ~66 solar mass BBH with 14 cycles in band.
Bayes factor ~ 17, 3, — (uniform, log-uni, expert prior)



BUT IS IT AN EVENT?


LVK 2021 found that the FAR is 5 per year. The p-astro is 70%.

Wang & Nitz 2025 constructed an eccentric template bank with pycbc and found that the FAR is 50 per year. So even worse!



GW200208_22


This is a ~66 solar mass BBH with 14 cycles in band.
Bayes factor ~ 17, 3, — (uniform, log-uni, expert prior)



BUT IS IT AN EVENT? PROBABLY NOT.

LVK 2021 found that the FAR is 5 per year. The p-astro is 70%.



Wang & Nitz 2025 constructed an eccentric template bank with pycbc and found that the FAR is 50 per year. So even worse!



Wang & Nitz 2025



AN ASIDE ON DEGENERACY & WAVEFORM LENGTH

The Bayes factors on the previous slides compared the **eccentric aligned-spin vs the quasi-circular spin-precessing hypothesis**. (Bustillo+ 2021, Romero-Shaw+2023, Divyajyoti+ 2025, Tibrewal+ 2026).



AN ASIDE ON DEGENERACY & WAVEFORM LENGTH

The Bayes factors on the previous slides compared the **eccentric aligned-spin vs the quasi-circular spin-precessing hypothesis**. (Bustillo+ 2021, Romero-Shaw+2023, Divyajyoti+ 2025, Tibrewal+ 2026).




This is a statement on $p(\text{confusion} \mid \text{number of cycles})$. It is **not** a statement about $p(\text{confusion} \mid \text{number of cycles, data})$.

Consequence: if two observations have the same Bayes factor but one is shorter it does not mean that the longer one is more trustworthy.


GW200105 – AN NSBH?

It was reported that there were signatures of eccentricity in a NSBH event GW200105. Note pyEFPE (Morras+ 2025) is inspiral only but **includes eccentricity and spin-precession!**



GW200105 – AN NSBH?

It was reported that there were signatures of eccentricity in a NSBH event GW200105. Note pyEFPE (Morras+ 2025) is inspiral only but **includes eccentricity and spin-precession!**



IS IT AN EVENT?


Criticism of this event was that GW200105 was reported as a low-significance single detector trigger (**FAR of 1 per 2.8 years**, (LVK 2021).

Davies & Harry 2022 showed with updated methods the **FAR is 1 per 50,000 yrs**. Phukon+ 2025 further showed that eccentric templates also decrease the FAR. The glitch also has negligible impact (Morrass+ 2025)

GW200105 – AN NSBH?

It was reported that there were signatures of eccentricity in a NSBH event GW200105. Note pyEFPE (Morras+ 2025) is inspiral only but **includes eccentricity and spin-precession!**


Bayes factor ~ 17, 1.3, __ (uniform, log-uni, expert prior).
See also Tiwari+ 2025




IS IT AN EVENT?

Criticism of this event was that GW200105 was reported as a low-significance single detector trigger (**FAR of 1 per 2.8 years**, (LVK 2021).


Davies & Harry 2022 showed with updated methods the **FAR is 1 per 50,000 yrs**. Phukon+ 2025 further showed that eccentric templates also decrease the FAR. The **glitch also has negligible impact** (Morras+ 2025)



GW190701




Orbital trajectory




There was a scattered light glitch in LIGO Livingston during the detection. Therefore, the “BayesWave” (Cornish+ 2015) glitch mitigation algorithm was employed to subtract the glitch.

GW190701



Orbital trajectory




> This is a heavy BBH with a source frame mass of 140


> There are waveform systematics and sampler systematics

> This event had a high Bayes factor 407, 51, — (uniform, log-uni, expert prior)

GW190701



Orbital trajectory




> This is a heavy BBH with a source frame mass of 140

> There are waveform systematics and sampler systematics

> This event had a high Bayes factor 407, 51, — (uniform, log-uni, expert prior)

There was a scattered light glitch in LIGO Livingston during the detection. Therefore, the “BayesWave” (Cornish+ 2015) glitch mitigation algorithm was employed to subtract the glitch.


HOW DOES GLITCH MITIGATION WORK?



LVK Collaboration 2017


HOW DOES GLITCH MITIGATION WORK?

- > How does “BayesWave” work?
- > Fit the signal coherently with a waveform model
- > Fit the glitch incoherently with sine-Gaussian wavelets
- > **Note this is not waveform model agnostic.**




Payne+ 2022

GW190701





GW190701 - GLITCH

GW190701



GW190701 - GLITCH



GW190701 - GLITCH



- > Just using one draw of the BayesWave glitch mitigation can massively differ compared to marginalizing over the glitch uncertainty
- > The majority of the eccentric evidence comes from L1.
- > To what extent is the glitch fitting the eccentricity and the eccentricity fitting the glitch?

GW200129

> This is a 70 solar mass BBH. The Bayes factors are 10,000, 1,698, ___ (uniform prior, log uniform prior, expert prior). There is also a glitch in this event!


GW200129 — prior: uniform-e

- Gupte+ 2024 (SEOBNRv4EHM, DINGO-IS)
- Planas+ 2025 (IMRPhenomTEHM, bilby)
- Malagon+ 2026 (SEOBNRv5EHM, RIFT)
- Malagon+ 2026 (TEOBResumS-Dali, RIFT)
- Pompili+ 2026 (SEOBNRv6EHM, bilby)




HOW DOES GLITCH MITIGATION WORK?

- > How does “BayesWave” work?
- > Fit the signal coherently with a waveform model
- > Fit the glitch incoherently with sine-Gaussian wavelets
- > **Note this is not waveform model agnostic.**




Adapated from Allen+1998




Payne+ 2022



- > Use auxillary channels to deglitch (Allen+ 1999, Davis+ 2022, Macas+ 2024)?
- > **Note, this is waveform model agnostic.** It uses more information, but it doesn't output uncertainty.





H1



L1








GW200129


In a recent study Gamboa+ in prep. showed that when analyzed with a **eccentric and spin-precessing waveform model** (SEOBNRv6EPHM) GW200129 **gains even more evidence for eccentricity, but does not show evidence for precession.**

GW200129

In a recent study Gamboa+ in prep. showed that when analyzed with a **eccentric and spin-precessing waveform model** (SEOBNRv6EPHM) GW200129 **gains even more evidence for eccentricity, but does not show evidence for precession.**




Gamboa+ in prep.



Gamboa+ in prep.


GW190521

> This is a ~146 solar mass BBH.
Bayes factors of 21 (log-uniform prior) from Romero-Shaw+ 2020



Romero-Shaw+ 2020

> There was also a study using NR simulations to compare the eccentric likelihood




Gayathri+ 2020


GW190521

> Problem is, subsequent papers were did not find eccentricity in GW190521 (Iglesias+ 2022, Ramos-Buades+ 2023, Gupte+2024, Planas+ 2025, Malagon+ 2026)

> Note, the waveform models used in these papers circularize towards merger




Ramos-Buades+ 2023



Iglesias+ 2022

GW190521 - HYPERBOLIC?

> Gamba+ 2021 proposed that GW190521 was a dynamical capture of two non-spinning black holes with Bayes factors of 4315




Gamba+ 2021


GW190521 - HYPERBOLIC?

> Gamba+ 2021 proposed that GW190521 was a **dynamical capture of two non-spinning black holes** with Bayes factors of 4315

> Recent works show that when using updated waveform models the Bayes factors become 4 (Pompili+ 2026 SEOBNRv6EHM) and 8 (Lange+ 2026 TEOBResumS-DALI)



Gamba+ 2021




Pompili+ 2026

> These papers sample in the energy and angular momentum at infinity. This allows one to see the posterior compared to a fiducial separatrix

PLENTY OF OTHER EVENTS, THOUGH NONE REACH HIGH SIGNIFICANCE


> Three independent catalogs (Xu+ 2025, Gupte+ 2026, Malagon+ 2026) have confirmed O4a has no promising eccentric event



Xu+ 2025

PLENTY OF OTHER EVENTS, THOUGH NONE REACH HIGH SIGNIFICANCE

> Three independent catalogs (Xu+ 2025, Gupte+ 2026, Malagon+ 2026) have confirmed O4a has no promising eccentric event



Xu+ 2025

> That's not to say these events are unimportant! They still constrain the prior

> A few ones with “mild” eccentricity support are GW190620, GW191109, GW231223, GW230706, GW230712.

Event name	Waveform Systematics	Noise Systematics	Sampler/Settings Systematics	Search Significance	Parameter Estimation Significance	Overall Significance
GW190701	High	High	Medium	High	Medium	Low
GW190521	Medium	Low	Medium	High	Low	Low
GW200208_22	Low	Low	Low	Low	Low	Low
GW200129	Low	Medium	Low	High	High	High
GW200105	Low	Low	Low	High	Low	Low


BUT WHAT IF OUR PRIORS ARE ALL WRONG?

Famously, before we started taking GW measurements, the expectation was that the majority of detections would be BNS not BBHs. So we need some way of updating our priors based off the data.

BUT WHAT IF OUR PRIORS ARE ALL WRONG?

Famously, before we started taking GW measurements, the expectation was that the majority of detections would be BNS not BBHs. So we need some way of updating our priors based off the data.

Step 1: Update our priors about the population using all the GW data




This [accounts for the look elsewhere effect](#). You are looking at a “global” property of the data. We can compare astrophysical predictions directly to this


BUT WHAT IF OUR PRIORS ARE ALL WRONG?

Famously, before we started taking GW measurements, the expectation was that the majority of detections would be BNS not BBHs. So we need some way of updating our priors based off the data.

Step 1: Update our priors about the population using all the GW data



Step 2: Update our belief about the individual events




This **accounts for the look elsewhere effect**. You are looking at a “global” property of the data. We can compare astrophysical predictions directly to this

By using data from other events, we can find $p(\text{eccentric} \mid \text{all GW data, our astrophysical priors})$. This **makes mildly eccentric events important** as they affect the population!

STUDIES LOOKING AT THE POPULATION PROPERTIES

Zeeshan+ 2026 defined a mixture between two Gaussians as a population model.


$$\pi(\epsilon) = (1 - \zeta_{\epsilon}) \cdot p(\epsilon \mid 0, \sigma_{\text{circ}}) + \zeta_{\epsilon} \cdot p(\epsilon \mid \mu_{\epsilon}, \sigma_{\epsilon}).$$




STUDIES LOOKING AT THE POPULATION PROPERTIES

Zeeshan+ 2026 defined a mixture between two Gaussians as a population model.

$$\pi(\epsilon) = (1 - \zeta_\epsilon) \cdot p(\epsilon \mid 0, \sigma_{\text{circ}}) + \zeta_\epsilon \cdot p(\epsilon \mid \mu_\epsilon, \sigma_\epsilon).$$




With a FAR cutoff of 1 per year, **<5% of binaries have eccentricity > 0.04**. When not including GW200129 this goes to <2%.




A PHYSICS INFORMED POPULATION MODEL

The distribution of eccentricities expected from dense stellar clusters with a dispersion, σ , due to BH captures




A PHYSICS INFORMED POPULATION MODEL

The distribution of eccentricities expected from dense stellar clusters with a dispersion, σ , due to BH captures



Gupte+ 2026


This implies that **less than 47% of binaries come from BH captures in nuclear star clusters**



Gupte+ 2026

SELECTION EFFECTS DUE TO QUASI-CIRCULAR TEMPLATES

> We do not use eccentric templates when searching for BBHs. This can lead to Malmquist bias.




Gupte+ 2026, Vijaykumar+ 2026

> The above plot was produced with a semi-analytical method that estimates sensitivity without using search pipelines.


SELECTION EFFECTS DUE TO QUASI-CIRCULAR TEMPLATES

> We do not use eccentric templates when searching for BBHs. This can lead to Malmquist bias.



> There are search papers that quantify our sensitivity to eccentricity

> Note Wang & Nitz 2025 did an eccentric search on O3 data but found no new candidates




Bhaumik+ 2024

WE COULD ALSO INCORPORATE THE POPULATION INFORMATION VIA PRIOR ODDS

We can “balance” the look elsewhere effect by obtaining a global property of the data. First, we can ask the probability that one event is eccentric.

$$p_{\text{eccentric}} = \frac{\mathcal{O}_{E/QC}}{1 + \mathcal{O}_{E/QC}} = \frac{R_E p(d|M_E)}{R_E p(d|M_E) + R_Q p(d|M_Q)}$$

We can then use all the GW data to obtain a posterior over the rates and marginalize our probability over this.




WE COULD ALSO INCORPORATE THE POPULATION INFORMATION VIA PRIOR ODDS

We can “balance” the look elsewhere effect by obtaining a global property of the data. First, we can ask the probability that one event is eccentric.

$$p_{\text{eccentric}} = \frac{\mathcal{O}_{E/QC}}{1 + \mathcal{O}_{E/QC}} = \frac{R_E p(d|M_E)}{R_E p(d|M_E) + R_Q p(d|M_Q)}$$

We can then use all the GW data to obtain a posterior over the rates and marginalize our probability over this.

Finally, we can use the same formalism to ask what the probability is that an event of the 57 is eccentric. This comes out to ~99%.



Gupte+ 2026

CONCLUSIONS

- > There is likely eccentricity in the current LVK data in at least one event

CONCLUSIONS

- > There is likely eccentricity in the current LVK data in at least one event
- > We should compare eccentric population observations with the astrophysics community (I'm sure will be many talks about this)

CONCLUSIONS

- > There is likely eccentricity in the current LVK data in at least one event
- > We should compare eccentric population observations with the astrophysics community (I'm sure will be many talks about this)
- > We **need** sophisticated **methods to account for non-Gaussian noise**. Simulation-based inference can be very useful here (Raymond+ 2024)

CONCLUSIONS

- > There is likely eccentricity in the current LVK data in at least one event
- > We should compare eccentric population observations with the astrophysics community (I'm sure will be many talks about this)
- > We **need** sophisticated **methods to account for non-Gaussian noise**. Simulation-based inference can be very useful here (Raymond+ 2024)
- > **Modelling eccentricity effects near merger is important**. Just because eccentricity is expected to approach zero near merger does not mean it will for all systems!

CONCLUSIONS

- > There is likely eccentricity in the current LVK data in at least one event
- > We should compare eccentric population observations with the astrophysics community (I'm sure will be many talks about this)
- > We **need** sophisticated **methods to account for non-Gaussian noise**. Simulation-based inference can be very useful here (Raymond+ 2024)
- > **Modelling eccentricity effects near merger is important**. Just because eccentricity is expected to approach zero near merger does not mean it will for all systems!
- > **Priors are not scary! We just need to choose a good one** (see recent work by Rozner+ 2026)