

# NMST - HOME ASSIGNMENT-6

Name: Kashish Singh

Stdt. Group: EC2

Reg. no.: 20195082

① a) We know that, Newton's forward difference interpolation formula is

$$y = y_0 + \mu \Delta y_0 + \frac{\mu(\mu-1)}{2!} \Delta^2 y_0 + \frac{\mu(\mu-1)(\mu-2)}{3!} \Delta^3 y_0 + \dots$$

where,  $\mu = \frac{x-a}{h}$

differentiating eqn. ① w.r.t.  $x$ :

$$\frac{dy}{dx} = \Delta y_0 + \frac{2\mu-1}{2} \Delta^2 y_0 + \frac{3\mu^2-6\mu+2}{6} \Delta^3 y_0 + \dots \quad (3)$$

differentiating eqn. (2) w.r.t.  $x$ :

$$\frac{du}{dx} = \frac{1}{h}$$

$$\text{So, } \frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} = \frac{1}{h} \left( \Delta y_0 + \frac{2\mu-1}{2} \Delta^2 y_0 + \frac{3\mu^2-6\mu+2}{6} \Delta^3 y_0 + \dots \right)$$

at  $x = a$ ,

$\mu = 0$  from eqn ②.

$$\text{So, } \frac{dy}{dx} = \frac{1}{h} \left( \Delta y_0 - \frac{\Delta^2 y_0}{2} + \frac{\Delta^3 y_0}{3} - \dots \right) \quad \text{Ans}$$

b)  $\frac{d^2 y}{dx^2} = \frac{1}{h^2} \left( \Delta^2 y_0 - \Delta^3 y_0 + \frac{11}{12} \Delta^4 y_0 + \dots \right)$

$$\Rightarrow \frac{d^2 y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d}{du} \left( \frac{dy}{dx} \times \frac{du}{dx} \right)$$

$$\begin{aligned}
 &= \frac{d}{du} \left[ \frac{1}{h} \left( \Delta y_0 + \left( \frac{2u-1}{2} \right) \Delta^2 y_0 + \frac{3u^2-6u+2}{6} \Delta^3 y_0 + \dots \right) \right] \times \frac{1}{h} \\
 &= \frac{1}{h} \left[ \Delta^2 y_0 + (u-1) \Delta^3 y_0 + \left( \frac{6u^2-18u+11}{12} \right) \Delta^4 y_0 + \dots \right] \times \frac{1}{h} \\
 &= \frac{1}{h^2} \left[ \Delta^2 y_0 + (u-1) \Delta^3 y_0 + \left( \frac{6u^2-18u+11}{12} \right) \Delta^4 y_0 + \dots \right]
 \end{aligned}$$

at  $x=a$ ,  
 $u=0$

$$\frac{d^2y}{dx^2} = \frac{1}{h^2} \left[ \Delta^2 y_0 - \Delta^3 y_0 + \frac{11}{12} \Delta^4 y_0 + \dots \right]$$

$$c) \left( \frac{d^3y}{dx^3} \right)_{x=a} = \frac{1}{h^3} \left[ \Delta^3 y_0 - \frac{3}{2} \Delta^4 y_0 + \dots \right]$$

$$\Rightarrow \left( \frac{d^3y}{dx^3} \right) = \frac{d}{dx} \left( \frac{d^2y}{dx^2} \right)$$

$$= \frac{d}{du} \left( \frac{d^2y}{dx^2} \right) \times \frac{du}{dx}$$

$$\begin{aligned}
 \frac{d^2y}{dx^2} &= \frac{1}{h^2} \left[ \Delta^2 y_0 + \left( \frac{12u-18}{12} \right) \Delta^3 y_0 + \dots \right] \times \frac{1}{h} \\
 &= \frac{1}{h^3} \left[ \Delta^3 y_0 + \frac{2u-3}{2} \Delta^4 y_0 + \dots \right]
 \end{aligned}$$

at  $x=a$ ,  
 $u=0$ .

$$\frac{d^3y}{dx^3} = \frac{1}{h^3} \left[ \Delta^3 y_0 - \frac{3}{2} \Delta^4 y_0 \right]$$

Aus

(a) We know that, Newton's backward difference formula is :-

$$y = y_m + u \nabla y_m + \frac{u(u+1)}{2!} \nabla^2 y_m + \frac{(u)(u+1)(u+2)}{3!} \nabla^3 y_m \quad \text{--- (1)}$$

$$u = \frac{x - x_m}{h} \quad \dots \quad (2)$$

Differentiating eqn (1) w.r.t.  $u$ , we get ...

$$\frac{dy}{du} = \nabla y_m + \frac{(2u+1)}{2!} \nabla^2 y_m + \frac{(3u^2+6u+2)}{3!} \nabla^3 y_m.$$

$$= \nabla y_m + \frac{(2u+1)}{2} \nabla^2 y_m + \frac{(3u^2+6u+2)}{3!} \nabla^3 y_m \quad \text{--- (2)}$$

$$\frac{du}{dx} = \frac{1}{h}$$

$$\text{Now, } \frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

$$\frac{dy}{dx} = \frac{1}{h} \left( \nabla y_m + \frac{\nabla^2 y_m + (2u+1)}{2} + \frac{(3u^2+6u+2)}{3!} \nabla^3 y_m \right)$$

at  $x = x_m$ , we have  $u = 0$ .

$\therefore$  putting  $u = 0$ , we get -

$$\left( \frac{dy}{dx} \right)_{x=x_m} = \frac{1}{h} \left[ \nabla y_m + \frac{1}{2} \nabla^2 y_m + \frac{1}{3} \nabla^3 y_m + \frac{1}{4!} \nabla^4 y_m + \dots \right] \quad \text{Ans.}$$

b) Again diff. eqn. (1), we get:-

$$\left( \frac{d^2 y}{dx^2} \right)_{x=x_m} = \frac{1}{h^2} \left[ \nabla^2 y_m + \nabla^3 y_m + \frac{11}{12} \nabla^4 y_m + \dots \right]$$

$$\Rightarrow \frac{d^2y}{dx^2} = \frac{d}{du} \left( \frac{dy}{du} \right) \frac{du}{dn}$$

$$= \frac{1}{h^2} \left[ \nabla^2 y_n + (u+1) \nabla^3 y_n + \left( \frac{6u^2 + 18u + 1}{12} \right) \nabla^4 y_n + \dots \right]$$

at  $n = m$ ,  $u = 0$ , we get:—

$$\therefore \left( \frac{d^2y}{dx^2} \right)_{x=x_m} = \frac{1}{h^2} \left[ \nabla^2 y_n + \nabla^3 y_n + \frac{11}{12} \nabla^4 y_n + \dots \right]$$

$$C) \therefore \frac{d^3y}{dx^3} = \frac{d}{du} \left( \frac{d^2y}{dx^2} \right) \frac{du}{dn}.$$

$$= \frac{1}{h^3} \left[ \frac{d}{du} (\nabla^2 y_n) + \frac{d}{du} (u+1) \nabla^3 y_n + \frac{d}{du} \left( \frac{6u^2 + 18u + 11}{12} \right) \nabla^4 y_n + \dots \right] \frac{1}{h}$$

$$\therefore \left( \frac{du}{dx} = \frac{1}{h} \right)$$

$$\therefore \frac{d^3y}{dx^3} = \frac{1}{h^3} \left( 0 + \nabla^3 y_n + \left( u + \frac{3}{2} \right) \nabla^4 y_n + \dots \right)$$

putting  $u = 0$  at  $n = m$ , we get:—

$$\left( \frac{d^3y}{dx^3} \right)_{x=x_m} = \frac{1}{h^3} \left[ \nabla^3 y_n + \frac{3}{2} \nabla^4 y_n + \dots \right]$$

Aus

(3) (a) Stirling central difference formula:-

$$y = y_0 + u \left( \frac{\Delta y_0 + \Delta y_1}{2} \right) + \frac{u^2}{2!} (\nabla^2 y_{-1}) + \frac{u(u^2 - 1)}{3!} \\ \left( \frac{\Delta^3 y_{-1} + \Delta^3 y_{-2}}{2} \right) + \frac{u^2(u^2 - 1^2)}{4!} \nabla^4 y_{-2} + \dots \quad (1)$$

where  $u = \frac{x-a}{h}$   $\rightarrow$  (2)

Differentiating eqn (1) w.r.t.  $u$ , we get:-

$$\frac{dy}{du} = \frac{\Delta y_0 + \Delta y_1}{2} + u \cdot \Delta^2 y_{-1} + \frac{(3u^2 - 1)}{6} \left( \frac{\Delta^3 y_{-1} + \Delta^3 y_{-2}}{2} \right) \dots \quad (2)$$

Differentiating eqn. (2) w.r.t.  $x$  ;—

$$\frac{du}{dx} = \frac{1}{h} \rightarrow (4)$$

$$\therefore \frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} \\ = \frac{1}{h} \left[ \frac{\Delta y_0 + \Delta y_1}{2} + u \cdot \Delta^2 y_{-1} + \frac{(3u^2 - 1)}{6} \left( \frac{\Delta^3 y_{-1} + \Delta^3 y_{-2}}{2} \right) \right. \\ + \left. \left( \frac{4u^2 - 2u}{4!} \right) \nabla^4 y_{-2} + \left( \frac{5u^4 - 15u^2 + 4}{5!} \right) \right. \\ \left. \left( \frac{\Delta^5 y_{-2} + \Delta^5 y_{-3}}{2!} \right) + \dots \right] \dots \quad (5)$$

when  $x = a$ ,  $u = 0$ .

$$\therefore \left( \frac{dy}{dx} \right)_{x=a} = \frac{1}{h} \left[ \frac{\Delta y_0 + \Delta y_1}{2} + \left( \frac{1}{6} \right) \left( \frac{\Delta^3 y_{-1} + \Delta^3 y_{-2}}{2} \right) + \dots \right]$$

∴ Proved.

b) Differentiating (3) wrt.  $n$  :-

$$\frac{d^2y}{dx^2} = \frac{d}{du} \left( \frac{dy}{du} \right) \frac{du}{dn}.$$

$$= \frac{1}{h^2} \left[ \Delta^2 y_1 + u \frac{(\Delta^3 y_{-1} + \Delta^3 y_{-2})}{2} + \frac{(6u^2 - 1)}{12} \Delta^4 y_{-2} \right. \\ \left. + \frac{(2u^3 - 3u)}{12} \left( \Delta^5 y_{-2} + \Delta^5 y_{-3} \right) + \dots \right]$$

putting  $u=0$ , at  $x=a$  in equation (3),

$$\left( \frac{d^2y}{dx^2} \right)_{x=a} = \frac{1}{h^2} \left( \Delta^2 y_1 - \frac{1}{12} \Delta^4 y_{-2} + \frac{1}{1920} \Delta^6 y_{-3} + \dots \right)$$

Ans.

(4) a) Bessel's central difference formula is :-

$$y = \frac{(y_0 - y_1)}{2} + (u - \frac{1}{2}) \Delta y_0 + \frac{u(u-1)}{2!} \left( \frac{\Delta^2 y_{-1} + \Delta^2 y_0}{2} \right) \\ + \frac{u(u-1)(u-1/2)}{3!} \Delta^3 y_{-1} + \dots \rightarrow ①$$

$$\text{where, } u = \frac{x-a}{h} \rightarrow ②$$

Differentiating eq. ① wrt.  $u$ , we get:-

$$\frac{dy}{du} = \Delta y_0 + \frac{(2u-1)}{2!} \left( \frac{\Delta^2 y_{-1} + \Delta^2 y_0}{2} \right) + \frac{(3u^2 - 3u + 1/2)}{3!}$$

$$\Delta^3 y_{-1} + \dots - ③$$

Differentiating with eqn ② wrt.  $n$  :-

$$\frac{du}{dx} = \frac{1}{h}$$

$$\therefore \frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

$$= \frac{1}{h} \left[ \Delta y_0 + \frac{(2u-1)}{2} \left( \frac{\Delta^2 y_1 + \Delta^2 y_0}{2} \right) + \left( \frac{3u^2 - 3u + 1/2}{3!} \right) \right]$$

$$\Delta^3 y_{-1} + \left( \frac{4u^3 - 6u^2 - 2u + 2}{4!} \right) \left( \frac{\Delta^4 y_{-2} + \Delta^4 y_{-1}}{2} \right) + \dots \right] \quad (5)$$

putting,  $u=0$ , at  $x=a$ , in eqn. (5) :-

$$\left( \frac{dy}{dx} \right)_{x=a} = \frac{1}{h} \left[ \Delta y_0 + \left( \frac{1}{2} \right) \left( \frac{\Delta^2 y_1 + \Delta^2 y_0}{2} \right) + \frac{1}{12} \Delta^3 y_{-1} + \right.$$

$$\left. + \frac{1}{12} \left( \frac{\Delta^4 y_{-2} + \Delta^4 y_{-1}}{2} \right) - \frac{1}{120} \Delta^5 y_{-2} + \dots \right] \quad (6)$$

Proved Ahs

b) Differentiating eqn. (6) w.r.t x, we get:-

$$\frac{d^2 y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d}{dy} \left( \frac{dy}{dx} \right) \frac{du}{dx}$$

$$= \frac{1}{h^2} \left[ \left( \frac{\Delta^2 y_1 + \Delta^2 y_0}{2} \right) + \left( \frac{2u-1}{2} \right) \Delta^3 y_1 + \left( \frac{6u^2 - 6u + 1}{12} \right) \right]$$

$$\left( \frac{\Delta^4 y_{-2} + \Delta^4 y_{-1}}{2} \right) + \left( \frac{4u^3 + 6u^2 + 1}{24} \right) \Delta^5 y_{-2} + \dots \right] \quad (7)$$

putting  $u=0$ , at  $x=a$ , in eqn (7), we get:-

$$\left( \frac{d^2 y}{dx^2} \right)_{x=a} = \frac{1}{h^2} \left[ \left( \frac{\Delta^2 y_1 + \Delta^2 y_0}{2} \right) - \frac{1}{2} \Delta^3 y_1 - \frac{1}{12} \left( \Delta^4 y_{-2} + \Delta^4 y_{-1} \right) \right]$$

$$+ \frac{1}{324} \Delta^5 y_{-2} + \frac{1}{90} \left( \frac{\Delta^6 y_{-3} + \Delta^6 y_{-2}}{2} \right) + \dots \right]$$

Hence Proved. Ahs

⑤ The difference table:-

$t$	$v$	$\Delta v$	$\Delta^2 v$	$\Delta^3 v$	$\Delta^4 v$
0	0	3			
5	3	11	8	36	
10	14	55	44	60	24
15	69		104		
20	228	159			

Acceleration at  $t=0$ , is given by:-

$$\left(\frac{dv}{dy}\right)_{t=0} = \frac{1}{h} \left[ \Delta v_0 - \frac{1}{2} \Delta^2 v_0 + \frac{1}{3} \Delta^3 v_0 - \frac{1}{4} \Delta^4 v_0 \right]$$

$$= \frac{1}{5} \left[ 3 \times \frac{1}{2} \times 8 + \frac{1}{3} \times 36 - \frac{1}{4} \times 24 \right]$$

$$= \frac{1}{5} [3 - 4 + 12 - 6] = \frac{1}{5} \times (15 - 10)$$

$$= 1.$$

Ans

$\therefore$  Acceleration =  $\frac{1 \text{ m/s}^2}{\text{m}}$ .

$x$	$y$	$\Delta y$	$\Delta^2 y$	$\Delta^3 y$	$\Delta^4 y$	$\Delta^5 y$
0.2	0.9182					
0.3	0.8975	-0.0207		0.0105		-0.0014

0.4	0.8873	0.0091	0.0007
		-0.0011	-0.0007
0.5	0.8862	0.0078	

0.0151

0.7 0.9086

using Newton's forward difference formula of differentiation, we get:-

$$\frac{dy}{du} = \Delta y_0 + \frac{2u-1}{2!} \Delta^2 y_0 + \frac{3u^2 - 6u + 2}{3!} \Delta^3 y_0 + \dots \quad (1)$$

for maxima or minima,

$$\frac{dy}{du} = c, \text{ for maxima}, \frac{d^2y}{du^2} = -ve.$$

$$\text{for minima, } \frac{d^2y}{du^2} = +ve.$$

$$\frac{dy}{du} = -0.0207 + \frac{(2u-1)}{2} \times 0.0105 + \frac{(3u^2 - 6u + 2) \times (-0.0011)}{3!}$$

$$= 207 + \frac{(2u-1) \times 105}{2} + \frac{(3u^2 - 6u + 2) \times -14}{3!} = 0.$$

$$= 42u^2 - 714u + (-899) = 0.$$

$$u_1 = 18.178, \quad u_2 = -1.1775.$$

$$\frac{d^2y}{du^2} = 105 + (u-1)(-14)$$

$$\left( \frac{d^2y}{du^2} \right)_{u=18.178} = \frac{-ve.}{}$$

$$\left( \frac{d^2y}{du^2} \right)_{u=-1.1775} = +ve.$$

$\therefore y$  is minimum at  $x = -1.1775$ .

$$\Rightarrow x = 0.2 - 1.1775 \times 0.1.$$

$$\therefore x = 0.08225.$$

$$\Rightarrow y = y_0 + x \Delta y_0 + \frac{x_0(x_0-1)}{2!} \Delta^2 y_0 + \frac{x_0(x_0-1)(x_0-2)}{3!} \Delta^3 y_0$$

$$\therefore y_{\min} = 0.9157.$$

Aus

3!

④ Let,  $I = \int_a^b y \cdot dx$ , where  $y$  takes the values  $y_0, y_1, y_2, \dots, y_n$  for  $x = x_0, x_1, x_2, \dots, x_n$ .

Let the interval of integration  $(a, b)$  be divided into  $n$  equal sub-intervals, each of width  $h$ .

where,  $h = \frac{b-a}{n}$ , so that

$$x_0 = a, x_1 = x_0 + h, x_2 = x_0 + 2h, \dots, x_n = x_0 + nh = b.$$

$$\Rightarrow I = \int_{x_0}^{x_0+nh} f(x) \cdot dx.$$

$\therefore$  any  $x$  is given by  $x = x_0 + sh$  &  $dx = h \cdot dx$

$$\therefore I = \int_0^n f(x_0 + sh) dx.$$

$$I = h \int_0^n \left( y_0 + \gamma \Delta y_0 + \frac{\gamma(\gamma-1)}{2} \Delta^2 y_0 + \frac{\gamma(\gamma-1)(\gamma+2)}{3} \Delta^3 y_0 + \dots \right) d\gamma.$$

(by Newton - forward interpolation formula).

$$= h \left[ y_0 + \frac{\gamma^2}{2} \Delta y_0 + \frac{1}{2} \left( \frac{\gamma^3}{3} - \frac{\gamma^3}{2} \right) \Delta^2 y_0 + \frac{1}{6} \left( \frac{\gamma^4}{4} - \gamma^3 + \gamma^2 \right) \Delta^3 y_0 + \dots \right]_n \\ = nh \left[ y_0 + \frac{n}{2} \Delta y_0 + \frac{n(2n-3)}{12} \Delta^2 y_0 + \frac{6n(n-1)^2}{24} \Delta^3 y_0 + \dots \right] \rightarrow ①$$

This is ~~the~~ general quadratic formula & is known as ~~Newton~~ - Cote's quadratic formula.

a) putting  $n=1$  in eqn. ① & taking the curve through  $(x_0, y_0)$  &  $(x_1, y_1)$  as a polynomial of degree one, so that differences of an order higher than one vanishes.

$$\int_{x_0}^{x_0+h} f(x) dx = h (y_0 + \frac{1}{2} \Delta y_0) \\ = \frac{h}{2} (2y_0 + (y_1 - y_0)) \\ = \frac{h}{2} (y_0 + y_1)$$

Similarly, for the next sub interval  
 $(x_0 + h, x_0 + 2h)$

$$\int_{x_0+h}^{x_0+2h} f(x) dx = \frac{h}{2} (y_1 + y_2)$$

$$\int_{x_0+(n-1)h}^{x_0+nh} f(x) dx = \frac{h}{2} (y_{n-1} + y_n).$$

Adding all the above equations; -

$$\int_{x_0}^{x_0+nh} f(x) dx = \frac{h}{2} [y_0 + 2(y_1 + y_2 + y_3 + \dots + y_{n-1}) + y_n]$$

$\therefore$  This is Trapezoidal rule.

b) Putting  $n=2$  in formula ① and taking the curve through  $(x_0, y_0)$ ,  $(x_1, y_1)$  &  $(x_2, y_2)$  as a polynomial of degree two so that the differences ~~are~~ of order higher than two vanishes, we get

$$\int_{x_0}^{x_0+2h} f(x) dx = 2h \left[ y_0 + \Delta y_0 + \frac{1}{6} D^2 y_0 \right]$$

$$= \frac{2h}{2} [Gy_0 + 6(y_1 - y_0) + (y_2 - 2y_1 + y_0)] \\ = \frac{h}{3} [y_0 + 4y_1 + y_2]$$

Similarly,

$$\int_{x_0+2h}^{x_0+4h} f(x) dx = \frac{h}{3} [y_2 + 4y_3 + y_4]$$

$$\int_{x_0+(n-2)h}^{x_0+nh} f(x) dx = \frac{h}{3} [y_{n-2} + 4y_{n-1} + y_n]$$

Adding all the above & equo. —

$$\int_{x_0}^{x_0+nh} f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) \\ + 2(y_2 + y_4 + \dots + y_{n-2})]$$

This is Simpson's one-third rule

C) Putting  $n=3$  in formula ① & taking the curve through  $(x_0, y_0), (x_1, y_1), (x_2, y_2)$  &  $(x_3, y_3)$  as a polynomial of degree  $\leq$  that differences of order higher than three vanishes, we get :-

$$\int_{x_0}^{x_0+3h} f(x) dx = 3h \left( y_0 + \frac{3}{2} \Delta y_0 + \frac{3}{4} \Delta^2 y_0 + \frac{1}{8} \Delta^3 y_0 \right)$$

$$= \frac{3h}{8} \left( 8y_0 + 12(y_1 - y_0) + 6(y_2 - 2y_1 + y_0) + (y_3 - 3y_2 + 3y_1 - y_0) \right)$$

$$= \frac{3h}{8} [y_0 + 3y_1 + 3y_2 + y_3]$$

Similarly,

$$\int_{x_0+3h}^{x_0+6h} f(x) \cdot dx = \frac{3h}{8} [y_3 + 3y_4 + 3y_5 + y_6]$$

$x_0+3h$ ,

$$\int_{x_0+(m-3)h}^{x_0+6h} f(x) \cdot dx = \frac{3h}{8} [y_{n-3} + 3y_{n-2} + 3y_{n-1} + y_n]$$

Adding the above integration,

$$\int_{x_0}^{x_0+nh} f(x) \cdot dx = \frac{3h}{8} [(y_0 + y_n) + 3(y_1 + y_2 + y_3 + y_4 + \dots + y_{n-2} + y_{n-1}) + 2(y_5 + y_6 + \dots + y_{n-3})]$$

$\therefore$  This is Dompson's three eight rule

d) Putting  $n=4$ , in formula ① and neglecting all differences of orders higher than 4, we get:-

$$\int_{x_0}^{x_0+4h} f(x) dx = h \int_0^4 [y_0 + \frac{\gamma}{1!} \Delta y_0 + \frac{\gamma(\gamma-1)}{2!} \Delta^2 y_0 + \frac{\gamma(\gamma-1)(\gamma-2)}{3!} \Delta^3 y_0 + \frac{\gamma(\gamma-1)(\gamma-2)(\gamma-3)}{4!} \Delta^4 y_0] d\gamma$$

By newton's forward interpolation

formula:-

$$= 4h \left[ y_0 + \frac{m}{2} \Delta y_0 + \frac{m(8m-3)}{12} \Delta^2 y_0 + \frac{m(m-2)^2}{24} \Delta^3 y_0 + \left( \frac{m^4}{5} - \frac{3m^3}{2} + \frac{11m^2}{3} - 3m \right) \frac{\Delta^4 y_0}{4!} \right]$$

$$= 4h \left[ y_0 + 2\Delta y_0 + \frac{5}{3} \Delta^2 y_0 + \frac{3}{2} \Delta^3 y_0 + \frac{7}{50} \Delta^4 y_0 \right]$$

$$= \frac{2h}{45} [7y_0 + 32y_1 + 12y_2 + 32y_3 + 7y_4]$$

Similarly,

$$\int_{x_0+4h}^{x_0+8h} f(x) dx = \frac{2h}{45} [7y_4 + 32y_5 + 12y_6 + 32y_7 + 7y_8]$$

Adding all the above equations, :-

$$\int_{x_0}^{x_0+6h} f(x) \cdot dx = \frac{2h}{45} [7y_0 + 32y_1 + 12y_2 + 32y_3 + 14y_4 + \dots]$$

∴ This is Boole's Rule.

- e) Putting  $n=6$ , in formula ① and neglecting all differences of order higher than six! —

$$\begin{aligned} \int_{x_0}^{x_0+6h} f(x) \cdot dx &= h \int_0^6 [y_0 + \frac{\gamma \Delta y_0 + \gamma(\gamma-1)}{2!} \Delta^2 y_0 + \\ &\quad \frac{\gamma(\gamma-1)(\gamma-2)}{3!} \Delta^3 y_0 + \dots + \\ &\quad \underbrace{\frac{\gamma(\gamma-1)(\gamma-2)(\gamma-3)(\gamma-4)(\gamma-5)}{6!} \Delta^6 y_0}] \\ &= h \left[ \gamma y_0 + \frac{\gamma^2}{2} \Delta y_0 + \frac{1}{2} \left( \frac{\gamma^3}{3} - \frac{\gamma^2}{2} \right) \Delta^2 y_0 + \dots + \right. \\ &\quad + \frac{1}{720} \left( \frac{\gamma^7}{7!} - \frac{5\gamma^6}{2} + \frac{17\gamma^5}{2} \right. \\ &\quad \left. - \frac{225}{4} \gamma^4 + \frac{274}{3} \gamma^3 - 60\gamma^2 \right] \left. \Delta^6 y_0 \right]_0^6 \end{aligned}$$

$$= \frac{6h}{20} [20y_0 + 60\Delta y_1 + 90\Delta^2 y_2 + 80\Delta^3 y_3 + \\ + 41\Delta^4 y_4 + 11\Delta^5 y_5 + \frac{41}{42}\Delta^6 y_6]$$

$$= \frac{3h}{10} [y_0 + 5y_1 + y_2 + 6y_3 + y_4 + 5y_5 + y_6]$$

Similarly,

$$\int_{x_0+12h}^{x_0+12h} f(x) dx = \frac{3h}{10} [y_8 + 5y_7 + y_9 + 6y_8 + y_{10} + 5y_9 + \\ + y_{12}]$$

$$\int_{x_0+(m-6)h}^{x_0+mh} f(x) dx = \frac{3h}{10} [y_{m-6} + 5y_{m-5} + y_{m-4} + \\ + 6y_{m-3} + 2y_{m-2} + 5y_{m-1} + y_m]$$

Adding all the above equations:-

$$\int_{x_0}^{x_0+mh} f(x) dx = \frac{3h}{10} [y_0 + 5y_1 + y_2 + 6y_3 + y_4 + 5y_5 + \\ + 2y_6 + \dots]$$

$\therefore$  This is Weddle's rule.

Here,  $m$  must be the multiple of 6.

(8) let  $h = 0.2$ ,  $a = 4$ ,  $b = 5.6$ ,  $n = 6$ .  
 $\therefore 5.2 = 4 + 6h$ .  
 $h = \frac{0.2}{\square}$ .

Now,  $x_n = x_0 + nh$  &  $y_n = \log x_n$ .

X	Y
4	1.3862
4.2	1.4350845
4.4	1.4816045
4.6	1.5260563
4.8	1.5686159
5.0	1.6094375
5.2	1.6486586

a) By using Trapezoidal rule:-

$$\int_{4}^{5.2} (\ln x) dx = \frac{h}{2} [y_0 + y_6 + 2(y_1 + y_2 + y_3 + y_4 + y_5)]$$

$$= \frac{0.2}{2} [3.334 + 2 \times 7.320799]$$

$$= 0.1 \bar{x} 18.276551$$

$$\int_{4}^{5.2} (\ln x) dx = 1.8276551.$$

b) By Weddle's rule :-

$$\int_{4}^{5.2} (\ln x) dx = \frac{3h}{10} [y_0 + y_6 + 5(y_1 + y_5) + y_2 + y_4 + 26y_3]$$

$$= \frac{0.6}{10} \times 30.464123$$

$$\therefore \int_4^{5.2} (\ln x) dx = \underline{1.827874}.$$

⑨ While using the Simpson's one third formula, the given interval of integration must be divided into an even number of sub-intervals.

$$\therefore \int_0^4 e^x dx = \frac{h}{3} [Y_0 + Y_n + 4(Y_1 + Y_3 + \dots + Y_{n-1}) + 2(Y_2 + Y_4 + \dots + Y_{n-2})]$$

$$\text{where } h = 1.$$

$$\therefore \int_0^4 e^x dx = \frac{1}{3} [(1+54.6) + 4(2.72+20.09) + (2 \times 7.39)]$$

$$= \frac{1}{3} \times 161.62$$

$$\therefore \int_0^4 e^x dx = 53.87333$$

By actual integration value it is :-

$$\int_0^4 e^x dx = [e^x]_0^4 = e^4 - 1$$

$$= 53.6$$

$\therefore$  Magnitude of error in Simpson's one third rule =  $153.6 - 53.87333$ .

$$= \frac{0.27333}{153.6} \quad \text{Ans}$$

⑩ Taking,  $n=6$  we get  $h=1$ ,

$$x_0 = 0, x_6 = 6 \rightarrow y = \frac{1}{1+x^2}$$

$$x_n = x_0 + n h$$

$$y_n = \frac{1}{1+x_n^2}$$

x.	0	1	2	3	4	5	6
y	1	0.5	0.2	0.1	$\frac{1}{17}$	$\frac{1}{26}$	$\frac{1}{37}$

ii) By Simpson's one third rule:-

$$\begin{aligned} \int_0^6 \frac{dx}{1+x^2} &= \frac{h}{3} [(Y_0 + Y_6) + 4(Y_1 + Y_3 + Y_5) \\ &\quad + 2(Y_2 + Y_4)] \\ &= \frac{1}{3} [(1 + \frac{1}{37}) + 4(0.5 + 0.1 + \frac{1}{26}) \\ &\quad + 2(0.2 + \frac{1}{17})] \\ &= \int_0^6 \frac{dx}{1+x^2} \\ &= \underline{1.366173413} \end{aligned}$$

i) By Trapezoidal rule

$$\begin{aligned} \int_0^6 \frac{dx}{1+x^2} &= \frac{h}{2} [(Y_0 + Y_6) + 2(Y_1 + Y_2 + Y_3 + Y_4 + Y_5)] \\ &= \frac{1}{2} ((1 + \frac{1}{37}) + 2(0.5 + 0.2 + \\ &\quad 0.1 + \frac{1}{17} + \frac{1}{26}) \\ \therefore \int_0^6 \frac{dx}{1+x^2} &= \underline{1.410798581} \text{ Ans.} \end{aligned}$$

By actual integration:-

$$\begin{aligned} \int_0^6 \frac{dx}{1+x^2} &= \tan^{-1} x \Big|_0^6 = \tan^{-1} 6 - \tan^{-1} 0 \\ &= 80.5376^\circ \\ &= \underline{1.40564765 \text{ rad.}} \text{ Ans.} \end{aligned}$$

$$\text{Magnitude of error in Trapezoidal rule: } - \\ = |1.40564765 - 1.410998581|$$

$$= 0.0051509$$

$$\text{Magnitude of error in Simpson's one-third rule: } - \\ = |1.40564765 - 1.366173413|$$

$$= \underline{0.039474} \quad \text{Ans.}$$

$\boxed{\text{Using Simpson's one-third formula,}}$

$$\text{when } n = 4$$

$$\int_{x_0}^{x_0+hn} f(x) dx = \frac{h}{3} [2(y_0 - y_n) + 4(y_1 - y_3) + 2y_2]$$

$$y_n = x_0 + nh, \quad y_n = \frac{1}{5+3n}, \quad h = 0.25.$$

x	1	1.25	1.50	1.75	2.00
y	0.125	0.1142	0.1052	0.0967	0.0809

$$\Rightarrow \int_1^2 \frac{dx}{5+3x} = \frac{1}{3} [(0.125 + 0.0809) + \\ 4(0.1142 + 0.0967) + 2 \times 0.1052] \\ = \frac{1}{12} \times (1.2715)$$

$$\therefore \int_1^2 \frac{dx}{5+3x} = \underline{0.1059}$$

b) when,  $n = 8$

$$h = \frac{2-1}{8} = \frac{1}{8}$$

$$y_n = x_0 + nh, \quad y_n = \frac{1}{5+3x}$$

X	1	1.125	1.25	1.375	1.5	1.625	1.75
Y	0.125	0.1194	0.1142	0.1095	0.1052	0.1012	0.097

X	1.875	2
Y	0.0941	0.0909

$$\therefore \int_1^2 \frac{dx}{5+3x} = \frac{y_8}{3} [y_0 + y_8 + 4(y_1 + y_3 + y_5 + y_7) + 2(y_2 + y_4 + y_6)] \\ = \frac{1}{24} \times 2.5455$$

$$\therefore \int_1^2 \frac{dx}{5+3x} = \underline{0.1060} \text{ Ans.}$$

By Actual Integration,

$$I = \int_1^2 \frac{dx}{5+3x} = \frac{1}{3} (\ln(5+3x))_1^2 \\ = \frac{1}{3} (\ln 11 - \ln 8) \\ = \frac{1}{3} \ln(11/8)$$

$$\therefore I = \underline{0.10615.} \text{ Ans.}$$

Absolute error when  $n=4$ ,

$$= |0.10615 - 0.10591| \\ = 0.00025.$$

Absolute error when  $m=8$ ,

$$= |0.10615 - 0.10601| \\ = \underline{0.00015} \text{ Ans.}$$

(12) Simpson's  $\frac{3}{8}$  formula is:

$$\int_{x_0}^{x_0+nh} f(x) \cdot dx = \frac{3h}{8} [y_0 + y_n + 3(y_1 + y_2 + \dots) + 2(y_3 + y_6 + \dots + y_{n-3})]$$

where,  $h = \frac{b-a}{m}$ ,  $m = m_0$  of intervals.

Case-1: when,  $n = 3$

$$x_m = x_0 + nh, y_m = \frac{1}{5+3x_m}$$

$$h = \frac{1}{3}$$

x	1	1.33	1.66	2
y	0.1250	0.1112	0.1002	0.0909

$$\begin{aligned} \therefore \int_1^2 \frac{dx}{5+3x} &= \frac{3 \times \frac{1}{3}}{8} [0.125 + 0.0909 + 3(0.1112 + \\ &\quad 0.1002)] \\ &= \frac{1}{8} (0.2159 + 0.4338) \\ &= \frac{1}{8} \times 0.8501 \end{aligned}$$

$$\Rightarrow \int_1^2 \frac{dx}{5+3x} = \underline{\underline{0.1062}}$$

Case-2: when  $n = 6$ ,  $h = \frac{2-1}{6} = \frac{1}{6}$

x	1	1.166	1.332	1.498	1.664	1.830	2.0
y	0.125	0.1176	0.1112	0.1053	0.1002	0.095	0.0909

$$\begin{aligned} \Rightarrow \int_1^2 \frac{dx}{5+3x} &= \frac{3h}{8} [y_0 + y_6 + 3(y_1 + y_2 + y_4 + y_5) + 2y_3] \\ &= \frac{3 \times \frac{1}{6}}{8} [0.125 + 0.0909 + 3[0.1176 + 0.112 \\ &\quad + 0.1002 + 0.095] + 2 \times 0.1053] \\ &= \frac{1}{16} \times 1.6992 \end{aligned}$$

$$\therefore \int_1^2 \frac{dx}{5+3x} = 0.1062$$

Now,

By exact integration,

$$\begin{aligned} \int_1^2 \frac{dx}{5+3x} &= \ln(5+3x) \Big|_1^2 \\ &= -\ln 8 + \ln 11 \\ &= 0.10613. \end{aligned}$$

when,  $n=6$

$$\text{Absolute errors} = |0.10615 - 0.1062| \\ = 0.00005.$$

(B) Boole's formula is:-

$$\int_{x_0}^{x_4} y dx = \frac{2h}{45} (7y_0 + 32y_1 + 12y_2 + 32y_3 + 7y_4)$$

where  $h = x_1 - x_0$ .

$x$	0.0	0.1	0.2	0.3	0.4
$y$	1.000	0.9975	0.99	0.9776	0.8604

Now, we have:

$$x_0 = 0.000, x_1 = 0.1.$$

$$x_4 = 0.4$$

$$\therefore h = (x_4 - x_0)/n$$

$$= (0.4 - 0.0)/4$$

$$h = \underline{\underline{0.1}}.$$

from ①, we have:-

$$\int_0^{0.4} y \cdot dx = \frac{2 \times 0.1}{45} (7x1.00 + 32 \times 0.9975 + 32 \times 0.9776 \times 0.8604)$$
$$= \int_0^{0.4} y \cdot dx = \frac{2 \times 0.1 \times 88.106}{45}$$
$$= 0.39158$$

$$\boxed{\int_0^{0.4} y \cdot dx = 0.39158}$$

Ans.

⑨  $y = y_m$ ,  $a=1$ ,  $b=2$ ,  $N=8$

$$\therefore h = \frac{b-a}{N} = \frac{2-1}{8} = \frac{1}{8}$$

$$\Rightarrow x_m = a + n \cdot h, y_m = y_{xm}$$

Now,

x	1	9/8	10/8	11/8	12/8	13/8	14/8	15/8	2
y	1	<del>8/9</del>	8/10	8/11	8/12	8/13	8/14	8/15	y <sub>2</sub>

Using Boole's formula,

$$\int_a^b y \cdot dx = \frac{2h}{45} (7y_0 + 32y_1 + 12y_2 + 32y_3 + 14y_4 + 32y_5 + 12y_6 + 32y_7 + 14y_8 + \dots)$$

$$\therefore \int_1^2 y \cdot dx = \frac{2 \times y_8}{45} \left( 7x1 + 32 \times \frac{8}{9} + 12 \times \frac{8}{10} + 14 \times \frac{8}{12} + 32 \times \frac{8}{13} + 12 \times \frac{8}{14} + 14 \times \frac{8}{15} + \dots \right)$$