### The Fundamental Groupoid

Riley Shahar

Objects

x y


Z



- Objects
- Morphisms




- Objects
- Morphisms
- Composition



- Objects
- Morphisms
- Composition
- Identities


### **Category Axioms**

### Category Axioms



Associative

### Category Axioms



► Set: Sets and Functions

Set: Sets and Functions

► Grp: Groups and Homomorphisms

Set: Sets and Functions

► Grp: Groups and Homomorphisms

► Top: Spaces and Continuous Functions

- Objects
- Morphisms
- Composition
- Identities

- ▶ Objects: There is a single object, \*.
- ► Morphisms
- Composition
- Identities

- ▶ Objects: There is a single object, \*.
- ▶ Morphisms: For each  $x \in G$ , there is a morphism x.
- Composition
- Identities

- ▶ Objects: There is a single object, \*.
- ▶ Morphisms: For each  $x \in G$ , there is a morphism x.
- ► Composition: Composition is the group multiplication.
- Identities

- Objects: There is a single object, \*.
- ▶ Morphisms: For each  $x \in G$ , there is a morphism x.
- ► Composition: Composition is the group multiplication.
- ▶ Identities: The identity  $1_*$  is the morphism e.

Take a group G. Make a category  $\mathbb{G}$ :

- Objects: There is a single object, \*.
- ▶ Morphisms: For each  $x \in G$ , there is a morphism x.
- Composition: Composition is the group multiplication.
- ▶ Identities: The identity  $1_*$  is the morphism e.

Moral: not all categories have structured sets for objects and structure-preserving functions for morphisms.

Goal: generalize bijections, group isomorphisms, homeomorphisms etc.

Goal: generalize bijections, group isomorphisms, homeomorphisms etc.

#### Definition

A morphism  $f: x \to y$  is an *isomorphism* when there exists an *inverse morphism*  $g: y \to x$  such that

$$\mathit{fg} = 1_{\mathsf{y}} \quad \mathsf{and} \quad \mathit{gf} = 1_{\mathsf{x}}.$$

Goal: generalize bijections, group isomorphisms, homeomorphisms etc.

#### Definition

A morphism  $f: x \to y$  is an isomorphism when there exists an inverse morphism  $g: y \to x$  such that

$$\mathit{fg} = 1_{\mathsf{y}} \quad \mathsf{and} \quad \mathit{gf} = 1_{\mathsf{x}}.$$

What are isomorphisms in  $\mathbb{G}$ ?



Goal: generalize bijections, group isomorphisms, homeomorphisms etc.

#### Definition

A morphism  $f: x \to y$  is an *isomorphism* when there exists an *inverse morphism*  $g: y \to x$  such that

$$fg = 1_y$$
 and  $gf = 1_x$ .

What are isomorphisms in  $\mathbb{G}$ ? Everything!



What are isomorphisms in  $\mathbb{G}?$  Everything!

What are isomorphisms in  $\mathbb{G}\mbox{.}$  Everything! Let's generalize!

What are isomorphisms in  $\mathbb{G}$ ? Everything! Let's generalize!


#### Definition


A *groupoid* is a category in which every morphism is an isomorphism.


What are isomorphisms in  $\mathbb{G}$ ? Everything! Let's generalize!


#### Definition

A *groupoid* is a category in which every morphism is an isomorphism.









If you relax the requirement that paths in the fundamental group are loops, you get the fundamental groupoid  $\Pi_1$ .

▶ Is the fundamental groupoid a homotopy invariant? By the most obvious notion of groupoid isomorphism, **no**.

- ▶ Is the fundamental groupoid a homotopy invariant? By the most obvious notion of groupoid isomorphism, **no**.
- The problem is that there are many more points in  $D^n$  than  $\{*\}$ , hence many more objects in  $\Pi_1(D^n)$  than  $\Pi_1(\{*\})$ .

- ▶ Is the fundamental groupoid a homotopy invariant? By the most obvious notion of groupoid isomorphism, **no**.
- The problem is that there are many more points in  $D^n$  than  $\{*\}$ , hence many more objects in  $\Pi_1(D^n)$  than  $\Pi_1(\{*\})$ .
- ➤ The naive notion of isomorphism is too strong: we need a kind of weak equivalence, much like homotopy for topological spaces, for categories.

- ▶ Is the fundamental groupoid a homotopy invariant? By the most obvious notion of groupoid isomorphism, **no**.
- The problem is that there are many more points in  $D^n$  than  $\{*\}$ , hence many more objects in  $\Pi_1(D^n)$  than  $\Pi_1(\{*\})$ .
- ► The naive notion of isomorphism is too strong: we need a kind of weak equivalence, much like homotopy for topological spaces, for categories.
- ► This line of thought naturally leads to "higher categories", with morphisms between morphisms, etc.