

# Exercises Week 8

## DSP

1. Compute the circular convolution of the two signals:

$$x_1[n] = [1, 3, 1, 3]$$

$$x_2[n] = [2, 2, 5, 5]$$

2. Compute the circular convolution in  $N = 7$  points of the same two signals (i.e. append zeros to make length 7, then do circular convolution)
3. Do the same for a periodic signal  $x[n]$  with period  $N = 6$  and the DFT coefficients:

$$X_k = [21.0000 + 0.0000i, -3.0000 + 5.1962i, -3.0000 + 1.7321i, -3.0000 + 0.0000i, -3.0000 - 1.7321i, -3.0000 - 5.1962i]$$

Write  $x[n]$  as a sum of sinusoids.

4. Consider a periodic signal  $x[n]$  with period  $N = 5$  and the DFT coefficients:

$$X_k = [15.0000 + 0.0000i, -2.5000 + 3.4410i, -2.5000 + 0.8123i, -2.5000 - 0.8123i, -2.5000 - 3.4410i]$$

Write  $x[n]$  as a sum of sinusoids.

5. Find the DFT coefficients of the periodic signal with period  $\{1, 1, 0, 0\}$ , and write the signal as a sum of sinusoidal components.
6. Find the DTFT of the signal  $\{1, 1, 0, 0\}$ , considering it is surrounded with infinitely-long zeros on both sides