

Tutorial 11

SC-220 Groups and Linear algebra Autumn 2019
(Inner Products spaces)

- (1) Let V be an inner product space, then show that $\|x\| = \sqrt{\langle x | x \rangle}$ defines a norm on V .

- (2) Parallelogram law

Let V be a real or complex vector space with an inner product. Show that the norm defined by the inner product satisfies the parallelogram law

$$\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2$$

- (3) In $M_2(\mathbb{R})$ we can define an inner product $\langle A | B \rangle = \text{Tr}(AB^T)$. Verify that the set

$$\mathcal{B} = \left\{ \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \frac{1}{2} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \right\}$$

is an orthonormal basis. Compute the Fourier expansion of $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ with respect to \mathcal{B}

- (4) Consider the space of real valued integrable functions on the interval $(-\pi, \pi)$ with inner product

$$\langle f | g \rangle = \int_{-\pi}^{\pi} f(t)g(t)dt$$

Verify that the set of trigonometric

$$\mathcal{B} = \left\{ \frac{1}{\sqrt{2\pi}}, \frac{\cos t}{\sqrt{\pi}}, \frac{\cos 2t}{\sqrt{\pi}}, \dots, \frac{\sin t}{\sqrt{\pi}}, \frac{\sin 2t}{\sqrt{\pi}}, \dots \right\}$$

is an orthonormal basis of this space. Find the expansion of the square wave function

$$f(t) = \begin{cases} -1 & \text{when } -\pi < t < 0 \\ 1 & \text{when } 0 < t < \pi \end{cases}$$

- (5) Apply the Gram-Schmidt procedure for \mathbb{C}^3 to the vectors $\left\{ \begin{pmatrix} i \\ i \\ i \end{pmatrix}, \begin{pmatrix} 0 \\ i \\ i \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ i \end{pmatrix} \right\}$

- (6) Consider the vector space P_3 of polynomials with rational coefficients that are of degree less than or equal to 3 on the domain $[-1, 1]$. Check that $\langle f | g \rangle = \int_{-1}^1 f(x)g(x)dx$ is an inner product on P_3 . The monomials $\{1, x, x^2, x^3\}$ form a basis of P_3 . Use the Gram-Schmidt orthogonalization process to produce an ONB of P_3 . Express the polynomial $x^3 + 3x^2 + 2x + 3$ as a linear combination of the ONB vectors.