

Probability Assignment-III

Nikhil Nair

I. PROBLEM

Find the mean number of heads in three tosses of a fair coin.

II. SOLUTION

Consider each trial results in success (i.e Heads) or failure (i.e Tails).

Let p and q = (1 - p) be the probability of success and failure respectively.

$$p = \frac{1}{2}; q = 1 - p = \frac{1}{2} \tag{1}$$

In n Bernoulli trials with x success and (n - x) failures, the probablity of x success in n- Bernoulli trials can be given as

$$^{n}C_{x}p^{x}q^{n-x} \tag{2}$$

Now the distribution of number of successes using (1) & (2) can be given as,

X	0	1	2	3
P(X)	${}^{3}C_{0}(\frac{1}{2})^{3}$	${}^3C_1(\frac{1}{2})^3$	${}^3C_2(\frac{1}{2})^3$	${}^3C_3(\frac{1}{2})^3$

Mean of X =
$$\mu = \sum_{i=1}^{n=3} x_i P(x_i)$$

$$\mu = 0 \times^{3} C_{0}(\frac{1}{2})^{3} + 1 \times^{3} C_{1}(\frac{1}{2})^{3} + 2 \times^{3} C_{2}(\frac{1}{2})^{3} + 3 \times^{3} C_{3}(\frac{1}{2})^{3}$$

$$\therefore$$
 Mean of $X = \mu = 1.5$