

Effects of drugs on Blood **Pressure**

Dr Apurv Patel

2nd year resident

Department of pharmacology

➤ **Arterial blood pressure** is the product of the **cardiac output** (CO) and the **peripheral vascular resistance (PVR)**.

- $BP = CO \times PVR$
- $CO = HR \times SV$

➤ This means:

- If either CO or PVR increases, BP rises.
- Conversely, if they decrease, BP falls.

➤ When blood pressure (BP) **drops**, it triggers a response in the body to bring it back to normal. Pressure-sensitive sensors called **baroreceptors** (found in the aortic arch and carotid sinuses) detect this change and send signals to the brain.

➤ The brain responds by **activating the sympathetic nervous system** and **reducing signals from the parasympathetic system** (vagus nerve). This leads to:

✓ **Vasoconstriction (narrowing of blood vessels).**

✓ **Increased heart rate and cardiac output.**

➤ Together, these changes raise the blood pressure to compensate for the drop.

➤ On the flip side, when blood pressure gets too high, the baroreceptors send more signals to the brain. This reduces sympathetic activity and increases parasympathetic signals, causing:

✓ Vasodilation (widening of blood vessels).

✓ Slower heart rate (bradycardia).

- These adjustments lower the blood pressure back to normal.
- This process helps maintain stable blood pressure in the body.

Effect of Ach

❖ Heart rate: ↓

- Ach hyperpolarizes the SA node cells (M_2).
- It increases the refractory period of the AV node and Purkinje fibers.

❖ Blood pressure: ↓

- Ach dilates blood vessels by acting on M_3 receptors present on vascular endothelial cells, leading to the release of **nitric oxide**.

Effect of Atropine

❖ Heart rate:



- Atropine blocks muscarinic receptors (M_2) on the SA node, reducing vagal tone and increasing heart rate.
- A brief initial bradycardia may occur after i.m./s.c. injection due to increased ACh release from blocked muscarinic autoreceptors (M_1).

❖ Blood pressure:

- Atropine has minimal direct effects on BP as cholinergic impulses do not regulate vascular tone.
- **Factors influencing BP:**
 - ✓ Tachycardia increases cardiac output, potentially raising BP.
 - ✓ High doses may cause vasodilation, possibly lowering BP.

Effect of Adrenaline($\alpha_1\alpha_2\beta_1\beta_2$)

❖ Heart rate:

- Adrenaline increases heart rate by enhancing pacemaker activity in the SA node.
- When BP rises significantly, **reflex bradycardia** occurs:
- The rise in BP stimulates baroreceptors, which activate the vagus nerve to slow the heart.

❖ Blood pressure:

- Causes a temporary rise in both systolic and diastolic BP (α -effects predominate at high doses, causing widespread vasoconstriction).
- BP normalizes quickly due to rapid uptake of adrenaline, followed by a secondary BP drop as β_2 effects (vasodilation) dominate at lower concentrations.

Effect of Noradrenaline($\alpha_1\alpha_2\beta_1$)

❖ Heart rate: ↑

- Direct action of noradrenaline on the heart is to increase the heart rate by stimulating β_1 receptors.
- However, the indirect action of noradrenaline on the heart is to decrease the heart rate due to **reflex bradycardia** caused by a rise in blood pressure.

❖ Blood pressure: ↑

- Causes **vasoconstriction** via α -receptors, with no β_2 -mediated vasodilation.
- Increases **systolic, diastolic, and mean BP.**
- Peripheral resistance consistently increases.

Effect of Isoprenaline($\beta_1\beta_2$)

❖ Heart rate:



- Isoprenaline strongly stimulates beta-1 adrenergic receptors in the heart.
- The result is enhanced calcium ion availability, which increases the rate and force of cardiac contractions.

❖ Blood pressure:



- **Systolic Blood Pressure:** Due to the increased cardiac output from beta-1 stimulation, **systolic pressure may rise slightly** or remain stable.
- **Diastolic Blood Pressure:** Beta-2 receptor activation causes vasodilation in peripheral blood vessels, leading to a **reduction in diastolic blood pressure**.

Effect of Phentolamine($\alpha_1\alpha_2$ #)

❖ Heart rate: 

- The decrease in blood pressure triggers a reflex tachycardia via the baroreceptor reflex.
- Additionally, blocking α_2 receptors on presynaptic nerve terminals leads to increased release of norepinephrine, further stimulating β_1 adrenergic receptors in the heart.


❖ Blood pressure: 

- Phentolamine blocks α_1 receptors on vascular smooth muscle, causing vasodilation.
- This reduces peripheral vascular resistance, leading to a decrease in blood pressure

Department of Pharmacology, PIMSR, Vadodara

Spot exercise

Following graph shows the effects of drugs A, B and C on dog mean blood pressure. The responses of these drugs are recorded after administration of prazosin.

DOG BLOOD PRESSURE

A :
B :
C : Isoprenaline (2.0 μ g/kg)

A: Adrenaline
B: Nor adrenaline

Identify the probable nature of drugs A and B with reasons.

Vasomotor reversal of Dale

- Normally, **adrenaline** primarily acts on two types of adrenergic receptors in blood vessels:
 - ✓ **Alpha-adrenergic receptors**, which cause **vasoconstriction**.
 - ✓ **Beta-adrenergic receptors**, which cause **vasodilation**.
- In most blood vessels, alpha receptors dominate, so adrenaline typically causes vasoconstriction.

❖ Mechanism of Vasomotor Reversal

- When an **alpha-adrenergic receptor blocker** (such as **phentolamine** or **ergot alkaloids**) is administered, the alpha receptors are inhibited, preventing adrenaline from causing vasoconstriction. As a result:
- The **beta-adrenergic receptors**, which mediate vasodilation, become unopposed.
- Subsequent administration of adrenaline leads to **vasodilation** instead of vasoconstriction.
- This "reversal" of vascular response from constriction to dilation is the **vasomotor reversal of Dale**.

Effect of Ephedrine

❖ Heart rate: ↑

- Directly stimulates beta-1 adrenergic receptors in the heart, increasing the rate and force of cardiac contractions.
- Indirectly causes the release of norepinephrine, which further stimulates beta-1 receptors.
- **Outcome: Increases heart rate.**


❖ Blood pressure: ↑

- **Systolic Blood Pressure:** Increases due to enhanced cardiac output (beta-1 stimulation) and vasoconstriction (alpha-adrenergic receptor stimulation).
- **Diastolic Blood Pressure:** May increase or remain stable due to vasoconstriction from alpha-adrenergic receptor activation.

Department of Pharmacology, PIMSR, Vadodara

Spot exercise

Following graph shows the effect of repeated administration of same doses of drug A on mean blood pressure of dog.



Identify the probable nature of drug. Explain the effect.

Tachyphylaxis

Effect of Propranolol($\beta_1\beta_2$)

❖ Heart rate:



- Propranolol blocks beta-1 receptors in the heart, which decreases the effects of sympathetic nervous system stimulation.
- Propranolol causes a **reduction in heart rate**, especially under conditions of increased sympathetic activity (e.g., stress or exercise).

❖ Blood pressure:




- Propranolol has no direct effect on blood vessels.
- Propranolol decreases blood pressure primarily by reducing cardiac output (due to lower heart rate and contractility).
- It also inhibits the renin release from the kidneys (via beta-1 receptor blockade).

Department of Pharmacology, PIMSR, Vadodara

Spot exercise

The Following graph shows effects of drugs A, B and C on dog mean blood pressure. The responses of drugs A, B and C are obtained after administration of propranolol.



C : Isoprenaline

Identify the probable nature of drug C with reasons

Effect of Histamine

❖ Heart rate:



- Histamine acts on **H2 receptors** in the heart, which increase cyclic AMP (cAMP). This enhances calcium influx into cardiac cells, increasing the rate and force of contraction.
- Additionally, reflex tachycardia may occur as a compensatory response to histamine-induced hypotension.

❖ Blood pressure:




- **H1 receptors** in endothelial cells promote the release of nitric oxide, leading to vasodilation.
- **H2 receptors** in vascular smooth muscle cells directly induce relaxation.
- The combined effect is **systemic vasodilation**, particularly in the skin, which decreases peripheral resistance.

Department of Pharmacology, PIMSR, Vadodara

Spot exercise

Following graph shows the effects of drugs A and B on dog mean blood pressure. The responses of drugs A and B are obtained again after administration of chlorpheniramine.



B : Histamine

A : Acetylcholine (3 μ g/kg)

Identify the probable nature of drug B with reasons.
















Effect of Mepyramine







❖ Heart rate:

- Mepyramine does not directly act on the heart but may influence heart rate indirectly.
- At therapeutic doses, it typically has minimal direct impact on heart rate.

❖ Blood pressure:

- By antagonizing H₁ receptors, mepyramine blocks histamine-induced vasodilation in peripheral blood vessels, preventing the associated drop in blood pressure.
- However, in individuals without elevated histamine levels, the blood pressure effect is usually mild or negligible.


No.	Drug	Blood pressure	Heart rate
1.	<u>Acetylcholine</u>		
2.	<u>Atropine</u>		
3.	<u>Adrenaline</u>	 	
4.	<u>Noradrenaline</u>		
5.	<u>Isoprenaline</u>		
6.	<u>Histamine</u>		
7.	<u>Propranolol</u>		

No.	Drug	Blood pressure	Heart rate
8.	<u>Phentolamine</u>		
9.	<u>Ephedrine</u>		
10.	<u>Mepyramine</u>		

Department of Pharmacology, PIMSR, Vadodara

Spot exercise

The given graph shows plasma concentration of a drug given by two different routes of administration verses time.




X: Intravenous
Y: Oral

Identify the route of administration of graph X and Y. Give reasons.

Department of Pharmacology, PIMSR, Vadodara**Spot exercise**

Following graph shows Dose response curves of drug X and drug X in presence of drug A.



Competitive
antagonist

X : - Dose Response Curve of Drug X.

A+X: - Dose Response Curve of Drug X in presence of Drug A


Answer the following questions-

1. What is the type of drug antagonism observed in the graph?
2. Mention the reasons to support your answer.

Department of Pharmacology, PIMSR, Vadodara

Spot exercise

Following graph shows Dose response curves of drug X and drug X in presence of drug A.



Non competitive
antagonism

X :- Dose Response Curve of Drug X.

A+X: - Dose Response Curve of Drug X in presence of Drug A

Answer the following questions-

1. What is the type of drug antagonism observed in the graph?
2. Mention the reasons to support your answer.