1/1 point

1/1 point

1/1 point

1/1 point

1/1 point

1. In this quiz, you will practice changing from the standard basis to a basis consisting of orthogonal vectors.

Given vectors  $\mathbf{v}=\begin{bmatrix}5\\-1\end{bmatrix}$ ,  $\mathbf{b_1}=\begin{bmatrix}1\\1\end{bmatrix}$  and  $\mathbf{b_2}=\begin{bmatrix}1\\-1\end{bmatrix}$  all written in the standard basis, what is  $\mathbf{v}$  in the basis defined by  $\mathbf{b_1}$  and  $\mathbf{b_2}$ ? You are given that  $\mathbf{b_1}$  and  $\mathbf{b_2}$  are orthogonal to each other.

- $\bigcirc$   $\mathbf{v_b} = \begin{bmatrix} -3 \\ 2 \end{bmatrix}$
- $\bigcirc$   $\mathbf{v_b} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$
- $\bigcirc$   $\mathbf{v_b} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$
- $\bullet$   $\mathbf{v_b} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$

 $\bigodot$  correct  $\label{eq:correct} \mbox{The vector } v \mbox{ is projected onto the two vectors } b_1 \mbox{ and } b_2.$ 

 $\textbf{2.} \quad \text{Given vectors } \textbf{v} = \begin{bmatrix} 10 \\ -5 \end{bmatrix}, \textbf{b}_1 = \begin{bmatrix} 3 \\ 4 \end{bmatrix} \text{ and } \textbf{b}_2 = \begin{bmatrix} 4 \\ -3 \end{bmatrix} \text{ all written in the standard basis, what is } \textbf{v} \text{ in the basis defined by } \textbf{b}_1 \text{ and } \textbf{b}_2 \text{? You are given that } \textbf{b}_1 \text{ and } \textbf{b}_2 \text{ are orthogonal to each other.}$ 

 $\bigcirc$   $\mathbf{v_b} = \begin{bmatrix} 2 \\ 11 \end{bmatrix}$ 

- $\bigcirc \quad \mathbf{v_b} = \begin{bmatrix} -2/5 \\ 11/5 \end{bmatrix}$
- $\odot$   $\mathbf{v_b} = \begin{bmatrix} 2/5 \\ 11/5 \end{bmatrix}$

 $\odot$   $\,$  correct  $\,$  The vector v is projected onto the two vectors  $b_1$  and  $b_2$ 

 $\textbf{3.} \quad \text{Given vectors } \mathbf{v} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \mathbf{b_1} = \begin{bmatrix} -3 \\ 1 \end{bmatrix} \text{ and } \mathbf{b_2} = \begin{bmatrix} 1 \\ 3 \end{bmatrix} \text{ all written in the standard basis, what is } \mathbf{v} \text{ in the basis defined by } \mathbf{b_1} \text{ and } \mathbf{b_2} ? \text{ You are given that } \mathbf{b_1} \text{ and } \mathbf{b_2} \text{ are orthogonal to each other.}$ 

- $\mathbf{v_b} = \begin{bmatrix} -2/5 \\ 5/4 \end{bmatrix}$
- $\bigcirc$   $\mathbf{v_b} = \begin{bmatrix} 5/4 \\ -5/2 \end{bmatrix}$
- $\bigcirc \quad \mathbf{v_b} = \begin{bmatrix} 2/5 \\ -4/5 \end{bmatrix}$

 $\odot$   $\,$  correct  $\,$  The vector v is projected onto the two vectors  $b_1$  and  $b_2$ 

4. Given vectors  $\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{b_1} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{b_2} = \begin{bmatrix} 1 \\ -2 \\ -1 \end{bmatrix}$  and  $\mathbf{b_3} = \begin{bmatrix} -1 \\ 2 \\ -5 \end{bmatrix}$  all written in the standard basis, what is  $\mathbf{v}$  in the basis defined by  $\mathbf{b_1}$ ,  $\mathbf{b_2}$  and  $\mathbf{b_3}$ ? You are given that  $\mathbf{b_1}$ ,  $\mathbf{b_2}$  and  $\mathbf{b_3}$  are all pairwise or because the part of the pa

- $\bigcirc \qquad \mathbf{v_b} = \begin{bmatrix} -3/5 \\ -1/3 \\ 2/15 \end{bmatrix}$

 $\odot$  correct  $\label{eq:correct} \text{The vector } v \text{ is projected onto the vectors } b_1, b_2 \text{ and } b_3$ 

 $v_b = \begin{bmatrix} 1\\1\\0 \end{bmatrix}$   $\otimes$  correct The vector  ${\bf v}$  is projected onto the vectors  ${\bf b_1}, {\bf b_2}, {\bf b_3}$  and  ${\bf b_4}$ .