



Take a closer look at the risks of calcification

Come and visit the Genzyme booth or try this for yourself at home

– we hope to give you some idea of what a calcified heart may look like.


How to view a virtual version of a living heart

Ask a Renvela representative to show you how to use this card.

OR

Follow the instructions below to view at home.

1. Turn on your computer's sound and webcam.
2. Go to www.renvela.eu and click on Virtual Heart.
3. Please click to allow the pop-up window when prompted, to activate your webcam.
4. Hold this card straight, about a meter away from the webcam, with the hourglass symbol facing it.



The imagery showing calcification in the heart has been endorsed by Dr John R Cockcroft, Professor of Cardiology at the Heart Research Institute in Cardiff, Wales.

Vascular calcification is prevalent and progressive in CKD patients,¹⁻³ and is a strong predictor of mortality.⁴⁻⁵

- The use of calcium-based binders has been associated with the progression of vascular calcification in CKD patients.^{2,3,6}
- Sevelamer is an effective monotherapy that has been shown to attenuate vascular calcification^{1,7,8} and improve outcomes in CKD patients on dialysis.^{4,9,10}
- Renvela is the only phosphate binder available in tablets and an innovative powder formulation.

Renvela
sevelamer carbonate

Prescribing Information

Product Summary Renvela® 800 mg film-coated tablets. Renvela® 2.4 g powder for oral suspension. **Active substance:** Sevelamer. Subject to medical prescription. **Composition:** Each tablet contains 800 mg sevelamer carbonate. Each tablet also contains microcrystalline cellulose, sodium chloride and zinc stearate. The tablet coating contains hypromellose and diacetylated monoglycerides. The printing ink contains iron oxide black (E172), propylene glycol, isopropyl alcohol and hypromellose. Each sachet contains 2.4 g sevelamer carbonate powder. The powder contains propylene glycol alginate, citrus cream flavour, sodium chloride, sucralose, ferric oxide (E172). **Therapeutic indications:** Renvela is indicated for the control of hyperphosphataemia in adult patients receiving haemodialysis or peritoneal dialysis. Renvela is also indicated for the control of hyperphosphataemia in adult patients with chronic kidney disease not on dialysis with serum phosphorus ≥ 1.78 mmol/L. Renvela should be used within the context of a multiple therapeutic approach, which could include calcium supplement, 1,25-dihydroxy Vitamin D₃ or one of its analogues to control the development of renal bone disease. **Contra-indications:** Hypersensitivity to the active substance or to any of the excipients. Hypophosphataemia. Bowel obstruction. **Possible side effects:** The safety of sevelamer (as either carbonate or hydrochloride salts) has been investigated in numerous clinical trials involving a total of 969 haemodialysis patients with treatment duration of

4 to 50 weeks (724 patients treated with sevelamer hydrochloride and 245 with sevelamer carbonate), 97 peritoneal dialysis patients with treatment duration of 12 weeks (all treated with sevelamer hydrochloride) and 128 patients with CKD not on dialysis with treatment duration of 8 to 12 weeks (79 patients treated with sevelamer hydrochloride and 49 with sevelamer carbonate). The most frequently occurring ($\geq 5\%$ of patients) undesirable effects possibly or probably related to sevelamer were all in the gastrointestinal disorders system organ class. Most of these adverse reactions were mild to moderate in intensity. Data possibly or probably related to sevelamer from these studies are listed by frequency here next. Gastrointestinal disorders: nausea, vomiting, upper abdominal pain, constipation (very common) and diarrhoea, dyspepsia, flatulence, abdominal pain (common). The reporting rate is classified as very common ($\geq 1/10$), common ($\geq 1/100$, $< 1/10$), uncommon ($\geq 1/1,000$, $< 1/100$), rare ($\geq 1/10,000$, $< 1/1,000$), very rare ($< 1/10,000$), including isolated. **Post-marketing experience:** In very rare cases (estimated post-marketing frequency of $< 1/10,000$), intestinal obstruction and ileus/subileus have been observed in patients during treatment with sevelamer hydrochloride, which contains the same active moiety as sevelamer carbonate. **Product licence holder:** Genzyme Europe BV, Gooimeer 10, 1411 DD Naarden, The Netherlands. **Date of revision of text:** June 2009. **Attention:** Please see full prescribing information.

References

1. Russo D et al. *Kidney Int* 2007;72(10):1255-1261.
2. Goodman WG et al. *N Engl J Med* 2000;342(20):1478-1483.
3. Guérin AP et al. *Nephrol Dial Transplant* 2000;15:1014-1021.
4. Block GA et al, on behalf of RIND outcomes. *Kidney Int* 2007;71:438-441.
5. Blacher J et al. *Hypertension* 2001;38:938-942.
6. London GM et al. *Nephrol Dial Transplant* 2003;18:173-1740.
7. Chertow GM et al. *Kidney Int* 2002;62:245-252.
8. Block GA et al. *Kidney Int* 2005;68(4):1815-1824.
9. Suki WD et al. *Kidney Int* 2007;72(9):1130-1137.
10. Borzecki AM et al. *J Clin Pharma Thera* 2007;32:617-624.

Date of Preparation: June 2010
RP230 06/10

genzyme
Renal