CS & IT ENGINEERING

Database

Introduction of RDBMS

DPP 01 Discussion Notes



Vijay Agarwal sir






TOPICS TO BE COVERED

01 Question

02 Discussion



According to RDBMS rules, choose the correct statement from the following.





A relation in RDBMS can have multiple attributes.



A relation in RDBMS is a set of rows and columns.




A tuple in a relation can have multiple values for an attribute.




All of the above



Multivalued






Consider the student relation shown below with schema stud (Sname, Sage, Smail, Smarks), [NAT]

id (W)

Stud

Sname	Sage	Smail	Smarks
Rohit	28	R@pw.live	68
Kanika	25	K@pw.live	75
Pankaj	25	K@pw.live	75
Rohit	28	R@pw.live	88
Anjali	26	A@pw.live	75



For the above given instance how many 2-set of attributes can determine a row uniquely?

Consider a relation schema R(A, B, C, D, E, F, H) with the given

Functional dependency set:

 $\{A \rightarrow BC, C \rightarrow AD, DE \rightarrow F, C \rightarrow F\}$

The attribute closure that contains all the attributes of the relation R is?








AEH+ [AFH] - [AFHBCDF] - (ABCD FFH)

All of the above





(G)

Consider the below relation schema Stud (Rid, name, course, mail, phone) with FD set as:



Rid → {Rid} -> Toivial FD

Rid → {name, mail} → Non Trivial FD

Semi course -> {course, phone}

phone → {phone} → Tovial

mail → {Rid, course} -> Non Tokial FD (5)

Servir Non Trivial: 1

name → {phone, mail, course} The number of non-trivial FD's in the given FD set is/are?



Tovial FD X -y is tovial

IR X = y

 $AB \rightarrow B$ $AB \rightarrow AB$

Non Trivial FD

X-y is Non Torvial

ibb XMy = \$ A Must Satisfy FD Defination

> A -> B Rall -> Branch

Semi Non Toivial FD

X & y 4 × Ny + o

A -> AB

Jeither x \$ y

exny + 6

Q.5


Consider the following set of FD's:

$$\{V \rightarrow W, W \rightarrow XZ, X \rightarrow YZ\}$$
 for relation

R(V, W, X, Y, Z)

Then the attribute closure of YZ⁺ contains how many elements?





For the given FD set: $\{P \to QT, Q \to SU, V \to U\}$ of a relation R(P, Q, T, S, U, V). Find the set of attributes that is Super key but not a [MCQ]



A PTQ
$$[PTQ]^{\dagger} = [PTQSU]$$

B. $[PV]$ $[PV]^{\dagger} = [PVQTSU] \Rightarrow [PQTSUV]$

PQV $[QV]^{\dagger} = [QVSU]$

QV $[QV]^{\dagger} = [QVSU]$

Keys Supel Set of Subel Keys is Super key getting all attribute of Relational Super key

Schena R then X is super key Candidate key: Minimal of Super Key


AB is c.k

(Note) Every Condidate key is a Subselley Ret Every Subser key is Not Candidate key

C.K

P.K are
Super
Key

Super key -> Candidate by (4 C.K Minimal I Select as & Poimally key Remaining 3CK AK/ Secondeen key



In a schema with attribute X, Y, Z, W, V, the following set of functional dependencies are given:



$$\{Y \rightarrow X, Y \rightarrow Z, ZW \rightarrow V, X \rightarrow W, V \rightarrow X\}.$$

Which of the following FD is not implied by the above set?





$$YX \rightarrow ZW$$

$$\underline{\mathbf{YX}} \to \underline{\mathbf{ZW}} \qquad (\underline{\mathbf{YX}}^{\dagger} - (\underline{\mathbf{YX}} \times \underline{\mathbf{Z}} \, \underline{\mathbf{W}} \, \underline{\mathbf{V}})$$



$$XV \rightarrow YZ$$



$$ZW \rightarrow V$$



$$XV \rightarrow XW$$

$$\frac{ZW \to V}{XV \to XW} \left(\frac{ZW}{Y} \right)^{+} = \left(\frac{ZW \times X}{X} \right)^{+} = \left(\frac$$

F [A>B, B>c]

Check ADC ? (A) = [ABC]


X-y Logically implied from the closure of X

(X3t-[...y.] determines y.



Choose the correct statement from the following.







The cardinality is defined as the number of attributes in a relation.




Degree of the relation is the number of tuples in the relation.




Relation instance is the set of tuples of a relation at a particular instance of time.



All of the above





Depoee

Arity: Number of Attorbute

(4)

Relational schema: STUPENT (Sid Sname Brand Marly) (6)

Relational Instance: Set of Records (Snapshot



