

Thermalisation of interacting bosons on a lattice

N. MITCHELL¹, D. HUTCHINSON¹ and D. SCHUMAYER¹

¹ *University of Otago - Address*

PACS **nn.mm.xx** – First pacs description
PACS **nn.mm.xx** – Second pacs description
PACS **nn.mm.xx** – Third pacs description

Abstract –This paper is inspired by work done by Marcos Rigol et al. ??, who showed numerically that a one-dimensional gas of hard-core bosons on an optical lattice (an integrable system) relaxes to an equilibrium state that is characterised by the Generalised Gibbs Ensemble (GGE). The GGE is a thermodynamic ensemble that has been extended to take into account all of the conserved quantities of a system. We explore the question of whether similar behaviour can be seen in smaller systems, which we can solve more exactly, where the strength of the interactions is increased to approach the hard-core limit.




Table 1: Table caption.

first	table	row
second	table	row

Fig. 1: Figure caption.

Section title. – Insert here the text. See fig. 1, table 1 and eq. (1). See also [1,2].

$$0 \neq 1 \quad (1)$$

* * *

Insert here the text.

REFERENCES

- [1] AUTHOR F., AUTHOR S. and AUTHOR T., *Some Rev. A*, **69** (1969) 9691.
- [2] AUTHOR F. and AUTHOR S., *Some Book of Interest*, edited by A. EDITOR, Vol. **9** (Publishing house, City) 1939, p. 666.
- [3] EDITOR A. (Editor), *Some Book of Interest*, Vol. **9** (Publishing house, City) 1939, sect. A.