

Testiranje hipoteza o učestalostima

Nedelja 7 - Vežbe

Dr Nikola N. Grubor

The NEW ENGLAND JOURNAL *of* MEDICINE

ESTABLISHED IN 1812

MAY 1, 2008

VOL. 358 NO. 18

Treatment of Hypertension in Patients 80 Years of Age or Older

Nigel S. Beckett, M.B., Ch.B., Ruth Peters, Ph.D., Astrid E. Fletcher, Ph.D., Jan A. Staessen, M.D., Ph.D.,
Lisheng Liu, M.D., Dan Dumitrescu, M.D., Vassil Stoyanovsky, M.D., Riitta L. Antikainen, M.D., Ph.D.,
Yuri Nikitin, M.D., Craig Anderson, M.D., Ph.D., Alli Belhani, M.D., Françoise Forette, M.D.,
Chakravarthi Rajkumar, M.D., Ph.D., Lutgarde Thijs, M.Sc., Winston Banya, M.Sc.,
and Christopher J. Bulpitt, M.D., for the HYVET Study Group*

ABSTRACT

METHODS

We randomly assigned 3845 patients from Europe, China, Australasia, and Tunisia who were 80 years of age or older and had a sustained systolic blood pressure of 160 mm Hg or more to receive either the diuretic indapamide (sustained release, 1.5 mg) or matching placebo. The angiotensin-converting–enzyme inhibitor perindopril (2 or 4 mg), or matching placebo, was added if necessary to achieve the target blood pressure of 150/80 mm Hg. The primary end point was fatal or nonfatal stroke.

RESULTS

The active-treatment group (1933 patients) and the placebo group (1912 patients) were well matched (mean age, 83.6 years; mean blood pressure while sitting, 173.0/90.8 mm Hg); 11.8% had a history of cardiovascular disease. Median follow-up was 1.8 years. At 2 years, the mean blood pressure while sitting was 15.0/6.1 mm Hg lower in the active-treatment group than in the placebo group. In an intention-to-treat analysis, active treatment was associated with a 30% reduction in the rate of fatal or nonfatal stroke (95% confidence interval [CI], -1 to 51; $P=0.06$), a 39% reduction in the rate of death from stroke (95% CI, 1 to 62; $P=0.05$), a 21% reduction in the rate of death from any cause (95% CI, 4 to 35; $P=0.02$), a 23% reduction in the rate of death from cardiovascular causes (95% CI, -1 to 40; $P=0.06$), and a 64% reduction in the rate of heart failure (95% CI, 42 to 78; $P<0.001$). Fewer serious adverse events were reported in the active-treatment group (358, vs. 448 in the placebo group; $P=0.001$).

(A.E.R.) both in London, the University of Leuven, Leuven, Belgium (J.A.S., L.T.); the Beijing Hypertension League Institute, Beijing (L.L.); Spitalul Judetean Cluj, Clinica Medicală 2, Cluj, Romania (D.D.); the National Transport Multi-Profile Hospital, Sofia, Bulgaria (V.S.); the University of Oulu, Oulu, Finland (R.L.A.); the State Scientific Research Institute of Internal Medicine, Novosibirsk, Russia (Y.N.); the George Institute for International Health, Sydney (C.A.); L'Etablissement Public de Santé Charles Nicolle, Service de Cardiologie, Tunis, Tunisia (A.B.); Hôpital Broca, University Paris V, Paris (F.F.); and the Brighton and Sussex Medical School, Brighton, United Kingdom (C.R.). Address reprint requests to Dr. Beckett at Care of the Elderly, Division of Medicine, Imperial College London, Du Cane Rd., London W12 0NN, United Kingdom.

*The committee members and investigators for the Hypertension in the Very Elderly Trial (HYVET) are listed in the Appendix.

This article (10.1056/NEJMoa0801369) was published at www.nejm.org on March 31, 2008.

METHODS

We randomly assigned 3845 patients from Europe, China, Australasia, and Tunisia who were 80 years of age or older and had a sustained systolic blood pressure of 160 mm Hg or more to receive either the diuretic indapamide sustained release, 1.5 mg) or matching placebo. The angiotensin-converting–enzyme inhibitor perindopril (2 or 4 mg), or matching placebo, was added if necessary to achieve the target blood pressure of 150/80 mm Hg. The primary end point was fatal or nonfatal stroke.

Šta je nulta hipoteza?

RESULTS

The active-treatment group (1933 patients) and the placebo group (1912 patients) were well matched (mean age, 83.6 years; mean blood pressure while sitting, 173.0/90.8 mm Hg); 11.8% had a history of cardiovascular disease. Median follow-up was 1.8 years. At 2 years, the mean blood pressure while sitting was 15.0/6.1 mm Hg lower in the active-treatment group than in the placebo group. In an intention-to-treat analysis, active treatment was associated with a 30% reduction in the rate of fatal or nonfatal stroke (95% confidence interval [CI], –1 to 51; $P=0.06$), a 39% reduction in the rate of death from stroke (95% CI, 1 to 62; $P=0.05$), a 21% reduction in the rate of death from any cause (95% CI, 4 to 35; $P=0.02$), a 23% reduction in the rate of death from cardiovascular causes (95% CI, –1 to 40; $P=0.06$), and a 64% reduction in the rate of heart failure (95% CI, 42 to 78; $P<0.001$). Fewer serious adverse events were reported in the active-treatment group (358, vs. 448 in the placebo group; $P=0.001$).

METHODS

We randomly assigned 3845 patients from Europe, China, Australasia, and Tunisia who were 80 years of age or older and had a sustained systolic blood pressure of 160 mm Hg or more to receive either the diuretic indapamide (sustained release, 1.5 mg) or matching placebo. The angiotensin-converting–enzyme inhibitor perindopril (2 or 4 mg), or matching placebo, was added if necessary to achieve the target blood pressure of 150/80 mm Hg. The primary end point was fatal or nonfatal stroke.

RESULTS

The active-treatment group (1933 patients) and the placebo group (1912 patients) were well matched (mean age, 83.6 years; mean blood pressure while sitting, 173.0/90.8 mm Hg); 11.8% had a history of cardiovascular disease. Median follow-up was 1.8 years. At 2 years, the mean blood pressure while sitting was 15.0/6.1 mm Hg lower in the active-treatment group than in the placebo group. In an intention-to-treat analysis, active treatment was associated with a 30% reduction in the rate of fatal or nonfatal stroke (95% confidence interval [CI], –1 to 51; $P=0.06$), a 39% reduction in the rate of death from stroke (95% CI, 1 to 62; $P=0.05$), a 21% reduction in the rate of death from any cause (95% CI, 4 to 35; $P=0.02$), a 23% reduction in the rate of death from cardiovascular causes (95% CI, –1 to 40; $P=0.06$), and a 64% reduction in the rate of heart failure (95% CI, 42 to 78; $P<0.001$). Fewer serious adverse events were reported in the active-treatment group (358, vs. 448 in the placebo group; $P=0.001$).

**Da li su uzorci
nezavisni?**


METHODS

We randomly assigned 3845 patients from Europe, China, Australasia, and Tunisia who were 80 years of age or older and had a sustained systolic blood pressure of 160 mm Hg or more to receive either the diuretic indapamide (sustained release, 1.5 mg) or matching placebo. The angiotensin-converting–enzyme inhibitor perindopril (2 or 4 mg), or matching placebo, was added if necessary to achieve the target blood pressure of 150/80 mm Hg. The primary end point was fatal or nonfatal stroke.

RESULTS

The active-treatment group (1933 patients) and the placebo group (1912 patients) were well matched (mean age, 83.6 years; mean blood pressure while sitting, 173.0/90.8 mm Hg); 11.8% had a history of cardiovascular disease. Median follow-up was 1.8 years. At 2 years, the mean blood pressure while sitting was 15.0/6.1 mm Hg lower in the active-treatment group than in the placebo group. In an intention-to-treat analysis, active treatment was associated with a 30% reduction in the rate of fatal or nonfatal stroke (95% confidence interval [CI], –1 to 51; $P=0.06$), a 39% reduction in the rate of death from stroke (95% CI, 1 to 62; $P=0.05$), a 21% reduction in the rate of death from any cause (95% CI, 4 to 35; $P=0.02$), a 23% reduction in the rate of death from cardiovascular causes (95% CI, –1 to 40; $P=0.06$), and a 64% reduction in the rate of heart failure (95% CI, 42 to 78; $P<0.001$). Fewer serious adverse events were reported in the active-treatment group (358, vs. 448 in the placebo group; $P=0.001$).

Šta mi govori interval poverenja od 95%?



METHODS

We randomly assigned 3845 patients from Europe, China, Australasia, and Tunisia who were 80 years of age or older and had a sustained systolic blood pressure of 160 mm Hg or more to receive either the diuretic indapamide (sustained release, 1.5 mg) or matching placebo. The angiotensin-converting–enzyme inhibitor perindopril (2 or 4 mg), or matching placebo, was added if necessary to achieve the target blood pressure of 150/80 mm Hg. The primary end point was fatal or nonfatal stroke.

RESULTS

The active-treatment group (1933 patients) and the placebo group (1912 patients) were well matched (mean age, 83.6 years; mean blood pressure while sitting, 173.0/90.8 mm Hg); 11.8% had a history of cardiovascular disease. Median follow-up was 1.8 years. At 2 years, the mean blood pressure while sitting was 15.0/6.1 mm Hg lower in the active-treatment group than in the placebo group. In an intention-to-treat analysis, active treatment was associated with a 30% reduction in the rate of fatal or nonfatal stroke (95% confidence interval [CI], –1 to 51; $P=0.06$), a 39% reduction in the rate of death from stroke (95% CI, 1 to 62; $P=0.05$), a 21% reduction in the rate of death from any cause (95% CI, 4 to 35; $P=0.02$), a 23% reduction in the rate of death from cardiovascular causes (95% CI, –1 to 40; $P=0.06$), and a 64% reduction in the rate of heart failure (95% CI, 42 to 78; $P<0.001$). Fewer serious adverse events were reported in the active-treatment group (358, vs. 448 in the placebo group; $P=0.001$).

Šta mi govori interval poverenja od 95%?

A p-vrednost?

METHODS

We randomly assigned 3845 patients from Europe, China, Australasia, and Tunisia who were 80 years of age or older and had a sustained systolic blood pressure of 160 mm Hg or more to receive either the diuretic indapamide (sustained release, 1.5 mg) or matching placebo. The angiotensin-converting–enzyme inhibitor perindopril (2 or 4 mg), or matching placebo, was added if necessary to achieve the target blood pressure of 150/80 mm Hg. The primary end point was fatal or nonfatal stroke.

RESULTS

The active-treatment group (1933 patients) and the placebo group (1912 patients) were well matched (mean age, 83.6 years; mean blood pressure while sitting, 173.0/90.8 mm Hg); 11.8% had a history of cardiovascular disease. Median follow-up was 1.8 years. At 2 years, the mean blood pressure while sitting was 15.0/6.1 mm Hg lower in the active-treatment group than in the placebo group. In an intention-to-treat analysis, active treatment was associated with a 30% reduction in the rate of fatal or nonfatal stroke (95% confidence interval [CI], -1 to 2 ; P=0.06), a 39% reduction in the rate of death from stroke (95% CI, 1 to 62; P=0.05), a 21% reduction in the rate of death from any cause (95% CI, 4 to 35; P=0.02), a 23% reduction in the rate of death from cardiovascular causes (95% CI, -1 to 40; P=0.06), and a 64% reduction in the rate of heart failure (95% CI, 42 to 78; P<0.001). Fewer serious adverse events were reported in the active-treatment group (358, vs. 448 in the placebo group; P=0.001).

**Šta ako je 95% CI
[-1, 2]; p = 0.06?**

METHODS

We randomly assigned 3845 patients from Europe, China, Australasia, and Tunisia who were 80 years of age or older and had a sustained systolic blood pressure of 160 mm Hg or more to receive either the diuretic indapamide (sustained release, 1.5 mg) or matching placebo. The angiotensin-converting–enzyme inhibitor perindopril (2 or 4 mg), or matching placebo, was added if necessary to achieve the target blood pressure of 150/80 mm Hg. The primary end point was fatal or nonfatal stroke.

RESULTS

The active-treatment group (1933 patients) and the placebo group (1912 patients) were well matched (mean age, 83.6 years; mean blood pressure while sitting, 173.0/90.8 mm Hg); 11.8% had a history of cardiovascular disease. Median follow-up was 1.8 years. At 2 years, the mean blood pressure while sitting was 15.0/6.1 mm Hg lower in the active-treatment group than in the placebo group. In an intention-to-treat analysis, active treatment was associated with a 30% reduction in the rate of fatal or nonfatal stroke (95% confidence interval [CI], −1 to 2 ; P=0.06), a 39% reduction in the **rate of death from stroke (95% CI, 1 to 62; P=0.05)**, a 21% reduction in the rate of death from any cause (95% CI, 4 to 35; P=0.02), a 23% reduction in the rate of death from cardiovascular causes (95% CI, −1 to 40; P=0.06), and a 64% reduction in the rate of heart failure (95% CI, 42 to 78; P<0.001). Fewer serious adverse events were reported in the active-treatment group (358, vs. 448 in the placebo group; P=0.001).

**Šta ako je p-vrednost
0.05 umesto 0.06?**

METHODS

We randomly assigned 3845 patients from Europe, China, Australasia, and Tunisia who were 80 years of age or older and had a sustained systolic blood pressure of 160 mm Hg or more to receive either the diuretic indapamide (sustained release, 1.5 mg) or matching placebo. The angiotensin-converting–enzyme inhibitor perindopril (2 or 4 mg), or matching placebo, was added if necessary to achieve the target blood pressure of 150/80 mm Hg. The primary end point was fatal or nonfatal stroke.

RESULTS

The active-treatment group (1933 patients) and the placebo group (1912 patients) were well matched (mean age, 83.6 years; mean blood pressure while sitting, 173.0/90.8 mm Hg); 11.8% had a history of cardiovascular disease. Median follow-up was 1.8 years. At 2 years, the mean blood pressure while sitting was 15.0/6.1 mm Hg lower in the active-treatment group than in the placebo group. In an intention-to-treat analysis, active treatment was associated with a 30% reduction in the rate of fatal or nonfatal stroke (95% confidence interval [CI], –1 to 2 ; P=0.06), a 39% reduction in the rate of death from stroke (95% CI, 1 to 62; P=0.05), a 21% reduction in the rate of death from any cause (95% CI, 4 to 35; P=0.02), a 23% reduction in the rate of death from cardiovascular causes (95% CI, –1 to 40; P=0.06), and a 64% reduction in the **rate of heart failure (95% CI, 42 to 78; P<0.001)**. Fewer serious adverse events were reported in the active-treatment group (358, vs. 448 in the placebo group; P=0.001).

**Šta ako je p-vrednost
< 0.001 umesto 0.06?**

Greške u Zaključivanju


Tabela 1. Ishodi odlučivanja u testiranju hipoteza

		Istina (populacija)
Odluka istraživača na osnovu analize uzoračkih podataka	H_0 istinita	H_0 neistinita
Odbacivanje H_0	Greška prvog tipa (α)	Ispravan zaključak ($1-\beta$ = snaga testa)
Prihvatanje H_0	Ispravan zaključak ($1-\alpha$ = nivo poverenja)	Greška drugog tipa (β)

Koraci u statističkom zaključivanju

1. Hipoteza
2. Izbor značajnosti (0.05, 0.01)
3. Izbor test statistike (t -statistika, χ^2 -statistika, itd.)
4. Izračunavanje statistike testa
5. Zaključak (p-vrednost)

Studentov t-test




1. Slučajno biran uzorak
2. Numerički podaci
3. Normalna raspodela u populaciji

Tabela kontigencije očekivanih vrednosti.

	Bronhitis	Bez bronhitisa	Ukupno
Kašalj			70
Bez kašlja			1249
Ukupno	273	1046	1319

5.3%

94.7%



**Tabela kontigencije
očekivanih
vrednosti.**

	Bronhitis	Bez bronhitisa	Ukupno
Kašalj	14.5	55.5	70
Bez kašlja	258.5	990.5	1249
Ukupno	273	1046	1319

Očekivane
učestalosti

	Bronhitis	Bez bronhitisa	Ukupno
Kašalj	26 (14.5)	44 (55.5)	70
Bez kašlja	247 (258.5)	1002 (990.5)	1249
Ukupno	273	1046	1319

Opservirane
učestalosti

247
(258.5)

1002
(990.5)

Hi-kvadrat test

Kategorijalni
podaci




1. **Apsolutne učestalosti (ne procenti, proporcije, merenja)**
2. **Uzorak čine nezavisne jedinice**
3. **2 kategorije, nijedna očekivana f-ja ne sme < 5**
4. **3+ kategorije, ne sme biti više od 20% očekivanih f-ja < 5**
5. **$N > 20$**

Hi-kvadrat test

Tipovi

- Test slaganja (goodness-of-fit)
- Test za $r \times k$ (2×2) tabele

$$\chi^2 = \sum \frac{(O_i - E_i)^2}{E_i}$$



Hi-kvadrat test

Test slaganja (goodness-of-fit)

	A	B	AB	O
Učestalost	59	19	7	55

Hi-kvadrat test

Test slaganja (goodness-of-fit)

	A	B	AB	O
Učestalost	59	19	7	55

O – 44.5%




A – 38.9%

B – 12.1%

AB – 4.5%

Hi-kvadrat test


Test slaganja (goodness-of-fit)



Hi-kvadrat test

Test za 2x2 tabele

Preuzeti i analizirati bazu podataka **Osteoporoz.xlsx**



Hi-kvadrat test

Test za formirane 2x2 tabele

		Patološki nalaz na jetri		Ukupno
		+	-	
Konzumiranje alkohola	+	10	16	26
	-	13	79	92
Ukupno		23	95	118

Šta je nulta hipoteza?

Hi-kvadrat test


Test za formirane 2x2 tabele

		Patološki nalaz na jetri		Ukupno
		+	-	
Konzumiranje alkohola	+	10	16	26
	-	13	79	92
Ukupno		23	95	118

Ne postoji razlika učestalosti patološkog nalaza na jetri u odnosu na konzumiranje alkohola.

Hi-kvadrat test

Test za formirane 2x2 tabele



The screenshot shows the 'Enter Two-Way Table' dialog box. The 'Table' tab is selected. The 'Number of Rows:' field shows 2, and the 'Number of Columns:' field shows 2. Below, a 2x2 table is displayed with the following counts:

	Patol	Bez p
Konz	10	16
Bez k	13	79

At the bottom, there are 'Help', 'Reset', and 'OK' buttons. The 'OK' button is highlighted with a red border.

McNemarov test


Test za vezane kategorije

- McNemarov test je neparametarski metod za testiranje dva vezana (zavisna) uzoraka.
- Iste jedinice opservirane više puta
- Mečovane jedinice dva uzorka

		Drugi uzorak (ili druga observacija)		N
		+	-	
Prvi uzorak (ili prva observacija)	+	a	b	a+b
	-	c	d	c+d
		a+c	b+d	

McNemarov test

Preuzeti i analizirati bazu podataka **Aritmije.xlsx**



Fisherov test tačne verovatnoće

Fisher's exact test


	Mleko prvo	Čaj prvo	Ukupno
Pogodak	4	4	8
Promašaj	0	0	0
Ukupno	4	4	8



Fisherov test tačne verovatnoće



Fisher's exact test

Preuzeti i analizirati bazu podataka **Alkohol i hipertenzija.xlsx**



Fisherov test tačne verovatnoće

Fisher's exact test




Numerički ili kategorijalni podaci?

Tip testa	Parametarski	Neparametarski
Jedan uzorak	t-test za jedan uzorak	Hi-kvadrat test slaganja
Dva nezavisna	t-test	Hi-kvadrat / Fisherov test tačne verovatnoće
Dva zavisna	t-test za zavisne uzorke	McNemarov test

Relation between insufficient response to antihypertensive treatment and poor compliance with treatment: a prospective case-control study

Reto Nuesch, Kerstin Schroeder, Thomas Dieterle, Benedict Martina, Edouard Battegay



Status	Method of BP measurement	
	Clinic	Ambulatory
Responder	43	55
Non-responder	62	50
Total	105	105

Fisherov test tačne verovatnoće
 $p = 0.127$

Relation between insufficient response to antihypertensive treatment and poor compliance with treatment: a prospective case-control study

Reto Nuesch, Kerstin Schroeder, Thomas Dieterle, Benedict Martina, Edouard Battegay

Status	Method of BP measurement	
	Clinic	Ambulatory
Responder	43	55
Non-responder	62	50
Total	105	105

**Fisherov test tačne verovatnoće
 $p = 0.127$**

Zašto ovaj test nije dobro upotrebljen?

Kako to utiče na p-vrednost?

Šta su trebali da urade?