

Statistical analysis

Graphs and tables

Tools

Help

Original menu

Discrete variables

Continuous variables

Nonparametric tests

Survival analysis

Accuracy of diagnostic test

Matched-pair analysis

Metaanalysis and metaregression

Calculate sample size

Numerical summaries

Smirnov-Grubbs test for outliers

Kolmogorov-Smimov test for normal distribution

Confidence interval for a mean

Single-sample t-test

Two-variances F-test

Two-sample t-test

Paired t-test

Bartlett's test

One-way ANOVA

Repeated-measures ANOVA

Multi-way ANOVA

ANCOVA

Test for Pearson's correlation

Linear regression

Model:  $\Sigma$  <No active model>

```
16)
,c("Kreatinin", "Urea"),
), quantiles=c(0,.25,.5,.
(0,10), main="Sample")
#####
```

Test for Pearson's correlation

Click pressing Ctrl key to select multiple variables

Variables (pick two)

Kreatinin

rb

Urea

Alternative Hypothesis

☒ Two-sided


☐ Correlation < 0


☐ Correlation > 0


Condition to limit samples for analysis. Ex1. age>50

<all valid cases>

<

 Help

 Reset

 OK



**Da li postoji povezanost depresije i sistolne arterijske tenzije? Testirati za nivo značajnosti 0.05.**

rb	skor	ta
1	23	139
2	19	109
3	26	113
4	23	128
5	19	124
6	17	105
7	23	116
8	26	135
9	20	120
10	19	124