
RedSnapper
nyears = 70 proyears = 50 3 sampled iterations
M
Msd
M by Year
M/K
M−at−length
First historical year
Last historical year
Last projected year
Length
M−at−age
Age
Mexp
Linf
Linfsd
Linf by Year
t0
K
Ksd
K by Year
Steepness (h)
procsd
AC
Rec Devs by Year
Stock−Recruit
Depletion
L50
L95
Maturity−at−length (current year)
Maturity−at−age (current year)


Sampled length−at−age curves
First historical year
Last historical year
Last projected year
Length
0
20
40
60
80
5
10
15
20
Age


Fleet: RedSnapper Stock: RedSnapper
nyears = 70 proyears = 50 3 sampled iterations
Esd
qinc
qcv
Spat_targ
Fishing effort (historical)
Change in future catchability
Selectivity−at−length
First historical year
Last historical year
Last projected year
L5
LFS
Vmaxlen
Selectivity−at−age
First historical year
Last historical year
Last projected year

Observation biases and sample sizes for observation object RedSnapper

































Observation time series plots for observation object RedSnapper

























Implementation error time series plots for implementation object Perfect_Imp




















Spawning biomass
0
20000
50000
1950
1960
1970
1980
1990
2000
2010
2020
Historical year


Stock depletion (SSB)
0
0.5
1
1.5
1950
1960
1970
1980
1990
2000
2010
2020
Historical year


Fishing mortality rate (apical)
0
0.2
0.4
0.6
0.8
1950
1960
1970
1980
1990
2000
2010
2020
Historical year


Annual catches
0
5000
10000
1950
1960
1970
1980
1990
2000
2010
2020
Historical year


Recruitment
0
2000
4000
1950
1960
1970
1980
1990
2000
2010
2020
Historical year


Recruitment
0
2000
4000
0
10000
20000
30000
40000
Spawning biomass


F/F_MSY
B/B_MSY
91.7%
8.3%
0%
0%
1950
2019
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5


Age
Age vulnerability (0.25, 0.75)
Simulation 1
Simulation 2
Simulation 3
5
10
15
20
1950
1960
1970
1980
1990
2000
2010
2020
Historical year