

How numerical simulations work: Simulating particles in electric and magnetic fields

Nikolaj Roager Christensen

Student Colloquium in Physics and Astronomy, Aarhus University

March 2021

TITLEIMAGE

How numerical simulations work: Simulating particles in electric and magnetic fields

Introduction [5 MIN]

Theory and physical background [5 MIN]

Euler's Method and the 4th order Runge-Kutta Method [15 MIN]

- Euler's Method

- Higher order Runge-Kutta methods

- Demonstration, particles in a solenoid

Introducing Adaptive step size [10 MIN]

- Demonstration: magnetic dipole

- Dormand Prince 5 (4) method

Conclusion [2 MIN]

- Extra, the cyclotron

- Extra, toroidal fields

Introduction, what and why

- ▶ Numerical simulations are important.

Introduction, what and why

- ▶ Numerical simulations are important.
- ▶ Testing setups, non-analytical systems.

Introduction, what and why

- ▶ Numerical simulations are important.
- ▶ Testing setups, non-analytical systems.
- ▶ Demonstration, charged particles in electric and magnetic fields.
- ▶ Analytically known and not.

Introduction, what and why

- ▶ Numerical simulations are important.
- ▶ Testing setups, non-analytical systems.
- ▶ Demonstration, charged particles in electric and magnetic fields.
- ▶ Analytically known and not.
- ▶ Simulations are not experiments.

Theory: Classical non-relativistic particles

- ▶ Some repetition from Electrodynamics

Theory: Classical non-relativistic particles

- ▶ Some repetition from Electrodynamics
- ▶ The Lorentz force (SI units):

$$\vec{F} = q(\vec{v} \times \vec{B} + \vec{E}).$$

Theory: Classical non-relativistic particles

- ▶ Some repetition from Electrodynamics
- ▶ The Lorentz force (SI units):

$$\vec{F} = q(\vec{v} \times \vec{B} + \vec{E}).$$

- ▶ Only 1 particle! so pre-programmed depending on the setup.


Theory: Classical non-relativistic particles

- ▶ Some repetition from Electrodynamics
- ▶ The Lorentz force (SI units):

$$\vec{F} = q(\vec{v} \times \vec{B} + \vec{E}).$$

- ▶ Only 1 particle! so pre-programmed depending on the setup.
- ▶ Could use potentials $\phi(\vec{r}, t)$ $\vec{A}(\vec{r}, t)$ and Hamiltonian.


Known results, cyclotron motion \vec{B} fields



Known results, cyclotron motion \vec{B} fields

- Magnetic forces do no work:

$$dW_{\vec{B}} = \vec{F}_B \cdot d\vec{r} \propto (\vec{v} \times \vec{B}) \cdot \vec{v} = 0.$$




Known results, cyclotron motion \vec{B} fields

- Magnetic forces do no work:

$$dW_{\vec{B}} = \vec{F}_B \cdot d\vec{r} \propto (\vec{v} \times \vec{B}) \cdot \vec{v} = 0.$$

- $(\vec{v} = \vec{v}_\perp + \vec{v}_\parallel)$:

$$|\vec{F}_B| = |q(\vec{v} \times \vec{B})| = |qv_\perp B|.$$



Known results, cyclotron motion \vec{B} fields


- Magnetic forces do no work:

$$dW_{\vec{B}} = \vec{F}_B \cdot d\vec{r} \propto (\vec{v} \times \vec{B}) \cdot \vec{v} = 0.$$

- $(\vec{v} = \vec{v}_\perp + \vec{v}_\parallel)$:

$$|\vec{F}_B| = |q(\vec{v} \times \vec{B})| = |qv_\perp B|.$$

- Same as Centripetal force:
Cyclotron motion



Known results, cyclotron motion \vec{B} fields

- Magnetic forces do no work:


$$dW_{\vec{B}} = \vec{F}_B \cdot d\vec{r} \propto (\vec{v} \times \vec{B}) \cdot \vec{v} = 0.$$

- $(\vec{v} = \vec{v}_{\perp} + \vec{v}_{\parallel})$:


$$|\vec{F}_B| = |q(\vec{v} \times \vec{B})| = |qv_{\perp}B|.$$

- Same as Centripetal force:
Cyclotron motion
- Cyclotron radius and
frequency:

$$R = \frac{v_{\perp} m}{|q|B} \quad \omega_c = \frac{|q|B}{m}.$$




Analytical solution: Protons in a Solenoid



Solenoid with $N = 1000$ turns per m , $I = 5$ A, $r = 1$ m, $|\vec{B}| \approx 6$ mT.
Proton with $E_{kin} = 1$ MeV/ c^2 ($|v| \approx 3.195 \times 10^5$ m/s)


Analytical solution: Protons in a Solenoid




$$R \approx 0.5 \text{ m} \sin(\theta) \quad T = \frac{2\pi}{\omega_c} \approx 10 \mu\text{s}$$

Analytical solution: Protons in a Solenoid

Analytical: proton in a solenoid, side/front-view




Analytical: proton in a solenoid, speed



$$R \approx 0.5 \text{ m} \sin(\theta) \quad T = \frac{2\pi}{\omega_c} \approx 10 \mu\text{s}$$

Analytical solution: Protons in a Solenoid


Analytical: proton in a solenoid



$$R \approx 0.5 \text{ m} \sin(\theta) \quad T = \frac{2\pi}{\omega_c} \approx 10 \mu s$$


Analytical solution: Protons in a Solenoid

Analytical: proton in a solenoid, speed



$$R \approx 0.5 \text{ m} \sin(\theta) \quad T = \frac{2\pi}{\omega_c} \approx 10 \mu s$$

Analytical solution: Protons in a Solenoid



Ordinary differential equation's.

- ▶ Sources: Zeigler et al. Theory of Modeling and Simulation (Third edition) chapter 3
- ▶ Algorithms exists for ODEs:

$$\dot{\mathbf{X}} = f_{ode}(\mathbf{X}(t), t).$$

Ordinary differential equation's.

- ▶ Sources: Zeigler et al. Theory of Modeling and Simulation (Third edition) chapter 3
- ▶ Algorithms exists for ODEs:

$$\dot{\mathbf{X}} = f_{ode}(\mathbf{X}(t), t).$$

- ▶ We have a:

$$\ddot{\vec{r}} = \frac{q}{m}(\dot{\vec{r}} \times \vec{B}(\vec{r}, t) + \vec{E}(\vec{r}, t)).$$

Ordinary differential equation's.

- Sources: Zeigler et al. Theory of Modeling and Simulation (Third edition) chapter 3
- Algorithms exists for ODEs:

$$\dot{\mathbf{X}} = f_{ode}(\mathbf{X}(t), t).$$

- We have a:

$$\ddot{\vec{r}} = \frac{q}{m}(\dot{\vec{r}} \times \vec{B}(\vec{r}, t) + \vec{E}(\vec{r}, t)).$$

- Here:

$$\mathbf{X} = \begin{pmatrix} \vec{r} \\ \dot{\vec{r}} \end{pmatrix} \quad f_{ode}(\vec{r}, t) = \begin{pmatrix} \dot{\vec{r}} \\ \frac{q}{m}(\dot{\vec{r}} \times \vec{B}(\vec{r}, t) + \vec{E}(\vec{r}, t)) \end{pmatrix}.$$

The ODE to solve


```
auto ODE = [...](const state_type Data, state_type &
    dDatadt, const double t){
    //Extract position and velocity from data
    vec pos = vec(Data[0],Data[1],Data[2]);
    vec velocity = vec(Data[3],Data[4],Data[5]);

    //Lorentz+Newtons 2nd law
    vec F = Charge*(Fields.get_Efield(pos,t)+
        cross(velocity,Fields.get_Bfield(pos,t)));
    vec dVdt = F*Inv_mass;

    //Save derivative of data
    dDatadt[0]=velocity.x;
    ...
};
```

The Forward Euler's Method

- Let $h = t_{i+1} - t_i > 0$ be constant.




- Bernard P. Zeigler et al.
Theory of Modeling and

The Forward Euler's Method

- ▶ Let $h = t_{i+1} - t_i > 0$ be constant.
- ▶ h , $\mathbf{X}(t)$, t_i and f_{ode} are known.

$$\dot{\mathbf{X}} = f_{ode}(\mathbf{X}(t), t).$$




- ▶ Bernard P. Zeigler et al.
Theory of Modeling and

The Forward Euler's Method

- ▶ Let $h = t_{i+1} - t_i > 0$ be constant.
- ▶ h , $\mathbf{X}(t)$, t_i and f_{ode} are known.

$$\dot{\mathbf{X}} = f_{ode}(\mathbf{X}(t), t).$$

- ▶ How would you find $\mathbf{X}(t_{i+1})$:



- ▶ Bernard P. Zeigler et al.
Theory of Modeling and


The Forward Euler's Method

- ▶ Let $h = t_{i+1} - t_i > 0$ be constant.
- ▶ h , $\mathbf{X}(t)$, t_i and f_{ode} are known.

$$\dot{\mathbf{X}} = f_{ode}(\mathbf{X}(t), t).$$

- ▶ How would you find $\mathbf{X}(t_{i+1})$:
- ▶ (Explicit) Forward Euler's Method:

$$\mathbf{X}(t_{i+1}) = \mathbf{X}(t_i) + hf_{ode}(\mathbf{X}(t_i), t_i).$$




The Forward Euler's Method

- ▶ Let $h = t_{i+1} - t_i > 0$ be constant.
- ▶ h , $\mathbf{X}(t)$, t_i and f_{ode} are known.

$$\dot{\mathbf{X}} = f_{ode}(\mathbf{X}(t), t).$$

- ▶ How would you find $\mathbf{X}(t_{i+1})$:
- ▶ (Implicit) Backward Euler's Method:

$$\mathbf{X}(t_{i+1}) = \mathbf{X}(t_i) + hf_{ode}(\mathbf{X}(t_{i+1}), t_i).$$



The Forward Euler's Method


- ▶ Let $h = t_{i+1} - t_i > 0$ be constant.
- ▶ h , $\mathbf{X}(t)$, t_i and f_{ode} are known.

$$\dot{\mathbf{X}} = f_{ode}(\mathbf{X}(t), t).$$

- ▶ How would you find $\mathbf{X}(t_{i+1})$:
- ▶ (Explicit) Forward Euler's Method:

$$\mathbf{X}(t_{i+1}) = \mathbf{X}(t_i) + hf_{ode}(\mathbf{X}(t_i), t_i).$$

- ▶ Chain together multiple steps from t_0 ...with step size h .



- ▶ Bernard P. Zeigler et al

Why does this work? What is the error

- ▶ Multiple justifications for why.

Why does this work? What is the error

- ▶ Multiple justifications for why.
- ▶ First 2 terms in Taylor series Zeigler et al.:

$$\mathbf{X}(t_{i+1}) = \mathbf{X}(t_i) + hf_{ode}(\mathbf{X}(t_i), t_i) + h^2 \dots + \dots$$

Why does this work? What is the error

- ▶ Multiple justifications for why.
- ▶ First 2 terms in Taylor series Zeigler et al.:

$$\mathbf{X}(t_{i+1}) = \mathbf{X}(t_i) + hf_{ode}(\mathbf{X}(t_i), t_i) + h^2 \dots + \dots$$

- ▶ “Local truncation error” h^2 .
- ▶ Global error h^1 .

Why does this work? What is the error

- ▶ Multiple justifications for why.
- ▶ First 2 terms in Taylor series Zeigler et al.:

$$\mathbf{X}(t_{i+1}) = \mathbf{X}(t_i) + hf_{ode}(\mathbf{X}(t_i), t_i) + h^2 \dots + \dots$$

- ▶ “Local truncation error” h^2 .
- ▶ Global error h^1 .
- ▶ Convergence, but not uniform.

Why does this work? The Runge Kutta family

- In general.

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = \int_{t_i}^{t_{i+1}} f_{ode}(\mathbf{X}(t), t) dt.$$

- L. Zheng, X. Zhang, Modeling and Analysis of Modern Fluid Problems, 2017, chapter 8:

Why does this work? The Runge Kutta family

- In general.

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = \int_{t_i}^{t_{i+1}} f_{ode}(\mathbf{X}(t), t) dt = hf_{ode}(\mathbf{X}(\tau), \tau).$$

- *Mean Value theorem for integrals* $t_i \leq \tau \leq t_{i+1}$.

- L. Zheng, X. Zhang, Modeling and Analysis of Modern Fluid Problems, 2017, chapter 8:

Why does this work? The Runge Kutta family

- In general.

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = \int_{t_i}^{t_{i+1}} f_{ode}(\mathbf{X}(t), t) dt = hf_{ode}(\mathbf{X}(\tau), \tau).$$

- *Mean Value theorem for integrals* $t_i \leq \tau \leq t_{i+1}$.
- Guess $\tau = t_j$.

- L. Zheng, X. Zhang, Modeling and Analysis of Modern Fluid Problems, 2017, chapter 8:

Why does this work? The Runge Kutta family

- In general.

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = \int_{t_i}^{t_{i+1}} f_{ode}(\mathbf{X}(t), t) dt = hf_{ode}(\mathbf{X}(\tau), \tau).$$

- Mean Value theorem for integrals $t_i \leq \tau \leq t_{i+1}$.
- Guess $\tau = t_j$.
- More generally, (*Explicit* and *single step*), Runge-Kutta family:

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = \int_{t_i}^{t'} f_{ode}(\mathbf{X}(t), t) dt + \dots \int_{t^{(m)}}^{t_{i+1}} f_{ode}(\mathbf{X}(t), t) dt$$

- L. Zheng, X. Zhang, Modeling and Analysis of Modern Fluid Problems, 2017, chapter 8:

Why does this work? The Runge Kutta family

- In general.

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = \int_{t_i}^{t_{i+1}} f_{ode}(\mathbf{X}(t), t) dt = hf_{ode}(\mathbf{X}(\tau), \tau).$$

- Mean Value theorem for integrals $t_i \leq \tau \leq t_{i+1}$.
- Guess $\tau = t_i$.
- More generally, (*Explicit and single step*), Runge-Kutta family:

$$\begin{aligned}\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) &= \int_{t_i}^{t'} f_{ode}(\mathbf{X}(t), t) dt + \dots \int_{t^{(m)}}^{t_{i+1}} f_{ode}(\mathbf{X}(t), t) dt \\ &= \sum_{j=1}^m h_j f_{ode}(\mathbf{X}(\tau_j), \tau_j)\end{aligned}$$

- Use $f_{ode}(\mathbf{X}(t_i), t_i)$ to approximate $\mathbf{X}(\tau_1)$ etc.
- L. Zheng, X. Zhang, Modeling and Analysis of Modern Fluid Problems, 2017, chapter 8:

Explicit Runge Kutta methods

- We want:

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j \mathbf{K}_j$$

- Martha L. Abell, James P. Braselton, Differential Equations with Mathematica (Fourth Edition), 2016:

Explicit Runge Kutta methods

- We want:

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j \mathbf{K}_j$$

- Want exact to p 'th order. Can be found with Taylor expansion of $\mathbf{X}(t_i)$.

- Martha L. Abell, James P. Braselton, Differential Equations with Mathematica (Fourth Edition), 2016:

Explicit Runge Kutta methods

- We want:

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j \mathbf{K}_j$$

- Want exact to p 'th order. Can be found with Taylor expansion of $\mathbf{X}(t_i)$.
- Local/global error h^{p+1} , Global error h^p .

- Martha L. Abell, James P. Braselton, Differential Equations with Mathematica (Fourth Edition), 2016:

Explicit Runge Kutta methods

- We want:

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j \mathbf{K}_j$$

- Want exact to p 'th order. Can be found with Taylor expansion of $\mathbf{X}(t_i)$.
- Local/global error h^{p+1} , Global error h^p .
- 2nd order (Heun's method):

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = \frac{h}{2}(\mathbf{k}_1 + \mathbf{k}_2)$$

$$\mathbf{k}_1 = f_{ode}(\mathbf{X}(t_i), t_i)$$

$$\mathbf{k}_2 = f_{ode}(\mathbf{X}(t_i) + h\mathbf{k}_1, t_i + h)$$

- Martha L. Abell, James P. Braselton, Differential Equations with Mathematica (Fourth Edition), 2016:

The 4th order Runge Kutta method

- RK4, often simply called the Runge Kutta method:


$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = \frac{h}{6}(\mathbf{k}_1 + 2\mathbf{k}_2 + 2\mathbf{k}_3 + \mathbf{k}_4)$$

$$\mathbf{k}_1 = f_{ode}(\mathbf{X}(t_i), t_i)$$

$$\mathbf{k}_2 = f_{ode}\left(\mathbf{X}(t_i) + \frac{h}{2}\mathbf{k}_1, t_i + \frac{h}{2}\right)$$

$$\mathbf{k}_3 = f_{ode}\left(\mathbf{X}(t_i) + \frac{h}{2}\mathbf{k}_2, t_i + \frac{h}{2}\right)$$

$$\mathbf{k}_4 = f_{ode}(\mathbf{X}(t_i) + h\mathbf{k}_3, t_i + h)$$



Wikipedia-user HilberTraum,
published under creative
commons: CC BY-SA 4.0

The General explicit Runge Kutta method

- General explicit, single step, fixed size, Runge Kutta method

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j \mathbf{K}_j$$

$$\mathbf{k}_1 = f_{ode}(\mathbf{X}(t_i), t_i)$$

$$\mathbf{k}_2 = f_{ode}(\mathbf{X}(t_i) + ha_{21}\mathbf{k}_1, t_i + c_2h)$$

$$\mathbf{k}_3 = f_{ode}(\mathbf{X}(t_i) + ha_{31}\mathbf{k}_1 + ha_{32}\mathbf{k}_2, t_i + c_3h) \quad \vdots$$

The General explicit Runge Kutta method

- General explicit, single step, fixed size, Runge Kutta method

$$\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j \mathbf{K}_j$$

$$\mathbf{k}_1 = f_{ode}(\mathbf{X}(t_i), t_i)$$

$$\mathbf{k}_2 = f_{ode}(\mathbf{X}(t_i) + ha_{21}\mathbf{k}_1, t_i + c_2h)$$

$$\mathbf{k}_3 = f_{ode}(\mathbf{X}(t_i) + ha_{31}\mathbf{k}_1 + ha_{32}\mathbf{k}_2, t_i + c_3h) \quad \vdots$$

- Expressed in Butcher tableau:

$c_1 = 0$			
c_2	a_{21}		
c_3	a_{31}	a_{32}	
c_n	a_{n1}	a_{n2}	\dots
<hr/>			
	b_1	b_2	\dots

Euler Implementations

```
state_type Data = Data0;
state_type dDatadt;
size_t time_res = T/timestep;
for (size_t i = 1; i < time_res; ++i)
{
    double t=i*dt;
    ODE(Data,dDatadt,t);
    //Euler time evolution
    Data+=timestep*dDatadt;

    save_step( Data , i*timestep );
};
```


RK4 Implementations (1/2)

```
state_type Data = Data0;
state_type temp=Data0;
state_type K1,K2,K3,K4;
size_t time_res = T/timestep;
for (size_t i = 1; i < time_res; ++i)
{
    double t=i*timestep;

    //substep 1
    ODE(Data,K1,t);

    //substep 2
    temp=Data+timestep*K1/2;
    ODE(temp,K2,t+timestep/2);
```

RK4 Implementations (2/2)

```
//substep 3
```

```
temp=Data+timestep*K2/2;  
ODE(temp,K3,t+timestep/2);
```

```
//substep 4
```

```
temp=Data+timestep*K3;  
ODE(temp,K4,t+timestep);
```

```
//Read data
```

```
Data+=timestep*(K1+2.0*K2+2.0*K3+K4)/6.0;  
save_step( Data , i*timestep );
```

```
}
```

“Correct” way

```
#include <boost/array.hpp>
#include <boost/numeric/odeint.hpp>
using namespace boost::numeric::odeint;
typedef boost::array< double, 6 > state_type;
...
size_t steps = integrate_const(
    runge_kutta4< state_type >(),
    ODE,    //Lorentz-force
    Data0 ,//{pos0,v0}
    0.0 ,   //t0=0
    T ,     //max time
    timestep ,//length of each step
    save_step //User defined save data function
);
```

Does it work

- ▶ Test, same proton in a solenoid use $\theta = 60^\circ$ reference, had:

$$R \approx 0.5 \text{ m} \sin(\theta) \approx 0.45 \text{ m} \quad T = \frac{2\pi}{\omega_c} \approx 10 \mu\text{s}$$

Does it work

- ▶ Test, same proton in a solenoid use $\theta = 60^\circ$ reference, had:

$$R \approx 0.5 \text{ m} \sin(\theta) \approx 0.45 \text{ m} \quad T = \frac{2\pi}{\omega_c} \approx 10 \mu\text{s}$$

- ▶ Compare Analytic, Euler, Runge-Kutta 4.

Does it work

- ▶ Test, same proton in a solenoid use $\theta = 60^\circ$ reference, had:

$$R \approx 0.5 \text{ m} \sin(\theta) \approx 0.45 \text{ m} \quad T = \frac{2\pi}{\omega_c} \approx 10 \mu\text{s}$$

- ▶ Compare Analytic, Euler, Runge-Kutta 4.
- ▶ Consider $\theta = 60^\circ$ at different h .

Does it work


- ▶ Test, same proton in a solenoid use $\theta = 60^\circ$ reference, had:

$$R \approx 0.5 \text{ m} \sin(\theta) \approx 0.45 \text{ m} \quad T = \frac{2\pi}{\omega_c} \approx 10 \mu\text{s}$$

- ▶ Compare Analytic, Euler, Runge-Kutta 4.
- ▶ Consider $\theta = 60^\circ$ at different h .
- ▶ Check error on $|\vec{v}|$, $R = \sqrt{y^2 + z^2}$ and $x(t)$.

At a glance, 3D view


$$h = t_{i+1} - t_i = 0.01 \mu\text{s}$$



3129 steps

At a glance, 3D view


$$h = t_{i+1} - t_i = 0.1 \mu\text{s}$$



312 steps

At a glance, 3D view


$$h = t_{i+1} - t_i = 1.0 \mu\text{s}$$



31 steps.


At a glance, front view, no border

Euler method, front-view




At a glance, front view, no border


RK4, front-view




Constant radius?




Constant radius?




Constant speed?



Constant speed?



Error as function of h



Adaptive step size, why

- ▶ h must be small “enough”

Adaptive step size, why

- ▶ h must be small “enough”
- ▶ Hard to pick, and may change:

Adaptive step size, why

- ▶ h must be small “enough”
- ▶ Hard to pick, and may change:
- ▶ Inhomogeneous fields (here Earth magnetic field)





Illustration originally from Nasa.
Published on wikipedia, in Public Domain

Adaptive step size, why

- ▶ h must be small “enough”
- ▶ Hard to pick, and may change:
- ▶ Inhomogeneous fields (here Earth magnetic field)
- ▶ time dependent fields (here the cyclotron, bad example)



Ernest O. Lawrence, 1934, U.S. Patent 1,948,384; image in Public Domain.


Adaptive step size, why

- ▶ h must be small “enough”
- ▶ Hard to pick, and may change:
- ▶ Inhomogeneous fields (here Earth magnetic field)
- ▶ time dependent fields (here the cyclotron, bad example)
- ▶ Let the computer pick h .

Example, magnetic dipole

- True dipole:

$$\vec{B} = \frac{\mu_0}{4\pi r^3} \left[3\hat{r}(\vec{m} \cdot \hat{r}) - \vec{m} \right].$$




Example, magnetic dipole

- True dipole:

$$\vec{B} = \frac{\mu_0}{4\pi r^3} \left[3\hat{r}(\vec{m} \cdot \hat{r}) - \vec{m} \right].$$

- Weak field, little change, long steps.



Arbitrarily set

$$\frac{\mu_0}{4\pi} |\vec{m}| = 0.155 \text{ T/m}^3.$$


Protons with speed around
10 000 m/s

Example, magnetic dipole

- True dipole:

$$\vec{B} = \frac{\mu_0}{4\pi r^3} \left[3\hat{r}(\vec{m} \cdot \hat{r}) - \vec{m} \right].$$

- Weak field, little change, long steps.
- strong field, large change, short steps.



Arbitrarily set

$$\frac{\mu_0}{4\pi} |\vec{m}| = 0.155 \text{ T/m}^3.$$


Protons with speed around
10 000 m/s

Example, magnetic dipole

- ▶ True dipole:

$$\vec{B} = \frac{\mu_0}{4\pi r^3} \left[3\hat{r}(\vec{m} \cdot \hat{r}) - \vec{m} \right].$$

- ▶ Weak field, little change, long steps.
- ▶ strong field, large change, short steps.
- ▶ No analytical solution (afaik.)



Arbitrarily set

$$\frac{\mu_0}{4\pi} |\vec{m}| = 0.155 \text{ T/m}^3.$$

Protons with speed around
10 000 m/s

Runge-Kutta Adaptive step size, how

- ▶ Reduce h until the “error” is small enough

Runge-Kutta Adaptive step size, how

- ▶ Reduce h until the “error” is small enough
- ▶ Use 2 different order methods, $\mathbf{x}^{(p+1)}(t_{i+1})$, $\mathbf{x}^{(p)}(t_{i+1})$.

Runge-Kutta Adaptive step size, how

- ▶ Reduce h until the “error” is small enough
- ▶ Use 2 different order methods, $\mathbf{x}^{(p+1)}(t_{i+1})$, $\mathbf{x}^{(p)}(t_{i+1})$.
- ▶ Approximate “Error” as $\mathbf{E} = |\mathbf{x}^{(p+1)}(t_{i+1}) - \mathbf{x}^{(p)}(t_{i+1})|$.

Runge-Kutta Adaptive step size, how

- ▶ Reduce h until the “error” is small enough
- ▶ Use 2 different order methods, $\mathbf{x}^{(p+1)}(t_{i+1})$, $\mathbf{x}^{(p)}(t_{i+1})$.
- ▶ Approximate “Error” as $\mathbf{E} = |\mathbf{x}^{(p+1)}(t_{i+1}) - \mathbf{x}^{(p)}(t_{i+1})|$.
- ▶ Adjust step size to keep the error(s) small (Implementations differ!).

Dormand, J. R.; Prince, P. J. (1980), "A family of embedded Runge-Kutta formulae", Journal of Computational and Applied Mathematics

(1)

Runge-Kutta Dormand Prince 5 (4)

- ode45 in Matlab, `scipy.solve_ivp` in Python ,
`RungeKutta_dopri5` in `boost::odeint`.

$$\mathbf{X}^{(5)}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j^{(5)} \mathbf{k}_j$$

$$\mathbf{X}^{(4)}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j^{(4)} \mathbf{k}_j.$$

$$\mathbf{k}_i = f_{ode}(\mathbf{X}(t_i) + h \sum_j^{i-1} a_{ki} \mathbf{k}_i, t_i + c_3 h).$$

Runge-Kutta Dormand Prince 5 (4)

- ▶ ode45 in Matlab, `scipy.solve_ivp` in Python ,
`RungeKutta_dopri5` in `boost::odeint`.

$$\mathbf{X}^{(5)}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j^{(5)} \mathbf{k}_j$$

$$\mathbf{X}^{(4)}(t_{i+1}) - \mathbf{X}(t_i) = h \sum_{j=1}^m b_j^{(4)} \mathbf{k}_j.$$

$$\mathbf{k}_i = f_{ode}(\mathbf{X}(t_i) + h \sum_j^{i-1} a_{ji} \mathbf{k}_j, t_i + c_3 h).$$

- ▶ 7 \mathbf{k}_i 's (actually 6).
Dormand, J. R.; Prince, P. J. (1980), "A family of embedded Runge-Kutta formulae", Journal of Computational and Applied Mathematics

Butcher Tableau of Dormand Prince (5) 4

c_i	a_{ij}	...						
0								
$\frac{1}{5}$	$\frac{1}{5}$							
$\frac{3}{10}$	$\frac{3}{40}$	$\frac{9}{40}$						
$\frac{4}{5}$	$\frac{40}{45}$	$-\frac{56}{15}$	$-\frac{32}{9}$					
$\frac{5}{8}$	$\frac{19372}{6561}$	$-\frac{25360}{2187}$	$\frac{64448}{6561}$	$-\frac{212}{729}$				
$\frac{9}{8}$	$\frac{9017}{3168}$	$-\frac{355}{33}$	$\frac{46732}{5247}$	$\frac{49}{176}$	$-\frac{5103}{18656}$			
1	$\frac{35}{384}$	0	$\frac{500}{1113}$	$\frac{125}{192}$	$-\frac{2187}{6784}$	$\frac{11}{84}$		
$/b^{(5)}$	$\frac{35}{384}$	0	$\frac{500}{1113}$	$\frac{125}{192}$	$-\frac{2187}{6784}$	$\frac{11}{84}$	0	
$/b^{(4)}$	$\frac{5179}{57600}$	0	$\frac{7571}{16695}$		$\frac{393}{640}$	$-\frac{92097}{339200}$	$\frac{187}{2100}$	$\frac{1}{40}$

Runge-Kutta Dormand Prince 5 (4)

- Step size correction (Dormand Prince):

$$h_{new} = 0.9h_{old} \left[\frac{\delta}{||E||} \right]^{\frac{1}{p+1}}$$

Runge-Kutta Dormand Prince 5 (4)

- Step size correction (Dormand Prince):

$$h_{new} = 0.9h_{old} \left[\frac{\delta}{||E||} \right]^{\frac{1}{p+1}}$$

- Step size correction (Me):

$$h_{new} = \min_j \left(0.9h_{old} \left[\frac{\delta_j}{E_j} \right]^{\frac{1}{p}} \right), \quad E_j > \delta_j : \text{reject}$$

$$\delta_j = \min(\delta_{abs}, |\mathbf{X}_j(t_i)|\delta_{rel}) \sqrt{\frac{h_{old}}{T}}.$$

Runge-Kutta Dormand Prince 5 (4)

- Step size correction (Dormand Prince):

$$h_{new} = 0.9h_{old} \left[\frac{\delta}{||E||} \right]^{\frac{1}{p+1}}$$


- Step size correction (Me):

$$h_{new} = \min_j \left(0.9h_{old} \left[\frac{\delta_j}{E_j} \right]^{\frac{1}{p}} \right), \quad E_j > \delta_j : \text{reject}$$

$$\delta_j = \min(\delta_{abs}, |\mathbf{X}_j(t_i)|\delta_{rel}) \sqrt{\frac{h_{old}}{T}}.$$


- Relative and absolute error at end. “Fail safe” $\sqrt{\dots}$.
Dormand, J. R.; Prince, P. J. (1980), "A family of embedded Runge-Kutta formulae", Journal of Computational and Applied Mathematics

Does it work?




My version relative and absolute error 10^{-6} . 94 steps (+ 14 rejected).

Does it work?




Odeint library relative and absolute error 10^{-7} . 92 steps.

Does it work?




Does it work?

dynamic timesteps




Another curious result




$T = 0.2 \text{ ms}$

Another curious result



$T = 0.3 \text{ ms}$


Another curious result



$$T = 0.6 \text{ ms}$$


Magnetic “mirror” or “bottle”.

Another curious result



Wikipedia user WikiHelper2134 , Public Domain.

Another curious result



Tandem Mirror Experiment The Lawrence Livermore National

Extra examples-“if time permits”

- ▶ The cyclotron
 - ▶ When the Dormand Prince method fails

Extra examples-“if time permits”

- ▶ The cyclotron
 - ▶ When the Dormand Prince method fails
- ▶ Toroidal fields [TBD]
 - ▶ Another non-analytic setup.

Conclusion

- ▶ Simulating particles in electric and magnetic fields.

Conclusion


- ▶ Simulating particles in electric and magnetic fields.
- ▶ Numerically solving ordinary differential equations
- ▶ Reasonable agreement with known results.
- ▶ Can easily be generalized to other systems.

Conclusion

- ▶ Simulating particles in electric and magnetic fields.
- ▶ Numerically solving ordinary differential equations
- ▶ Reasonable agreement with known results.
- ▶ Can easily be generalized to other systems.
- ▶ Limitation, simulations are not experiments.

Example, cyclotron


- Electric field accelerates, magnetic contains.



Ernest O. Lawrence, 1934, U.S. Patent 1,948,384; image in Public Domain.

Example, cyclotron


- ▶ Electric field accelerates, magnetic contains.
- ▶ Single gap, oscillating field.



Ernest O. Lawrence, 1934, U.S.
Patent 1,948,384; image in
Public Domain.

Example, cyclotron

- ▶ Electric field accelerates, magnetic contains.
- ▶ Single gap, oscillating field.
- ▶ Uses classical Cyclotron frequency




Ernest O. Lawrence, 1934, U.S.
Patent 1,948,384; image in
Public Domain.

Example, cyclotron

- ▶ Electric field accelerates, magnetic contains.
- ▶ Single gap, oscillating field.
- ▶ Uses classical Cyclotron frequency
- ▶ Analytical final speed, in principle path.


$$\frac{R|q|B}{m} = v_{\perp}$$



Ernest O. Lawrence, 1934, U.S. Patent 1,948,384; image in Public Domain.

can it be simulated


- ▶ with fixed step size, looks bad 4999 points



4999 points.

can it be simulated


- ▶ with fixed step size, looks bad 4999 points
- ▶ my adabtive method, error: 10^{-6}



500000 points.

can it be simulated


- ▶ with fixed step size, looks bad 4999 points
- ▶ my adabtive method, error: 10^{-6}
- ▶ odeint library



2070 points.

can it be simulated

- ▶ with fixed step size, looks bad 4999 points
- ▶ my adabtive method, error: 10^{-6}
- ▶ odeint library
- ▶ non-continuous ode are bad.



2070 points.

TBD