

University of Thessaly



Neuro-Fuzzy Computing

ECE447

1st Problem Set

Alexandra Gianni Nikos Stylianou

ID: 3382

ID: 2917

December 13, 2023

1 Problem 1

The contour lines of $f(x, y)$ are plotted with the following MATLAB code and are presented in figure 1.

```
function [Z] = plot_contour(start_num , end_num)

    x = linspace(start_num , end_num , 100);
    y = x;
    [X, Y] = meshgrid(x, y);
    Z = X.^2 + 4*X.*Y + Y.^2;
    contour(X, Y, Z, 40);
    xlabel('X');
    ylabel('Y');
end
```




Figure 1: Contour lines of $f(x, y)$

A general formula of a quadratic equation is $f(x, y) = ax^2 + 2bxy + cy^2$. Writing our formula in the previous form, we find that $a = 1$, $b = 2$, $c = 1$. Calculation of the discriminant can help us calculate the location of the function's local minimum/maximum.

$$D = \begin{bmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{bmatrix} = f_{xx}f_{yy} - f_{xy}^2 = 2 \times 2 - 4^2 = -12 < 0, \quad \text{where} \quad (1)$$

$$f_{xx} = \frac{\partial^2 f}{\partial x^2} = 2, \quad f_{yy} = \frac{\partial^2 f}{\partial y^2} = 2, \quad f_{xy} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) = 4D = 2 \times 2 - 4^2 = -12 < 0.$$

So, we only have to find the point at which $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ are equal to 0. This point will be a saddle point at which the gradients in each orthogonal direction are 0, but this point is neither a local minimum nor a maximum. More precisely:

$$\begin{cases} \frac{\partial f}{\partial x} = 2x + 4y = 0 \\ \frac{\partial f}{\partial y} = 4x + 2y = 0 \end{cases} \Rightarrow \begin{cases} x = 0 \\ y = 0 \end{cases} \quad (2)$$

Thus, the point $(x, y) = (0, 0)$ is the saddle point mentioned for the given function and this can be justified using the plotted contour lines.

2 Problem 4

In this problem, we will express the derivatives in respect to the matching activation function.

2.1 LogSig

This activation function is expressed as

$$S(x) = \frac{1}{1 + e^{-x}}$$

Multiplying itself with $(1 + e^{-x})$ gives

$$(1 + e^{-x}) S(x) = 1 \Leftrightarrow e^{-x} = \frac{1}{S(x)} - 1$$

So, activation function's derivative will be

$$\begin{aligned} \frac{dS}{dx} &= \frac{d\left((1 + e^{-x})^{-1}\right)}{dx} = (1 + e^{-x})^{-2} e^{-x} = S^2(x) \left(\frac{1}{S(x)} - 1\right) \\ &= S(x) - S^2(x) = S(x) (1 - S(x)) \end{aligned}$$

2.2 TanSig

Activation function is

$$S(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$$

Its derivative is

$$\begin{aligned} \frac{dS}{dx} &= \frac{(e^x + e^{-x})(e^x + e^{-x}) - (e^x - e^{-x})(e^x - e^{-x})}{(e^x + e^{-x})^2} = \\ &= \frac{(e^x + e^{-x})^2 - (e^x - e^{-x})^2}{(e^x + e^{-x})^2} = 1 - \left(\frac{e^x - e^{-x}}{e^x + e^{-x}}\right)^2 = 1 - S^2(x) \end{aligned} \quad (3)$$

2.3 Swish

Activation function is

$$S(x) = \frac{x}{1 + e^{-x}}$$

The derivative in respect to x is

$$\frac{dS}{dx} = \frac{1 + e^{-x} + xe^{-x}}{(1 + e^{-x})^2} = \frac{1 + e^{-x}}{(1 + e^{-x})^2} + \frac{xe^{-x}}{(1 + e^{-x})^2} = \frac{1}{1 + e^{-x}} + x \frac{e^{-x}}{(1 + e^{-x})^2}$$

Rewriting the function gives us

$$\frac{S(x)}{x} = \frac{1}{1 + e^{-x}} \quad \text{and} \quad e^{-x} = \frac{x - S(x)}{S(x)}$$

So, continuing with the derivative:

$$\begin{aligned}\frac{dS}{dx} &= \frac{1}{1+e^{-x}} + x \frac{e^{-x}}{(1+e^{-x})^2} = \frac{S(x)}{x} + \frac{e^{-x}}{1+e^{-x}} \frac{x}{1+e^{-x}} = \frac{S(x)}{x} + S(x) \frac{x}{1+e^{-x}} = \\ &= \frac{S(x)}{x} + S(x) \frac{S(x)}{x} e^{-x} = \frac{S(x)}{x} (1 + S(x)e^{-x}) = \frac{S(x)}{x} \left(1 + S(x) \frac{x - S(x)}{S(x)} \right) = \\ &= \frac{S(x)}{x} (1 + x - S(x)) = \frac{S(x)}{x} + S(x) - \frac{S^2(x)}{x}\end{aligned}$$

Let $\sigma = \frac{1}{1+e^{-x}}$, thus

$$\frac{dS}{dx} = \frac{S(x)}{x} + S(x) - \frac{S^2(x)}{x} = \sigma + S(x) - \sigma S(x) = S(x) + \sigma(1 - S(x)) \quad (4)$$

2.4 Custom tanh

Activation function is

$$S(x) = x \times \tanh(\ln(1 + e^x)) = \frac{x(1 + e^x) - \frac{x}{1 + e^x}}{1 + e^x + \frac{1}{1 + e^x}}$$

After some calculations, we can observe a relationship to the Swish function:

$$S(x) = \frac{x(1 + e^x) - T(-x)}{1 + e^x + \frac{1}{1 + e^x}} = \frac{\frac{-x^2}{T(-x)} - T(-x)}{\frac{-x}{T(-x)} + \frac{T(-x)}{x}} = \frac{-x^3 - T^2(-x)}{-x^2 + T^2(-x)}$$

3 Problem 8

ADALINE is a single-layer artificial neural network that can learn and adapt to non-linear relationships between inputs and outputs.

It consists of a single neuron with a linear activation function. Each input of the neuron has a corresponding weight, which is adapted during training to minimize the error between the network's output and the desired output. To adjust the weights the algorithm uses the learning rule α .

Suppose that we have the following three reference patterns and their targets:

$$\left\{ p_1 = \begin{bmatrix} 2 \\ 4 \end{bmatrix}, t_1 = [26] \right\} \quad \left\{ p_2 = \begin{bmatrix} 4 \\ 2 \end{bmatrix}, t_2 = [26] \right\} \quad \left\{ p_3 = \begin{bmatrix} -2 \\ -2 \end{bmatrix}, t_3 = [-26] \right\}$$

The probability of vector p_1 is $P_1 = 0.20$, the probability of vector p_2 is $P_2 = 0.70$, and the probability of vector p_3 is $P_3 = 0.10$.

3.1 Question a

The number of inputs to an ADALINE network for any neural network is determined by the dimensionality of our data, not the number of data points (patterns) that we have. In our case, each pattern is a 2-dimensional vector. Therefore, our ADALINE network has two inputs, one for each dimension.

In an ADALINE network, the number of weights is equal to the number of inputs. In our case, each pattern is a 2-dimensional vector. Thus, our neural network will have two weights, one for each dimension of the input.

4 Problem 11

Fuzzy logic is a type of logic that deals with vague, imprecise, or uncertain information. It is based on the concept of fuzzy sets, which are sets that can have any degree of membership between 0 and 1. The value zero is used to represent complete non-membership, the value one is used to represent complete membership, and values in between are used to represent intermediate degrees of membership. This means that an element can be a member of a fuzzy set to some degree, rather than all or nothing.

The uniqueness of fuzzy logic is that fuzzy logic can handle imprecise and uncertain information, which makes it a valuable tool for dealing with real-life problems that are inherently vague or fuzzy.

On this exercise, we are dealing with the linguistic variable *Truth* with a possible membership set:

$$T = \text{Absolutely false, Very false, False, Fairly true, True, Very true, Absolutely true}$$

Based on that set we may define the membership function of truth as:

$$\text{True}(u) = u \quad \text{False}(u) = 1-u$$

for each $u \in [0, 1]$.

5 Problem 12

In order to evaluate the expression "not (A(x) OR B(x))", we must first take a look at how fuzzy logic differs from binary logic at operation level. In binary logic we have three basic operations: AND(x, y), OR(x, y) and NOT(x). But, in fuzzy logic, where a function can have a value in the range of [0...1], things are slightly different. The binary operation AND(x, y) is equivalent to MIN(x, y) from fuzzy logic, OR(x, y) to MAX(x, y) and NOT(x) to 1-x.

We need to find the proper x for which the previous expression has the maximum value. First, we calculate the expression and then find the correct x . To achieve this, we need to divide our calculations into ranges. Starting for $x \leq 2$, "A(x) AND B(x)" is equal to " $\max(A(x), B(x))$ " = 1. Applying De Morgan's law, we therefore have $\max(A(x), B(x)) \Rightarrow \text{not}(A(x) \text{ or } B(x)) = 0$. Exactly the same result is obtained with $x \geq 7$.

Things are a bit different in $2 \leq x \leq 7$. The function $A(x)$ starts to fall while $B(x)$ starts to rise. The point at which the two functions cross is important for the definition of the required expression and can be obtained by solving the equation:

$$A(x_{crit}) = B(x_{crit}) \Leftrightarrow 1 - \frac{x_{crit} - 2}{3} = \frac{x_{crit} - 3}{4} \Rightarrow x_{crit} = \frac{29}{7}$$

For $2 \leq x \leq \frac{29}{7}$, $\max(A(x), B(x)) = 1 - \frac{x-2}{3} = A(x)$, because in this region $A(x)$ lies above $B(x)$. Thus, $\text{not}(A(x) \text{ or } B(x)) = \frac{x-2}{3}$.

Using the same logic, we find out that for $\frac{29}{7} \leq x \leq 7$, $\text{not}(A(x) \text{ or } B(x)) = 1 - \frac{x-3}{4}$.

Therefore, the expression $f(x) = \text{not}(A(x) \text{ or } B(x))$ is summarized below:

$$f(x) = \begin{cases} 0 & x \leq 2, \\ \frac{x-2}{3} & 2 \leq x \leq \frac{29}{7}, \\ 1 - \frac{x-3}{4} & \frac{29}{7} \leq x \leq 7, \\ 0 & x \geq 7 \end{cases} \quad (5)$$




Figure 2: Plot of A(x), B(x)

By plotting this function in figure 3, we can clearly see that the maximum occurs at $x = x_{crit} = \frac{29}{7}$ and its value is 0.715465.




Figure 3: Expression's plot