# A very brief introduction to the Einstein Toolkit (Open-source codes)

3rd HEL.A.S. Summer School and DAAD school for Neutron Stars and Gravitational Waves

Theodoros Soultanis

IMPRS PhD student

at

Max Planck Institute for Astronomy, Heidelberg (MPIA)
Heidelberg Institute for Theoretical Studies (HITS)

Email: theodoros. soultanis@h-its.org

October 9, 2018









## Outline

Einstein Toolkit: General information

Setting up a simulation: Jupyter notebook










### Einstein Toolkit: What is it?

Einstein Toolkit, Link: https://einsteintoolkit.org/

- A three-dimensional code for the dynamical evolution of spacetimes in full General Relativity (BBH, BNS, Isolated stars)
- Open source, community-driven, and freely accessible infrastructure with highly active users
- Consists of diverse software components contributed by many different people and used by many more (210 members from 129 different groups and 36 countries)






## Einstein Toolkit: Structure

- Highly modular code which supports C, C++, FORTRAN, OpenCL, CUDA.
- Built on the Cactus framework, which provides the basic infrastructure for the numerical simulations (the modules in Einstein Toolkit's language are called Thorns)
- Thorns are developed independently (interchangeable with others)
  - Usually do not directly interact with each other
  - Each of them interacts with the Cactus framework
- Documentation is tricky





• Implements an Adaptive Mesh Refinement

Cactus code: http://cactuscode.org/










## Einstein Toolkit: Structure



 Scalability tested on HPC infrastructure (Galileo, ARIS)

Scalability tests for the Einstein Toolkit









# Einstein Toolkit: 3+1 decomposition

- The McLachlan module implements the dynamical evolution of spacetime in the BSSNOK formalism
- The code is auto-generated from tensor equations via Kranc

#### **Definitions**

$$\begin{split} \widetilde{\gamma}_{ij} &= \mathrm{e}^{-4\phi} \gamma_{ij} \\ \mathrm{e}^{4\phi} &= \gamma^{1/3} \\ A_{ij} &= K_{ij} - \frac{1}{3} \gamma_{ij} K \\ \widetilde{A}_{ij} &= \mathrm{e}^{-4\phi} A_{ij} \\ \widetilde{\Gamma}^i &= \widetilde{\gamma}^{jk} \widetilde{\Gamma}^i_{jk} = -\partial_j \widetilde{\gamma}^{ij} \end{split}$$

#### Time evolution

$$\begin{split} &\frac{d}{dt}\widetilde{\gamma}_{ij} = -2\widetilde{A}_{ij}, \quad \frac{d}{dt}\phi = -\frac{1}{6}\alpha K, \quad \frac{d}{dt} = \partial_t - \mathcal{L}_\beta \\ &\frac{d}{dt}K = -\gamma^{ij}D_jD_i\alpha + \alpha(\widetilde{A}_{ij}\widetilde{A}^{ji} + \frac{1}{3}K^2) + 4\pi\alpha(\rho + S) \\ &\frac{d}{dt}\widetilde{A}_{ij} = e^{-4\phi}\left(-(D_jD_i\alpha)^{TF} + \alpha(R_{ij}^{TF} - 8\pi S_{ij}^{TF})\right) + \alpha(K\widetilde{A}_{ij} - 2\widetilde{A}_{il}\widetilde{A}_j^l) \\ &\partial_t\widetilde{\Gamma}^i = -2\widetilde{A}^{ij}\partial_j\alpha + 2\alpha\left(\widetilde{\Gamma}^i{}_{jk}\widetilde{A}^{kj} - \frac{2}{3}\widetilde{\gamma}^{ij}\partial_jK - 8\pi\widetilde{\gamma}^{ij}S_j + 6\widetilde{A}^{ij}\partial_j\phi\right) \\ &+ \beta^k\partial_j\widetilde{\Gamma}^i - \widetilde{\Gamma}^j\partial_j\beta^i + \frac{2}{3}\widetilde{\Gamma}^i\partial_j\beta^j + \frac{1}{3}\widetilde{\gamma}^{li}\partial_l\partial_j\beta^j + \widetilde{\gamma}^{lj}\partial_j\partial_l\beta^i \end{split}$$

$$\frac{d\alpha}{dt} = -2\alpha K$$

#### Gamma driver

$$\partial_t \beta^i = kB^i$$
$$\partial_t B^i = \partial_t \widetilde{\Gamma}^i - \eta B^i$$

# Einstein Toolkit: Hydrodynamics

- The GRHydro module implements a grid based High Resolution Shock Capturing Scheme (HRSC) for the evolution of matter
- It supports GRHD, GRMHD (Valencia formulation)
- The EOS\_Omni module provides an interface for all the EOS calls
  - polytropes, gamma law, piecewise polytropes, hybrid

- Hydro\_RNSID generates initial configurations of rapidly (uniformly or differentially) rotating stars for the GRHydro code.
- So far, two kinds of rotation law profiles are available: uniform and differential, described the j-law profile:

$$\Omega_{c} - \Omega = \frac{1}{A^{2} r_{e}^{2}} \left[ \frac{(\Omega - \omega) r^{2} \sin^{2} \theta e^{-2\nu}}{1 - (\Omega - \omega)^{2} r^{2} \sin^{2} \theta e^{-2\nu}} \right]$$

 Initial data for BNS models are generated externally LORENE, and subsequently imported to the Einstein Toolkit



# Jupyter notebook session

Clone the repository containing useful material such as thornlist files, parameter files, configuration files, EOS tables:

- 1 git clone https://thsoulta@bitbucket.org/thsoulta/et\_thessaloniki.git
  - Download and build the Finstein Toolkit
    - Personal computer
    - Clusters
  - Set up a simulation
    - Isolated rotating (uniformly, differentially) or non-rotating stars with Hydro\_RNSID
    - ▶ Binary neutron star mergers with LORENE









## References



Einstein Toolkit, https://einsteintoolkit.org/



T. W. Baumgarte and S. L. Shapiro, Phys. Rev. D 59, 024007 (1998), gr-qc/9810065.



M. Shibata and T. Nakamura, Phys. Rev. D 52, 5428 (1995).



T. Nakamura, K. Oohara, and Y. Kojima, Prog. Theor. Phys. Suppl. 90, 1 (1987)



Stergioulas N., 1999, Rapidly rotating neutron star, https://www.gravity.phys.uwm.edu/rns



LORENE, http://www.lorene.obspm.fr/



Kranc, http://kranccode.org/



Mösta, P., Mundim, B. C., Faber, J. A., et al. 2014, Classical and Quantum Gravity, 31, 15005.

Introduction to the Einstein Toolkit







