

PROBING THE NATURE OF COMPACT OBJECTS WITH QNMs

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Sebastian H. Völkel


Theoretical Astrophysics
University of Tübingen,
Germany



**3rd HEL.A.S. Summer School
and DAAD School**

8-12 October
Thessaloniki 2018

- 1 COMPACT OBJECTS
- 2 WKB METHOD AND
BOHR-SOMMERFELD RULES
- 3 INVERSE PROBLEM
- 4 RESULTS



PART I

What are Compact Objects?

COMPACT OBJECTS

- Characteristic property is the **compactness** $C \equiv M/R$

COMPACT OBJECTS

- Characteristic property is the **compactness** $C \equiv M/R$
- **Compact objects** require **general relativity**

COMPACT OBJECTS

- Characteristic property is the **compactness** $C \equiv M/R$
- **Compact objects** require **general relativity**
 - **Neutron stars:** $C \approx 0.1 - 0.3$
 - **Black holes:** $C = 0.5$

COMPACT OBJECTS

- Characteristic property is the **compactness** $C \equiv M/R$
- **Compact objects** require **general relativity**
 - **Neutron stars:** $C \approx 0.1 - 0.3$
 - **Black holes:** $C = 0.5$
- Are there **exotic compact objects**?¹

¹Cardoso and Pani (2017)

COMPACT OBJECTS

- Characteristic property is the **compactness** $C \equiv M/R$
- **Compact objects** require **general relativity**
 - **Neutron stars:** $C \approx 0.1 - 0.3$
 - **Black holes:** $C = 0.5$
- Are there **exotic compact objects**?¹
 - gravastars, boson stars, wormholes, alternative black holes, ...
 - objects which could have $C \approx 0.3 - 0.5$

¹Cardoso and Pani (2017)

HOW CAN WE STUDY THEM?

- Traditionally by **electromagnetic (EM) radiation**: radio, x-ray, ...

HOW CAN WE STUDY THEM?

- Traditionally by **electromagnetic (EM) radiation**: radio, x-ray, ...
 - Often complicated source modeling, “**environmental effects**”, ...
 - Black holes are black, only **indirect measurements**

HOW CAN WE STUDY THEM?

- Traditionally by **electromagnetic (EM) radiation**: radio, x-ray, ...
 - Often complicated source modeling, “**environmental effects**”, ...
 - Black holes are black, only **indirect measurements**
- In contrast, **gravitational wave (GW) emission**

HOW CAN WE STUDY THEM?

- Traditionally by **electromagnetic (EM) radiation**: radio, x-ray, ...
 - Often complicated source modeling, “**environmental effects**”, ...
 - Black holes are black, only **indirect measurements**
- In contrast, **gravitational wave (GW) emission**
 - Our **new window** to the universe (**Nobel Prize in Physics 2017**)
 - **Testing compact objects** and **general relativity**

HOW DO WE DESCRIBE GW EMISSION?

- Solving **Einstein's equations** for **dynamical space-times**

HOW DO WE DESCRIBE GW EMISSION?

- Solving **Einstein's equations** for **dynamical space-times**
- **Complicated**, in general different approaches (examples):

HOW DO WE DESCRIBE GW EMISSION?

- Solving **Einstein's equations** for **dynamical space-times**
- **Complicated**, in general different approaches (examples):
 - **Post Newtonian theory** (weak gravity, less compact objects)

HOW DO WE DESCRIBE GW EMISSION?

- Solving **Einstein's equations** for **dynamical space-times**
- **Complicated**, in general different approaches (examples):
 - **Post Newtonian theory** (weak gravity, less compact objects)
 - **Perturbation theory** (weak+strong gravity, but small perturbations)

HOW DO WE DESCRIBE GW EMISSION?

- Solving **Einstein's equations** for **dynamical space-times**
- **Complicated**, in general different approaches (examples):
 - **Post Newtonian theory** (weak gravity, less compact objects)
 - **Perturbation theory** (weak+strong gravity, but small perturbations)
 - **Numerical relativity** (solving the full Einstein equations numerically)

PERTURBATION THEORY

- **Small perturbations** $h_{\mu\nu}$ to a known **background solution** $g_{\mu\nu}^0$

$$g_{\mu\nu} = g_{\mu\nu}^0 + h_{\mu\nu} \quad (1)$$

²Reviews: Kokkotas and Schmidt (1999); Nollert (1999); Berti et al. (2009)
Maggiore (2018)

PERTURBATION THEORY

- **Small perturbations** $h_{\mu\nu}$ to a known **background solution** $g_{\mu\nu}^0$

$$g_{\mu\nu} = g_{\mu\nu}^0 + h_{\mu\nu} \quad (1)$$

- **Spherically symmetric** and **non-rotating objects** (background)

²Reviews: Kokkotas and Schmidt (1999); Nollert (1999); Berti et al. (2009)
Maggiore (2018)

PERTURBATION THEORY

- **Small perturbations** $h_{\mu\nu}$ to a known **background solution** $g_{\mu\nu}^0$

$$g_{\mu\nu} = g_{\mu\nu}^0 + h_{\mu\nu} \quad (1)$$

- **Spherically symmetric** and **non-rotating objects** (background)
- A lengthy calculation yields (for **axial perturbations**)²

$$\frac{d^2}{dr^{*2}} \Psi(r) + [\omega_n^2 - V(r)] \Psi(r) = 0, \quad (2)$$

²Reviews: Kokkotas and Schmidt (1999); Nollert (1999); Berti et al. (2009)
Maggiore (2018)

PERTURBATION THEORY

- **Small perturbations** $h_{\mu\nu}$ to a known **background solution** $g_{\mu\nu}^0$

$$g_{\mu\nu} = g_{\mu\nu}^0 + h_{\mu\nu} \quad (1)$$

- **Spherically symmetric** and **non-rotating objects** (background)
- A lengthy calculation yields (for **axial perturbations**)²

$$\frac{d^2}{dr^{*2}} \Psi(r) + [\omega_n^2 - V(r)] \Psi(r) = 0, \quad (2)$$

- $\Psi(r)$ is related to $h_{\mu\nu}$

²Reviews: Kokkotas and Schmidt (1999); Nollert (1999); Berti et al. (2009)
Maggiore (2018)

PERTURBATION THEORY

- **Small perturbations** $h_{\mu\nu}$ to a known **background solution** $g_{\mu\nu}^0$

$$g_{\mu\nu} = g_{\mu\nu}^0 + h_{\mu\nu} \quad (1)$$

- **Spherically symmetric** and **non-rotating objects** (background)
- A lengthy calculation yields (for **axial perturbations**)²

$$\frac{d^2}{dr^{*2}} \Psi(r) + [\omega_n^2 - V(r)] \Psi(r) = 0, \quad (2)$$

- $\Psi(r)$ is related to $h_{\mu\nu}$
- ω_n^2 are the so-called **quasi-normal modes (QNMs)**

²Reviews: Kokkotas and Schmidt (1999); Nollert (1999); Berti et al. (2009) Maggiore (2018)

PERTURBATION THEORY

- **Small perturbations** $h_{\mu\nu}$ to a known **background solution** $g_{\mu\nu}^0$

$$g_{\mu\nu} = g_{\mu\nu}^0 + h_{\mu\nu} \quad (1)$$

- **Spherically symmetric** and **non-rotating objects** (background)
- A lengthy calculation yields (for **axial perturbations**)²

$$\frac{d^2}{dr^{*2}} \Psi(r) + [\omega_n^2 - V(r)] \Psi(r) = 0, \quad (2)$$

- $\Psi(r)$ is related to $h_{\mu\nu}$
- ω_n^2 are the so-called **quasi-normal modes (QNMs)**
- $V(r)$ acts as **effective potential**

²Reviews: Kokkotas and Schmidt (1999); Nollert (1999); Berti et al. (2009) Maggiore (2018)

PERTURBATION THEORY

- **Small perturbations** $h_{\mu\nu}$ to a known **background solution** $g_{\mu\nu}^0$

$$g_{\mu\nu} = g_{\mu\nu}^0 + h_{\mu\nu} \quad (1)$$

- **Spherically symmetric** and **non-rotating objects** (background)
- A lengthy calculation yields (for **axial perturbations**)²

$$\frac{d^2}{dr^{*2}} \Psi(r) + [\omega_n^2 - V(r)] \Psi(r) = 0, \quad (2)$$

- $\Psi(r)$ is related to $h_{\mu\nu}$
- ω_n^2 are the so-called **quasi-normal modes (QNMs)**
- $V(r)$ acts as **effective potential**
- $r^*(r) \equiv \int^r \sqrt{\frac{g_{11}(r)}{g_{00}(r)}} dr$ **tortoise coordinate**

²Reviews: Kokkotas and Schmidt (1999); Nollert (1999); Berti et al. (2009) Maggiore (2018)

PART II

How can one calculate QNMs?

HOW TO GET THE QNMs?

- **Potential** and **tortoise transformation** are **non-trivial**

HOW TO GET THE QNMs?

- **Potential** and **tortoise transformation** are **non-trivial**
- **Numerical methods** (integrating wave equation, continued fraction, ...)

HOW TO GET THE QNMs?

- **Potential** and **tortoise transformation** are **non-trivial**
- **Numerical methods** (integrating wave equation, continued fraction, ...)
- **Approximate methods** (here WKB)

HOW TO GET THE QNMs?

- **Potential** and **tortoise transformation** are **non-trivial**
- **Numerical methods** (integrating wave equation, continued fraction, ...)
- **Approximate methods** (here WKB)
- Result of **WKB theory** are **Bohr-Sommerfeld (BS) rules**

HOW TO GET THE QNMs?

- **Potential** and **tortoise transformation** are **non-trivial**
- **Numerical methods** (integrating wave equation, continued fraction, ...)
- **Approximate methods** (here WKB)
- Result of **WKB theory** are **Bohr-Sommerfeld (BS) rules**
- **Approximate** but useful for **semi-analytic studies**

THE WKB METHOD I

- Named after **Wentzel**, **Kramer** and **Brillouin** (st. also Jeffreys)

THE WKB METHOD I

- Named after **Wentzel**, **Kramer** and **Brillouin** (st. also Jeffreys)
- Powerful **approximate method** for **linear differential equations**

THE WKB METHOD I

- Named after **Wentzel**, **Kramer** and **Brillouin** (st. also Jeffreys)
- Powerful **approximate method** for **linear differential equations**
- Describes **oscillatory** and **exponentially** “behaving” systems

THE WKB METHOD I

- Named after **Wentzel**, **Kramer** and **Brillouin** (st. also Jeffreys)
- Powerful **approximate method** for **linear differential equations**
- Describes **oscillatory** and **exponentially** “behaving” systems
- **Important** for **quantum mechanics**, but also for **gravitational perturbations**

THE WKB METHOD I

- Named after **Wentzel**, **Kramer** and **Brillouin** (st. also Jeffreys)
- Powerful **approximate method** for **linear differential equations**
- Describes **oscillatory** and **exponentially** “behaving” systems
- **Important** for **quantum mechanics**, but also for **gravitational perturbations**
- **Asymptotic series**, does not necessarily converge!

THE WKB METHOD II

- Time independent **Schrödinger equation** ³

$$\epsilon^2 y''(x) = Q(x)y(x), \quad Q(x) \equiv V(x) - E_n \neq 0 \quad (3)$$

³Details can be found in Bender and Orszag (1978)

THE WKB METHOD II

- Time independent **Schrödinger equation** ³

$$\epsilon^2 y''(x) = Q(x)y(x), \quad Q(x) \equiv V(x) - E_n \neq 0 \quad (3)$$

- **Ansatz** of the form

$$y(x) \sim \exp\left(\frac{1}{\delta} \sum_{n=0}^{\infty} \delta^n S_n(x)\right), \quad \delta \rightarrow 0 \quad (4)$$

³Details can be found in Bender and Orszag (1978)

THE WKB METHOD III

- **Inserting** and **ordering** in powers of ε and δ yields

$$S_0(x) = \pm \int^x \sqrt{Q(t)} dt \quad (5)$$

$$S_1(x) = -\frac{1}{4} \ln(Q(x)) \quad (6)$$

THE WKB METHOD III

- **Inserting** and **ordering** in powers of ε and δ yields

$$S_0(x) = \pm \int^x \sqrt{Q(t)} dt \quad (5)$$

$$S_1(x) = -\frac{1}{4} \ln(Q(x)) \quad (6)$$

- Leading order WKB approximation for $y(x)$

$$y(x) \sim c_1 Q^{-1/4} \exp\left(\frac{1}{\varepsilon} \int_a^x \sqrt{Q(t)} dt\right) \quad (7)$$

$$+ c_2 Q^{-1/4} \exp\left(-\frac{1}{\varepsilon} \int_a^x \sqrt{Q(t)} dt\right), \quad \varepsilon \rightarrow 0 \quad (8)$$

THE WKB METHOD IV

- Valid for **slowly changing potentials** and **not too small** $Q(x)$

$$\left| \frac{d}{dx} \frac{1}{Q(x)} \right| = \left| \frac{Q'(x)}{Q(x)^2} \right| \ll 1 \quad (9)$$

THE WKB METHOD IV

- Valid for **slowly changing potentials** and **not too small** $Q(x)$

$$\left| \frac{d}{dx} \frac{1}{Q(x)} \right| = \left| \frac{Q'(x)}{Q(x)^2} \right| \ll 1 \quad (9)$$

- WKB** solutions are **not valid** at **turning points** ($E_n = V(x)$)!

THE WKB METHOD IV

- Valid for **slowly changing potentials** and **not too small** $Q(x)$

$$\left| \frac{d}{dx} \frac{1}{Q(x)} \right| = \left| \frac{Q'(x)}{Q(x)^2} \right| \ll 1 \quad (9)$$

- WKB** solutions are **not valid** at **turning points** ($E_n = V(x)$)!
- Solve locally** the wave equation and **match** with **neighboring WKB** solutions

THE WKB METHOD IV

- Valid for **slowly changing potentials** and **not too small** $Q(x)$

$$\left| \frac{d}{dx} \frac{1}{Q(x)} \right| = \left| \frac{Q'(x)}{Q(x)^2} \right| \ll 1 \quad (9)$$


- WKB** solutions are **not valid** at **turning points** ($E_n = V(x)$)!
- Solve locally** the wave equation and **match** with **neighboring WKB** solutions
- Use **expanded potential around turning points** (Airy functions for linear expansion, “Kramer’s relations”)

THE BOHR-SOMMERFELD RULES I

- Depending on the type of potential, “shortcuts appear”

THE BOHR-SOMMERFELD RULES I

- Depending on the type of potential, “shortcuts appear”
- Example: **potential well** $V(x)$ with **two turnings points** (x_0, x_1)



THE BOHR-SOMMERFELD RULES II

- **Different regions** exist:

THE BOHR-SOMMERFELD RULES II

- **Different regions** exist:
 - One “**classically allowed**” region: $E_n > V(x)$
 - Two “**classically forbidden**” regions: $E_n < V(x)$
 - Two **regions close to the turning points**: $E_n \approx V(x)$

THE BOHR-SOMMERFELD RULES II

- **Different regions** exist:
 - One “**classically allowed**” region: $E_n > V(x)$
 - Two “**classically forbidden**” regions: $E_n < V(x)$
 - Two **regions close to the turning points**: $E_n \approx V(x)$
- Take **WKB solutions** in **allowed** and **forbidden regions**

THE BOHR-SOMMERFELD RULES II

- **Different regions** exist:
 - One “**classically allowed**” region: $E_n > V(x)$
 - Two “**classically forbidden**” regions: $E_n < V(x)$
 - Two **regions close to the turning points**: $E_n \approx V(x)$
- Take **WKB solutions** in **allowed** and **forbidden regions**
- Get **expanded solution** for regions around **turning points**

THE BOHR-SOMMERFELD RULES II

- **Different regions** exist:
 - One “**classically allowed**” region: $E_n > V(x)$
 - Two “**classically forbidden**” regions: $E_n < V(x)$
 - Two **regions close to the turning points**: $E_n \approx V(x)$
- Take **WKB solutions** in **allowed** and **forbidden regions**
- Get **expanded solution** for regions around **turning points**
- **Matching** solutions yields the classical **Bohr-Sommerfeld rule**

$$\int_{x_0}^{x_1} \sqrt{E_n - V(x)} dx = \pi \left(n + \frac{1}{2} \right), \quad n = 0, 1, 2, \dots \quad (10)$$

GENERALIZED BOHR-SOMMERFELD RULES: I

- Generalization to **quasi-stationary states**⁴

$$\int_{x_0}^{x_1} \sqrt{E_n - V(x)} dx = \pi \left(n + \frac{1}{2} \right) - \frac{i}{4} \exp \left(2i \int_{x_1}^{x_2} \sqrt{E_n - V(x)} dx \right) \quad (11)$$

⁴Popov et al. (1991)

⁵Völkel and Kokkotas (2017,1)

GENERALIZED BOHR-SOMMERFELD RULES: I

- Generalization to **quasi-stationary states**⁴

$$\int_{x_0}^{x_1} \sqrt{E_n - V(x)} dx = \pi \left(n + \frac{1}{2} \right) - \frac{i}{4} \exp \left(2i \int_{x_1}^{x_2} \sqrt{E_n - V(x)} dx \right) \quad (11)$$




FIGURE 1: Völkel and Kokkotas (2017,1).


⁴Popov et al. (1991)

⁵Völkel and Kokkotas (2017,1)

GENERALIZED BOHR-SOMMERFELD RULES: I

- Generalization to **quasi-stationary states**⁴

$$\int_{x_0}^{x_1} \sqrt{E_n - V(x)} dx = \pi \left(n + \frac{1}{2} \right) - \frac{i}{4} \exp \left(2i \int_{x_1}^{x_2} \sqrt{E_n - V(x)} dx \right) \quad (11)$$



- 3 classical turning points
- Complex** spectrum E_n
- Two widths $\mathcal{L}_1, \mathcal{L}_2$
- Application to ultra compact stars⁵

FIGURE 1: Völkel and Kokkotas (2017,1).

⁴Popov et al. (1991)

⁵Völkel and Kokkotas (2017,1)

GENERALIZED BOHR-SOMMERFELD RULES: II

- Estimate for repulsive potentials with a jump and QNM condition⁶

$$\int_{x_0}^R \sqrt{E_n - V(x)} dx = \left(n + \frac{3}{4}\right) + i \arctanh \left(\frac{\sqrt{E_n - V(R_-)}}{\sqrt{E_n - V(R_+)}} \right) \quad (12)$$


⁶Völkel and Kokkotas (2018, TBS)

⁷First derived in Zhang et al. (2011)

GENERALIZED BOHR-SOMMERFELD RULES: II

- Estimate for repulsive potentials with a jump and QNM condition⁶

$$\int_{x_0}^R \sqrt{E_n - V(x)} dx = \left(n + \frac{3}{4}\right) + i \operatorname{arctanh} \left(\frac{\sqrt{E_n - V(R_-)}}{\sqrt{E_n - V(R_+)}} \right) \quad (12)$$




⁶Völkel and Kokkotas (2018, TBS)

⁷First derived in Zhang et al. (2011)

GENERALIZED BOHR-SOMMERFELD RULES: II

- Estimate for repulsive potentials with a jump and QNM condition⁶

$$\int_{x_0}^R \sqrt{E_n - V(x)} dx = \left(n + \frac{3}{4}\right) + i \arctanh \left(\frac{\sqrt{E_n - V(R_-)}}{\sqrt{E_n - V(R_+)}} \right) \quad (12)$$



- 1 classical turning point
- Complex** spectrum
- Asymptotic $\Delta\omega_{r,n} = \pi/R^*$ ⁷
- Applies to **neutron stars**

⁶Völkel and Kokkotas (2018, TBS)

⁷First derived in Zhang et al. (2011)

PART III

*Can one hear the shape of a drum?*⁸

⁸Kac, M. (1966); Gordon et al. (1992)

CAN ONE HEAR THE SHAPE OF A DRUM?

INVERT BS RULES

- Known for **single** wells and **single** barriers (**classical BS rule**)⁹

⁹Wheeler (1976); Chadan and Sabatier (1989); Lazenby and Griffiths (1980); Gandhi and Efthimiou (2006)

¹⁰Völkel and Kokkotas (2017,2); Völkel (2018,1); Völkel and Kokkotas (2018,2)

¹¹Völkel and Kokkotas (2018 TBS)

INVERT BS RULES

- Known for **single** wells and **single** barriers (**classical BS rule**)⁹
- Extended to **quasi-stationary states** (3 or 4 turning points)¹⁰

⁹Wheeler (1976); Chadan and Sabatier (1989); Lazenby and Griffiths (1980); Gandhi and Efthimiou (2006)

¹⁰Völkel and Kokkotas (2017,2); Völkel (2018,1); Völkel and Kokkotas (2018,2)

¹¹Völkel and Kokkotas (2018 TBS)

INVERT BS RULES

- Known for **single** wells and **single** barriers (**classical BS rule**)⁹
- Extended to **quasi-stationary states** (3 or 4 turning points)¹⁰
- **Neutron star potentials** with discontinuity (1 turning point)¹¹

⁹Wheeler (1976); Chadan and Sabatier (1989); Lazenby and Griffiths (1980); Gandhi and Efthimiou (2006)

¹⁰Völkel and Kokkotas (2017,2); Völkel (2018,1); Völkel and Kokkotas (2018,2)

¹¹Völkel and Kokkotas (2018 TBS)

INVERT BS RULES

- Known for **single** wells and **single** barriers (**classical BS rule**)⁹
- Extended to **quasi-stationary states** (3 or 4 turning points)¹⁰
- **Neutron star potentials** with discontinuity (1 turning point)¹¹

$$\mathcal{L}_1(E) = x_1 - x_0 = 2 \frac{\partial}{\partial E} \int_{E_{\min}}^E \frac{n(E') + 1/2}{\sqrt{E - E'}} dE' \quad (13)$$

$$\mathcal{L}_2(E) = x_2 - x_1 = -\frac{1}{\pi} \int_E^{E_{\max}} \frac{(dT(E')/dE')}{T(E')\sqrt{E' - E}} dE' \quad (14)$$

⁹Wheeler (1976); Chadan and Sabatier (1989); Lazenby and Griffiths (1980); Gandhi and Efthimiou (2006)

¹⁰Völkel and Kokkotas (2017,2); Völkel (2018,1); Völkel and Kokkotas (2018,2)


¹¹Völkel and Kokkotas (2018 TBS)

PART IV

Results

ULTRA COMPACT STARS


Example for ultra compact constant density star $C \approx 0.44$



Reconstructed axial perturbation potential, constant density star, $l = 3$, taken from Völkel and Kokkotas (2017,2).

DAMOUR-SOLODUKHIN WORMHOLE


Example for Damour-Solodukhin wormhole $C \approx 0.5$



Reconstructed axial perturbation potential, Damour-Solodukhin wormhole, $l = 3$, taken from Völkel and Kokkotas (2018,2).

NEUTRON STARS


Example for neutron star polytrope $C \approx 0.15$



Reconstructed axial perturbation potential, neutron star polytrope, $l = 3$, Völkel and Kokkotas (2018 TBS).

NEUTRON STARS

Fit spectrum to analytic model potential for neutron star
Bohr-Sommerfeld rule to reconstruct internal space-time



Reconstructed metric function $v(r)$, for two neutron star polytropes, $l = 3$,
Völkel and Kokkotas (2018 TBS).

UNIQUENESS?

- Inverted BS rules give **approximate results**

UNIQUENESS?

- Inverted BS rules give **approximate results**
- Reconstruction of the **potential widths**

UNIQUENESS?

- Inverted BS rules give **approximate results**
- Reconstruction of the **potential widths**
- Usually **no unique potential** possible (expected)

UNIQUENESS?

- Inverted BS rules give **approximate results**
- Reconstruction of the **potential widths**
- Usually **no unique potential** possible (expected)
- **More information** is required for unique reconstruction, e.g.:
 - Birkhoff's theorem (ultra compact stars)
 - Symmetric potential (some wormholes)
 - Radius of the star (normal neutron stars)

UNIQUENESS?

- Inverted BS rules give **approximate results**
- Reconstruction of the **potential widths**
- Usually **no unique potential** possible (expected)
- **More information** is required for unique reconstruction, e.g.:
 - Birkhoff's theorem (ultra compact stars)
 - Symmetric potential (some wormholes)
 - Radius of the star (normal neutron stars)
- **One has to assume the underlying type of potential!**

CONCLUSIONS

- **Perturbations** of simple compact objects are described by the one-dimensional **wave equation**

CONCLUSIONS

- **Perturbations** of simple compact objects are described by the one-dimensional **wave equation**
- Different types of **compact objects** have different **potentials & QNMs**

CONCLUSIONS

- **Perturbations** of simple compact objects are described by the one-dimensional **wave equation**
- Different types of **compact objects** have different **potentials & QNMs**
- **WKB/Bohr-Sommerfeld** rules are useful for semi-analytic work

CONCLUSIONS

- **Perturbations** of simple compact objects are described by the one-dimensional **wave equation**
- Different types of **compact objects** have different **potentials & QNMs**
- **WKB/Bohr-Sommerfeld** rules are useful for semi-analytic work
- Can be inverted for the **inverse spectrum problem**

CONCLUSIONS

- **Perturbations** of simple compact objects are described by the one-dimensional **wave equation**
- Different types of **compact objects** have different **potentials & QNMs**
- **WKB/Bohr-Sommerfeld** rules are useful for semi-analytic work
- Can be inverted for the **inverse spectrum problem**
- **Potential** can be approximately **reconstructed**, depends on the type of object

Details for the presented work can be found in:

- Völkel and Kokkotas, *Class. Quantum Grav.* 34, 125006 (2017)
- Völkel and Kokkotas, *Class. Quantum Grav.* 34, 175015 (2017)
- Maselli, Völkel, and Kokkotas, *Phys. Rev. D* 96, 064045 (2017)
- Völkel, *J. Phys. Commun.* 2, 025029 (2018)
- Völkel and Kokkotas, *Class. Quantum Grav.* 35, 105018 (2018)

BIBLIOGRAPHY I

- C. M. Bender and S. A. Orszag. *Advanced Mathematical Methods for Scientists and Engineers*. New York: McGraw-Hill, 1978.
- E. Berti, V. Cardoso, and A. O. Starinets. TOPICAL REVIEW: Quasinormal modes of black holes and black branes. *Classical and Quantum Gravity*, 26(16):163001, August 2009. doi: 10.1088/0264-9381/26/16/163001.
- Vitor Cardoso and Paolo Pani. Tests for the existence of horizons through gravitational wave echoes. *Nat. Astron.*, 1:586–591, 2017. doi: 10.1038/s41550-017-0225-y.
- K. Chadan and P. C. Sabatier. *Inverse problems in quantum scattering theory*. Texts and Monographs in Physics. Springer-Verlag, New York, second edition, 1989. doi: 10.1007/978-3-642-83317-5.
- S. C. Gandhi and C. J. Efthimiou. Inversion of Gamow's formula and inverse scattering. *Am. J. Phys.*, 74:638–643, July 2006. doi: 10.1119/1.2190683.
- C. Gordon, D. Webb, and S. Wolpert. Isospectral plane domains and surfaces via Riemannian orbifolds. *Inventiones mathematicae*, 110(1), 1992. ISSN 1432-1297. doi: 10.1007/BF01231320. URL <http://dx.doi.org/10.1007/BF01231320>.
- Kac, M. Can One Hear the Shape of a Drum? *The American Mathematical Monthly*, 73(4), 1966. doi: 10.2307/2313748. URL <http://www.jstor.org/stable/2313748>.
- K. D. Kokkotas and B. G. Schmidt. Quasi-Normal Modes of Stars and Black Holes. *Living Reviews in Relativity*, 2:2, December 1999. doi: 10.12942/lrr-1999-2.
- J. C. Lazenby and D. J. Griffiths. Classical inverse scattering in one dimension. *Am. J. Phys.*, 48: 432–436, June 1980. doi: 10.1119/1.11998.

BIBLIOGRAPHY II

- M. Maggiore. *Gravitational Waves: Astrophysics and Cosmology*. Number τ . 2. Oxford University Press, 2018. ISBN 9780198570899. URL <https://books.google.gr/books?id=3ZNODwAAQBAJ>.
- H.-P. Nollert. TOPICAL REVIEW: Quasinormal modes: the characteristic ‘sound’ of black holes and neutron stars. *Classical and Quantum Gravity*, 16:R159–R216, December 1999. doi: 10.1088/0264-9381/16/12/201.
- V. S. Popov, V. D. Mur, and A. V. Sergeev. Quantization rules for quasistationary states. *Physics Letters A*, 157:185–191, July 1991. doi: 10.1016/0375-9601(91)90048-D.
- S. H. Völkel. Inverse spectrum problem for quasi-stationary states. *J. Phys. Commun.*, 2(2): 025029, 2018,1. doi: 10.1088/2399-6528/aaae2. URL <http://stacks.iop.org/2399-6528/2/i=2/a=025029>.
- S. H. Völkel and K. D. Kokkotas. A semi-analytic study of axial perturbations of ultra compact stars. *Class. Quant. Grav.*, 34(12):125006, June 2017,1. doi: 10.1088/1361-6382/aa68cc.
- S. H. Völkel and K. D. Kokkotas. Ultra Compact Stars: Reconstructing the Perturbation Potential. *Class. Quant. Grav.*, 34(17):175015, 2017,2. doi: 10.1088/1361-6382/aa82de.
- S. H. Völkel and K. D. Kokkotas. Wormhole Potentials and Throats from Quasi-Normal Modes. *Class. Quant. Grav.*, 35(10):105018, 2018,2. doi: 10.1088/1361-6382/aabce6.
- J. A. Wheeler. *Studies in Mathematical Physics: Essays in Honor of Valentine Bargmann*. Princeton Series in Physics. Princeton University Press, 1976. ISBN 9780608066288. URL <https://press.princeton.edu/titles/861.html>.
- Y. J. Zhang, J. Wu, and P. T. Leung. High-frequency behavior of w -mode pulsations of compact stars. *Phys. Rev. D*, 83:064012, Mar 2011. doi: 10.1103/PhysRevD.83.064012. URL <https://link.aps.org/doi/10.1103/PhysRevD.83.064012>.