

Gravitational Waves (A General Overview)

Theocharis Apostolatos

National and Kapodistrian University of Athens



October 3, 2018

Outline

- 1 A first course in Gravitational Radiation
- 2 Detection of GW
- 3 Generation of GW

Electromagnetism

- The basic equation of E/M field (connecting the field with its sources)

$$F^{\mu\nu}_{,\nu} = 4\pi J^\mu$$

where the whole information about the field is included in the components of $F^{\mu\nu}$ while J^μ describes the sources of the field.

No interaction with spacetime.

Electromagnetism

- The basic equation of E/M field (connecting the field with its sources)

$$F^{\mu\nu}_{,\nu} = 4\pi J^{\mu}$$

where the whole information about the field is included in the components of $F^{\mu\nu}$ while J^{μ} describes the sources of the field.

The background spacetime is assumed to be fixed (Minkowski).

No interaction with spacetime.

Electromagnetism

- The basic equation of E/M field (connecting the field with its sources)

$$F^{\mu\nu}_{,\nu} = 4\pi J^\mu$$

where the whole information about the field is included in the components of $F^{\mu\nu}$ while J^μ describes the sources of the field.

The background spacetime is assumed to be fixed (Minkowski).

No interaction with spacetime.

- This is a 1st order p.d.e. with respect to F but 2nd order with respect to the real fundamental field A :

$$F^{\mu\nu} = A^{\nu,\mu} - A^{\mu,\nu}.$$

Electromagnetism

- The basic equation of E/M field (connecting the field with its sources)

$$F^{\mu\nu}_{,\nu} = 4\pi J^\mu$$

where the whole information about the field is included in the components of $F^{\mu\nu}$ while J^μ describes the sources of the field.

The background spacetime is assumed to be fixed (Minkowski).

No interaction with spacetime.

- This is a 1st order p.d.e. with respect to F but 2nd order with respect to the real fundamental field A :

$$F^{\mu\nu} = A^{\nu,\mu} - A^{\mu,\nu}.$$

- Given the distribution of sources (currents, non-moving charges) the field $F^{\mu\nu}$ (or A^μ) can be found.

Electromagnetic waves

- Away from the sources the potential field A^μ satisfies the following equation

$$A^{\nu,\mu}_{,\nu} - A^{\mu,\nu}_{,\nu} = 0$$

which is not exactly a wave equation, except if the 1st term is 0. But A^μ is not unambiguously related with a real E/M field.

$$A'^\mu = A^\mu + \Lambda'^\mu$$

offers an alternative description of the same E/M field.

Electromagnetic waves

- Away from the sources the potential field A^μ satisfies the following equation

$$A^{\nu,\mu}_{,\nu} - A^{\mu,\nu}_{,\nu} = 0$$

which is not exactly a wave equation, except if the 1st term is 0. But A^μ is not unambiguously related with a real E/M field.

$$A'^\mu = A^\mu + \Lambda^{,\mu}$$

offers an alternative description of the same E/M field.

- There is some freedom in defining A^μ (gauge). In the Lorentz gauge

$$A^\nu_{,\nu} = 0$$

the propagation equation is a pure wave equation:

$$\Box A^\mu = 0$$

Propagation of Electromagnetic waves

- Electromagnetic waves (disturbances of the field) propagate in flat spacetime:

Propagation of Electromagnetic waves

- Electromagnetic waves (disturbances of the field) propagate in flat spacetime:
 - ① ...With the speed of light.

Propagation of Electromagnetic waves

- Electromagnetic waves (disturbances of the field) propagate in flat spacetime:
 - ① ...With the speed of light.
 - ② ...Without any dispersion.

Propagation of Electromagnetic waves

- Electromagnetic waves (disturbances of the field) propagate in flat spacetime:
 - ① ...With the speed of light.
 - ② ...Without any dispersion.
 - ③ ...The characteristics of the waves are related to the spatial and temporal properties of the source (charges).

Propagation of Electromagnetic waves

- Electromagnetic waves (disturbances of the field) propagate in flat spacetime:
 - ① ...With the speed of light.
 - ② ...Without any dispersion.
 - ③ ...The characteristics of the waves are related to the spatial and temporal properties of the source (charges).
 - ④ ... An isolated source is expected to produce a spherical E/M wave far from the source, and the field will scale like $A \approx 1/r$.

Propagation of Electromagnetic waves

- Electromagnetic waves (disturbances of the field) propagate in flat spacetime:
 - ① ...With the speed of light.
 - ② ...Without any dispersion.
 - ③ ...The characteristics of the waves are related to the spatial and temporal properties of the source (charges).
 - ④ ... An isolated source is expected to produce a spherical E/M wave far from the source, and the field will scale like $A \approx 1/r$.
- Only accelerating charges generate waves. Linearly moving charges generate static fields (E and M).

Gravity

- The basic equation of a gravitational field (connecting the field with its sources)

$$G^{\mu\nu} = \frac{4\pi G}{c^2} T^{\mu\nu}.$$

$G^{\mu\nu}$, the Einstein tensor, comes from the Riemann tensor (describing all the details of spacetime curvature), while $T^{\mu\nu}$ includes all info about the distribution and motion of masses/energy.

Gravity

- The basic equation of a gravitational field (connecting the field with its sources)

$$G^{\mu\nu} = \frac{4\pi G}{c^2} T^{\mu\nu}.$$

$G^{\mu\nu}$, the Einstein tensor, comes from the Riemann tensor (describing all the details of spacetime curvature), while $T^{\mu\nu}$ includes all info about the distribution and motion of masses/energy.

Spacetime is NOT fixed. Its own disturbances is the field itself.

Gravity

- The basic equation of a gravitational field (connecting the field with its sources)

$$G^{\mu\nu} = \frac{4\pi G}{c^2} T^{\mu\nu}.$$

$G^{\mu\nu}$, the Einstein tensor, comes from the Riemann tensor (describing all the details of spacetime curvature), while $T^{\mu\nu}$ includes all info about the distribution and motion of masses/energy.

Spacetime is NOT fixed. Its own disturbances is the field itself.

- This is again a 2nd order p.d.e. with respect to the fundamental field, the metric $g_{\mu\nu}$:

$$G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R_{\kappa}^{\kappa}$$

and

$$R^{\mu\nu} = R^{\mu\alpha\nu\beta} g_{\alpha\beta}.$$

Things to have in mind

- In order to describe the metric we need some coordinates to pinpoint each spacetime point (event). We could choose them freely, thus affecting the metric function, but not the way the spacetime is wrapped.

Things to have in mind

- In order to describe the metric we need some coordinates to pinpoint each spacetime point (event). We could choose them freely, thus affecting the metric function, but not the way the spacetime is wrapped.
- If we want to look for waves we should somehow “discriminate” them from the background. (An ocean wave with the dimensions of the Earth (tidal) is truly a wave?)

Things to have in mind

- In order to describe the metric we need some coordinates to pinpoint each spacetime point (event). We could choose them freely, thus affecting the metric function, but not the way the spacetime is wrapped.
- If we want to look for waves we should somehow “discriminate” them from the background. (An ocean wave with the dimensions of the Earth (tidal) is truly a wave?)
- Waves in a flat background

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$$

with $h_{\mu\nu} \ll 1$. h could be “seen” as a field living on a flat spacetime as long as all equations are kept linearized with respect to h . [Not only h but $h_{,a}$ should be small as well.]

Wave equations for a suitable h

- Einstein equations for $g = \eta + h$:

$$R_{\alpha\beta\mu\nu} = \frac{1}{2}(h_{\alpha\nu,\beta\mu} + h_{\beta\mu,\alpha\nu} - h_{\alpha\mu,\beta\nu} - h_{\beta\nu,\alpha\mu}) \Rightarrow$$

$$R_{\mu\nu} = R^{\alpha}_{\mu\alpha\nu} = \frac{1}{2}(h^{\alpha}_{\mu,\nu\alpha} + h^{\alpha}_{\nu,\mu\alpha} - h^{\alpha}_{\mu\nu,\alpha} - h_{,\mu\nu}) \Rightarrow$$

$$G_{\mu\nu} = \frac{1}{2}(h^{\alpha}_{\mu\alpha,\nu} + h^{\alpha}_{\nu\alpha,\mu} - h^{\alpha}_{\mu\nu,\alpha} - h_{,\mu\nu} \\ - \eta_{\mu\nu}(h^{\alpha\beta}_{\alpha\beta} - h^{\beta}_{,\beta}))$$

But by using a new field alternative, h with inverted trace

$$\bar{h}_{\mu\nu} \equiv h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h,$$

$$G_{\mu\nu} = \frac{1}{2}(-\bar{h}^{\alpha}_{\mu\nu,\alpha} - \eta_{\mu\nu}\bar{h}^{\alpha\beta}_{\alpha\beta} + \bar{h}^{\alpha}_{\mu\alpha,\nu} + \bar{h}^{\alpha}_{\nu\alpha,\mu})$$

Freedom in choosing h

- If only

$$\bar{h}_{\alpha\beta}{}^{\alpha} = 0 \dots$$

then

$$\square \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^2} T_{\mu\nu}$$

and \bar{h} would propagate in vacuum exactly like a wavy E/M field.

Freedom in choosing h

- If only

$$\bar{h}_{\alpha\beta}{}^{\alpha} = 0 \dots$$

then

$$\square \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^2} T_{\mu\nu}$$

and \bar{h} would propagate in vacuum exactly like a wavy E/M field.

- The new requirement reminds the Lorentz gauge ($A_{\mu,\mu} = 0$) but with one more free index. Could this be somehow satisfied?

Freedom in choosing h

- If only

$$\bar{h}_{\alpha\beta,}{}^{\alpha} = 0 \dots$$

then

$$\square \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^2} T_{\mu\nu}$$

and \bar{h} would propagate in vacuum exactly like a wavy E/M field.

- The new requirement reminds the Lorentz gauge ($A_{\mu,}{}^{\mu} = 0$) but with one more free index. Could this be somehow satisfied?
- YES. Just shift the coordinates a bit to redefine the almost flat metric.

$$\begin{aligned} x^{\mu (NEW)} &= x^{\mu (OLD)} + \xi^{\mu} \quad \text{with} \quad |\xi^{\mu}| \ll 1 \\ h_{\mu\nu}^{(NEW)} &= h_{\mu\nu}^{(OLD)} - \xi_{\mu,\nu} - \xi_{\nu,\mu} \end{aligned}$$

The Lorentz gauge

- Adjusting ξ_μ to nullify $\bar{h}_{\mu\nu}{}^\nu$: One has to solve the system

$$\square \xi_\mu = \bar{h}^{(OLD)}{}_{\mu\nu}{}^\nu$$

Of course an additional coordinate shift $\xi \rightarrow \xi + \Xi$, with

$$\square \Xi_\mu = 0$$

is still a Lorentz gauge (no exhaustion of available freedom in defining coords).

The Lorentz gauge

- Adjusting ξ_μ to nullify $\bar{h}_{\mu\nu}{}^\nu$: One has to solve the system

$$\square \xi_\mu = \bar{h}^{(OLD)}{}_{\mu\nu}{}^\nu$$

Of course an additional coordinate shift $\xi \rightarrow \xi + \Xi$, with

$$\square \Xi_\mu = 0$$

is still a Lorentz gauge (no exhaustion of available freedom in defining coords).

- In this type of coords the new metric field \bar{h} satisfies the wave equation

$$\square \bar{h}_{\mu\nu} = 0$$

far from any matter.

The wavy solution

- The wave should have the following form

$$\bar{h}_{\mu\nu} = A_{\mu\nu} e^{ik_\alpha x^\alpha} \quad (1)$$

x^α are the new shifted coords. while k_α should be null
($k_\alpha k^\alpha = 0$).

The wavy solution

- The wave should have the following form

$$\bar{h}_{\mu\nu} = A_{\mu\nu} e^{ik_\alpha x^\alpha} \quad (1)$$

x^α are the new shifted coords. while k_α should be null ($k_\alpha k^\alpha = 0$).

On the other hand $\bar{h}_{\mu\nu}$ should be consistent with Lorentz gauge. Thus

$$A_{\mu\nu} k^\nu = 0.$$

$A_{\mu\nu}$ should be a symmetric 2nd order tensor orthogonal to a null vector.

Counting freedom

- $10-4-4=2$. Analytically:

Counting freedom

- $10-4-4=2$. Analytically:

$A_{\mu\nu}$ should be a symmetric tensor (metric components), that is 10 arbitrary components.

Counting freedom

- $10-4-4=2$. Analytically:

$A_{\mu\nu}$ should be a symmetric tensor (metric components), that is 10 arbitrary components.

$A_{\mu\nu}k^\nu = 0$ with $\mu = 0, 1, 2, 3$ are 4 constraints (from Lorentz gauge).

Counting freedom

- $10-4-4=2$. Analytically:

$A_{\mu\nu}$ should be a symmetric tensor (metric components), that is 10 arbitrary components.

$A_{\mu\nu}k^\nu = 0$ with $\mu = 0, 1, 2, 3$ are 4 constraints (from Lorentz gauge).

On the other hand we could choose as we wish the 4 solutions of Ξ^μ that define the coordinate shift, without affecting the Lorentz gauge.

Counting freedom

- $10-4-4=2$. Analytically:

$A_{\mu\nu}$ should be a symmetric tensor (metric components), that is 10 arbitrary components.

$A_{\mu\nu}k^\nu = 0$ with $\mu = 0, 1, 2, 3$ are 4 constraints (from Lorentz gauge).

On the other hand we could choose as we wish the 4 solutions of Ξ^μ that define the coordinate shift, without affecting the Lorentz gauge.

Therefore only 2 components of $A_{\mu\nu}$ remain to be fixed. These are related to the polarizations of GW.

A wise choice of freedom

- Choose the Ξ solutions so that

$$A_{\mu\nu}u^\nu = 0$$

for a specific 4-velocity (for a specific observer O). This is only 3 constraints since $A_{\mu\nu}k^\nu = 0$ already.

A wise choice of freedom

- Choose the Ξ solutions so that

$$A_{\mu\nu} u^\nu = 0$$

for a specific 4-velocity (for a specific observer O). This is only 3 constraints since $A_{\mu\nu} k^\nu = 0$ already.

- In O's frame ($u^\nu = (1, 0, 0, 0)$)

$$A_{\mu 0} = 0$$

$$A_{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & A_{11} & A_{12} & A_{13} \\ 0 & A_{12} & A_{22} & A_{23} \\ 0 & A_{13} & A_{23} & A_{33} \end{pmatrix}$$

A wise choice of freedom

- Choose the Ξ solutions so that

$$A_{\mu\nu}u^\nu = 0$$

for a specific 4-velocity (for a specific observer O). This is only 3 constraints since $A_{\mu\nu}k^\nu = 0$ already.

- In O's frame ($u^\nu = (1, 0, 0, 0)$)

$$A_{\mu 0} = 0$$

- From $A_{\mu\nu}k^\nu = 0$ (say $k^\nu = k(1, 0, 0, 1)$)

$$A_{\mu 3} = 0$$

$$A_{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & A_{11} & A_{12} & 0 \\ 0 & A_{12} & A_{22} & 0 \\ 0 & A_{13} & A_{23} & 0 \end{pmatrix}$$

A wise choice of freedom

- Choose the Ξ solutions so that

$$A_{\mu\nu}u^\nu = 0$$

for a specific 4-velocity (for a specific observer O). This is only 3 constraints since $A_{\mu\nu}k^\nu = 0$ already.

- In O's frame ($u^\nu = (1, 0, 0, 0)$)

$$A_{\mu 0} = 0$$

- From $A_{\mu\nu}k^\nu = 0$ (say $k^\nu = k(1, 0, 0, 1)$)

$$A_{\mu 3} = 0$$

- Require $A^\mu_\mu = 0$ (the 4th coordinate adjustment).

$$A_{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & A_{11} & A_{12} & 0 \\ 0 & A_{12} & -A_{11} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

The final metric (in this special frame)

$$g_{\mu\nu}^{TT} = \eta_{\mu\nu} + A_{\mu\nu} e^{ik_\mu x^\mu}$$

with 2 independent parameters.

The final metric (in this special frame)

$$g_{\mu\nu}^{TT} = \eta_{\mu\nu} + A_{\mu\nu} e^{ik_\mu x^\mu}$$

with 2 independent parameters.

It is the metric in the TT (**transverse-traceless**) gauge; that is, in a frame of reference where **no deviations from flatness** arise in the direction of wave propagation, **no deviations in time-components of the metric** (nothing new about clocks besides SR) arise, and **no area distortion** show up in the transverse plane (traceless).

The final metric (in this special frame)

$$g_{\mu\nu}^{TT} = \eta_{\mu\nu} + A_{\mu\nu} e^{ik_{\mu}x^{\mu}}$$

with 2 independent parameters.

It is the metric in the TT (**transverse-traceless**) gauge; that is, in a frame of reference where **no deviations from flatness** arise in the direction of wave propagation, **no deviations in time-components of the metric** (nothing new about clocks besides SR) arise, and **no area distortion** show up in the transverse plane (traceless).

- As the wave propagates, nearby points in the plane vertical to propagation approach or deviate from each other in an oscillatory manner. Actually they do **NOT** move. The spacetime background simply inflates and deflates, so the oscillating distance corresponds to proper distance $\int ds = \int \sqrt{g_{\mu\nu} dx^{\mu} dx^{\nu}}$.

The physical interpretation

- In the TT gauge, a particle, initially at rest, when hit by a GW will remain at a constant coordinate position. (The gauge corresponds to the frame of this particle.) The same is true for any other particle (initially at rest).

The physical interpretation

- In the TT gauge, a particle, initially at rest, when hit by a GW will remain at a constant coordinate position. (The gauge corresponds to the frame of this particle.) The same is true for any other particle (initially at rest).

- The proper distance between such particles is

$$s = \int_0^\epsilon \sqrt{|g_{11}|} dx = \int_0^\epsilon \sqrt{1 + A_{11} e^{ik(z-ct)}} dx$$

for particles A, B at $(t, 0, 0, z)$ and at $(t, \epsilon, 0, z)$.

Therefore this distance will oscillate like

$$s \simeq \left[1 + \frac{A_{11}}{2} \right] e^{ik(z-ct)} \epsilon$$

The physical interpretation

- In the TT gauge, a particle, initially at rest, when hit by a GW will remain at a constant coordinate position. (The gauge corresponds to the frame of this particle.) The same is true for any other particle (initially at rest).

- The proper distance between such particles is

$$s = \int_0^\epsilon \sqrt{|g_{11}|} dx = \int_0^\epsilon \sqrt{1 + A_{11} e^{ik(z-ct)}} dx$$

for particles A, B at $(t, 0, 0, z)$ and at $(t, \epsilon, 0, z)$.

Therefore this distance will oscillate like

$$s \simeq \left[1 + \frac{A_{11}}{2} \right] e^{ik(z-ct)} \epsilon$$

- Similar for the cases:

- B: $(t, 0, \epsilon, z)$ (s as above with $A_{11} \rightarrow -A_{11}$),
- B: $(t, \epsilon/\sqrt{2}, \epsilon/\sqrt{2}, z)$ (s as above with $A_{11} \rightarrow A_{12}$)
- B: (t, ϵ, δ, z) ($s \simeq \sqrt{\epsilon^2 + \delta^2} \left[1 + \frac{A_{11}(\epsilon^2 - \delta^2) + 2A_{12}\epsilon\delta}{2(\epsilon^2 + \delta^2)} \right] e^{i\dots}$)

The polarizations

- The components A_{11} and A_{12} are the amplitudes of 2 independent waves propagating with the same k^μ . They correspond to the 2 possible polarizations of a GW (like the polarizations of EM wave).

The polarizations

- The components A_{11} and A_{12} are the amplitudes of 2 independent waves propagating with the same k^μ . They correspond to the 2 possible polarizations of a GW (like the polarizations of EM wave).
- One of them stretches and compresses alternatively the (proper) distances along the x - and the y -axes. This is called the **+polarization**.

The polarizations

- The components A_{11} and A_{12} are the amplitudes of 2 independent waves propagating with the same k^μ . They correspond to the 2 possible polarizations of a GW (like the polarizations of EM wave).
- One of them stretches and compresses alternatively the (proper) distances along the x - and the y -axes. This is called the **+polarization**.
- The other one stretches and compresses alternatively the (proper) distances along the bisectors of the right angle between the x - and y -axes. This is called the **×polarization**.

The polarizations

- The components A_{11} and A_{12} are the amplitudes of 2 independent waves propagating with the same k^μ . They correspond to the 2 possible polarizations of a GW (like the polarizations of EM wave).
- One of them stretches and compresses alternatively the (proper) distances along the x - and the y -axes. This is called the **+polarization**.
- The other one stretches and compresses alternatively the (proper) distances along the bisectors of the right angle between the x - and y -axes. This is called the **×polarization**.
- A wave could have both polarizations (linear combination of them). Both carry information about the geometry of the source.

What to measure?

- Look for oscillations of the proper distance between free particles.

What to measure?

- Look for oscillations of the proper distance between free particles.
- They should be proportional to the distance between the particles: $\epsilon \rightarrow 2\epsilon$ and $\delta \rightarrow 2\delta$ makes $\Delta s \rightarrow 2\Delta s$.

What to measure?

- Look for oscillations of the proper distance between free particles.
- They should be proportional to the distance between the particles: $\epsilon \rightarrow 2\epsilon$ and $\delta \rightarrow 2\delta$ makes $\Delta s \rightarrow 2\Delta s$.
- Is there a tool to measure proper distance? YES, light. The proper distance for a photon is 0 (null). Thus it could reveal metric disturbances by measuring travel time of light back and forth between two free particles.

$$0 = -c^2 dt^2 + g_{11} dx^2 \Rightarrow \epsilon = \int_0^{T_{\rightarrow}} \frac{cdt}{1 + \frac{A_{11}}{2} \cos[k(z - ct)]}$$

Note: If the distance is small compared to λ_{GW} ($\ll 10^6\text{m}$) ct is almost constant along the flight.

What to measure?

- Look for oscillations of the proper distance between free particles.
- They should be proportional to the distance between the particles: $\epsilon \rightarrow 2\epsilon$ and $\delta \rightarrow 2\delta$ makes $\Delta s \rightarrow 2\Delta s$.
- Is there a tool to measure proper distance? YES, light. The proper distance for a photon is 0 (null). Thus it could reveal metric disturbances by measuring travel time of light back and forth between two free particles.

$$0 = -c^2 dt^2 + g_{11} dx^2 \Rightarrow \epsilon = \int_0^{T_{\rightarrow}} \frac{cdt}{1 + \frac{A_{11}}{2} \cos[k(z - ct)]}$$

Note: If the distance is small compared to λ_{GW} ($\ll 10^6\text{m}$) ct is almost constant along the flight.

- We could measure T_{\rightleftharpoons} and monitor that time.

Interferometric detectors

- Instead of measuring very short time intervals we could make light measure light. (Use interference techniques to measure differences of flight times along different paths.)

Interferometric detectors

- Instead of measuring very short time intervals we could make light measure light. (Use interference techniques to measure differences of flight times along different paths.)

Remember that proper distances along the x -axis and along the y -axis follow opposite behavior (stretch-compression) in $+$ polarization.

Interferometric detectors

- Instead of measuring very short time intervals we could make light measure light. (Use interference techniques to measure differences of flight times along different paths.)

Remember that proper distances along the x -axis and along the y -axis follow opposite behavior (stretch-compression) in $+$ polarization.

- When the two beams interfere, if initially (before the wave) they were adjusted to interfere destructively, they will have partial positive interference when the 2 paths change (when GW present). The light output will be proportional to the small shift, thus to the amplitude of the wave and the length of the beams.

Other detector schemes

- Resonators: Solid bars with specific oscillatory modes.

Other detector schemes

- Resonators: Solid bars with specific oscillatory modes.
- When hit by a GW it will be excited to resonate.

Other detector schemes

- Resonators: Solid bars with specific oscillatory modes.
- When hit by a GW it will be excited to resonate.
- The lower the internal friction the more pronounced the resonance curve. [Low temperature, expensive solid state materials]. One measures its oscillations electronically.

Other detector schemes

- Resonators: Solid bars with specific oscillatory modes.
- When hit by a GW it will be excited to resonate.
- The lower the internal friction the more pronounced the resonance curve. [Low temperature, expensive solid state materials]. One measures its oscillations electronically.
- The disadvantage: Good for only one frequency.

Other detector schemes

- Resonators: Solid bars with specific oscillatory modes.
- When hit by a GW it will be excited to resonate.
- The lower the internal friction the more pronounced the resonance curve. [Low temperature, expensive solid state materials]. One measures its oscillations electronically.
- The disadvantage: Good for only one frequency.
- One could create a xylophone with some of them, so as to resonate one after the other in case of frequency shifting source.

Other detector schemes

- Resonators: Solid bars with specific oscillatory modes.
- When hit by a GW it will be excited to resonate.
- The lower the internal friction the more pronounced the resonance curve. [Low temperature, expensive solid state materials]. One measures its oscillations electronically.
- The disadvantage: Good for only one frequency.
- One could create a xylophone with some of them, so as to resonate one after the other in case of frequency shifting source.

The interferometers are broad-band detectors.

Back to equations; now with matter



$$\square \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^2} T_{\mu\nu}$$

Back to equations; now with matter

- $$\square \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^2} T_{\mu\nu}$$
- For an isolated source (e.g., a binary of compact objects) we expect a spherical wave solution characterized by the frequency of T -oscillations, Ω . Thus

$$\bar{h}_{\mu\nu} = \frac{B_{\mu\nu}}{r} e^{i\Omega(t-r/c)}$$

outside the source.

Back to equations; now with matter

$$\square \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^2} T_{\mu\nu}$$

- For an isolated source (e.g., a binary of compact objects) we expect a spherical wave solution characterized by the frequency of T -oscillations, Ω . Thus

$$\bar{h}_{\mu\nu} = \frac{B_{\mu\nu}}{r} e^{i\Omega(t-r/c)}$$

outside the source.

- A bit complicated calculation (even for low density-slowly moving objects) relates the amplitude $B_{\mu\nu}$ with the 2nd time derivative of the quadrupole moment of the source:

$$B_{jk} e^{i\Omega t} = 2 \frac{d^2}{dt^2} I_{jk}(t)$$

where

$$I_{jk} = \int T^{00} x^j x^k d^3x$$

is the quadrupole moment of the source.

Energy radiated

- The amplitude of strong sources

Estimation of strong energy emitters



Estimation of amplitude at Earth

